

# Valerie A Odero-Marah

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/4903750/publications.pdf>

Version: 2024-02-01

26  
papers

1,739  
citations

394421

19  
h-index

552781

26  
g-index

26  
all docs

26  
docs citations

26  
times ranked

3034  
citing authors

#	ARTICLE	IF	CITATIONS
1	Val16A SOD2 Polymorphism Promotes Epithelial-Mesenchymal Transition Antagonized by Muscadine Grape Skin Extract in Prostate Cancer Cells. <i>Antioxidants</i> , 2021, 10, 213.	5.1	2
2	Proteomics-Metabolomics Combined Approach Identifies Peroxidase as a Protector against Metabolic and Oxidative Stress in Prostate Cancer. <i>International Journal of Molecular Sciences</i> , 2019, 20, 3046.	4.1	32
3	Larrea tridentata Extract Mitigates Oxidative Stress-Induced Cytotoxicity in Human Neuroblastoma SH-SY5Y Cells. <i>Antioxidants</i> , 2019, 8, 427.	5.1	15
4	Cancer-bone microenvironmental interactions promotes STAT3 signaling. <i>Molecular Carcinogenesis</i> , 2019, 58, 1349-1361.	2.7	3
5	CCAAT-displacement protein/cut homeobox transcription factor (CUX1) represses estrogen receptor-alpha (ER- $\alpha$ ) in triple-negative breast cancer cells and can be antagonized by muscadine grape skin extract (MSKE). <i>PLoS ONE</i> , 2019, 14, e0214844.	2.5	8
6	Epithelial-Mesenchymal Transition (EMT) and Prostate Cancer. <i>Advances in Experimental Medicine and Biology</i> , 2018, 1095, 101-110.	1.6	122
7	Association of Epithelial Mesenchymal Transition with prostate and breast health disparities. <i>PLoS ONE</i> , 2018, 13, e0203855.	2.5	7
8	High mobility group A2 (HMGA2) promotes EMT via MAPK pathway in prostate cancer. <i>Biochemical and Biophysical Research Communications</i> , 2018, 504, 196-202.	2.1	48
9	Targeting the Nuclear Cathepsin L CCAAT Displacement Protein/Cut Homeobox Transcription Factor-Epithelial Mesenchymal Transition Pathway in Prostate and Breast Cancer Cells with the Z-FY-CHO Inhibitor. <i>Molecular and Cellular Biology</i> , 2017, 37, .	2.3	26
10	Snail transcription factor NLS and importin $\beta$ 1 regulate the subcellular localization of Cathepsin L and Cux1. <i>Biochemical and Biophysical Research Communications</i> , 2017, 491, 59-64.	2.1	14
11	Muscadine Grape Skin Extract Induces an Unfolded Protein Response-Mediated Autophagy in Prostate Cancer Cells: A TMT-Based Quantitative Proteomic Analysis. <i>PLoS ONE</i> , 2016, 11, e0164115.	2.5	31
12	Snail promotes cell migration through PI3K/AKT-dependent Rac1 activation as well as PI3K/AKT-independent pathways during prostate cancer progression. <i>Cell Adhesion and Migration</i> , 2015, 9, 255-264.	2.7	58
13	The impact of low-dose carcinogens and environmental disruptors on tissue invasion and metastasis. <i>Carcinogenesis</i> , 2015, 36, S128-S159.	2.8	40
14	Assessing the carcinogenic potential of low-dose exposures to chemical mixtures in the environment: the challenge ahead. <i>Carcinogenesis</i> , 2015, 36, S254-S296.	2.8	239
15	Muscadine grape skin extract can antagonize Snail-cathepsin L-mediated invasion, migration and osteoclastogenesis in prostate and breast cancer cells. <i>Carcinogenesis</i> , 2015, 36, 1019-1027.	2.8	48
16	Snail Promotes Epithelial Mesenchymal Transition in Breast Cancer Cells in Part via Activation of Nuclear ERK2. <i>PLoS ONE</i> , 2014, 9, e104987.	2.5	94
17	Camalexin-Induced Apoptosis in Prostate Cancer Cells Involves Alterations of Expression and Activity of Lysosomal Protease Cathepsin D. <i>Molecules</i> , 2014, 19, 3988-4005.	3.8	21
18	Muscadine grape skin extract reverts snail-mediated epithelial mesenchymal transition via superoxide species in human prostate cancer cells. <i>BMC Complementary and Alternative Medicine</i> , 2014, 14, 97.	3.7	22

#	ARTICLE	IF	CITATIONS
19	The phytoalexin camalexin mediates cytotoxicity towards aggressive prostate cancer cells via reactive oxygen species. <i>Journal of Natural Medicines</i> , 2013, 67, 607-618.	2.3	16
20	The role of Snail in prostate cancer. <i>Cell Adhesion and Migration</i> , 2012, 6, 433-441.	2.7	99
21	Snail-mediated regulation of reactive oxygen species in ARCaP human prostate cancer cells. <i>Biochemical and Biophysical Research Communications</i> , 2011, 404, 34-39.	2.1	61
22	Snail negatively regulates cell adhesion to extracellular matrix and integrin expression via the MAPK pathway in prostate cancer cells. <i>Cell Adhesion and Migration</i> , 2011, 5, 249-257.	2.7	41
23	Snail transcription factor regulates neuroendocrine differentiation in LNCaP prostate cancer cells. <i>Prostate</i> , 2010, 70, 982-992.	2.3	86
24	Epithelial to mesenchymal transition (EMT) in human prostate cancer: lessons learned from ARCaP model. <i>Clinical and Experimental Metastasis</i> , 2008, 25, 601-610.	3.3	147
25	Receptor activator of NF- $\kappa$ B Ligand (RANKL) expression is associated with epithelial to mesenchymal transition in human prostate cancer cells. <i>Cell Research</i> , 2008, 18, 858-870.	12.0	123
26	Insulin-like Growth Factor-1-Dependent Up-regulation of ZEB1 Drives Epithelial-to-Mesenchymal Transition in Human Prostate Cancer Cells. <i>Cancer Research</i> , 2008, 68, 2479-2488.	0.9	336