

Nicola J Wiles

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/4909224/publications.pdf>

Version: 2024-02-01

85
papers

3,947
citations

172457

29
h-index

133252

59
g-index

91
all docs

91
docs citations

91
times ranked

5809
citing authors

#	ARTICLE	IF	CITATIONS
1	Therapist-delivered internet psychotherapy for depression in primary care: a randomised controlled trial. <i>Lancet</i> , The, 2009, 374, 628-634.	13.7	340
2	Cognitive behavioural therapy as an adjunct to pharmacotherapy for primary care based patients with treatment resistant depression: results of the CoBaIT randomised controlled trial. <i>Lancet</i> , The, 2013, 381, 375-384.	13.7	278
3	Self-reported psychotic symptoms in the general population. <i>British Journal of Psychiatry</i> , 2006, 188, 519-526.	2.8	214
4	Insomnia, worry, anxiety and depression as predictors of the occurrence and persistence of paranoid thinking. <i>Social Psychiatry and Psychiatric Epidemiology</i> , 2012, 47, 1195-1203.	3.1	190
5	Facilitated physical activity as a treatment for depressed adults: randomised controlled trial. <i>BMJ</i> , The, 2012, 344, e2758-e2758.	6.0	163
6	User Experience of Cognitive Behavioral Therapy Apps for Depression: An Analysis of App Functionality and User Reviews. <i>Journal of Medical Internet Research</i> , 2018, 20, e10120.	4.3	147
7	Validity of the Short Mood and Feelings Questionnaire in late adolescence.. <i>Psychological Assessment</i> , 2014, 26, 752-762.	1.5	137
8	Birth weight and psychological distress at age 45â€“51 years. <i>British Journal of Psychiatry</i> , 2005, 187, 21-28.	2.8	118
9	Prevalence of treatment-resistant depression in primary care: cross-sectional data. <i>British Journal of General Practice</i> , 2013, 63, e852-e858.	1.4	115
10	Early Parental and Child Predictors of Recurrent Abdominal Pain at School Age: Results of a Large Population-Based Study. <i>Journal of the American Academy of Child and Adolescent Psychiatry</i> , 2006, 45, 729-736.	0.5	110
11	Is socioeconomic status in early life associated with drug use? A systematic review of the evidence. <i>Drug and Alcohol Review</i> , 2009, 28, 142-153.	2.1	110
12	Cost-effectiveness of therapist-delivered online cognitiveâ€“behavioural therapy for depression: randomised controlled trial. <i>British Journal of Psychiatry</i> , 2010, 197, 297-304.	2.8	103
13	Long-term effectiveness and cost-effectiveness of cognitive behavioural therapy as an adjunct to pharmacotherapy for treatment-resistant depression in primary care: follow-up of the CoBaIT randomised controlled trial. <i>Lancet Psychiatry</i> , the, 2016, 3, 137-144.	7.4	103
14	Fetal Growth and Childhood Behavioral Problems: Results from the ALSPAC Cohort. <i>American Journal of Epidemiology</i> , 2006, 163, 829-837.	3.4	95
15	Is screen time associated with anxiety or depression in young people? Results from a UK birth cohort. <i>BMC Public Health</i> , 2019, 19, 82.	2.9	81
16	The clinical effectiveness of sertraline in primary care and the role of depression severity and duration (PANDA): a pragmatic, double-blind, placebo-controlled randomised trial. <i>Lancet Psychiatry</i> , the, 2019, 6, 903-914.	7.4	78
17	Maintenance or Discontinuation of Antidepressants in Primary Care. <i>New England Journal of Medicine</i> , 2021, 385, 1257-1267.	27.0	75
18	Physical activity and emotional problems amongst adolescents. <i>Social Psychiatry and Psychiatric Epidemiology</i> , 2008, 43, 765-772.	3.1	74

#	ARTICLE	IF	CITATIONS
19	The process and delivery of CBT for depression in adults: a systematic review and network meta-analysis. <i>Psychological Medicine</i> , 2019, 49, 1937-1947.	4.5	70
20	Socioeconomic status in childhood and later alcohol use: a systematic review*. <i>Addiction</i> , 2007, 102, 1546-1563.	3.3	69
21	Clinical effectiveness and cost-effectiveness of cognitive behavioural therapy as an adjunct to pharmacotherapy for treatment-resistant depression in primary care: the CoBaT randomised controlled trial. <i>Health Technology Assessment</i> , 2014, 18, 1-167, vii-viii.	2.8	69
22	Physical activity and depression in adolescents: cross-sectional findings from the ALSPAC cohort. <i>Social Psychiatry and Psychiatric Epidemiology</i> , 2012, 47, 1023-1033.	3.1	62
23	Physical Activity and Common Mental Disorder: Results from the Caerphilly Study. <i>American Journal of Epidemiology</i> , 2007, 165, 946-954.	3.4	55
24	Polymorphism of the 5-HT transporter and response to antidepressants: randomised controlled trial. <i>British Journal of Psychiatry</i> , 2011, 198, 464-471.	2.8	55
25	Adverse effects from antidepressant treatment: randomised controlled trial of 601 depressed individuals. <i>Psychopharmacology</i> , 2014, 231, 2921-2931.	3.1	55
26	Psychological therapies for treatment-resistant depression in adults. <i>The Cochrane Library</i> , 2018, 2018, CD010558.	2.8	54
27	How much change is enough? Evidence from a longitudinal study on depression in UK primary care. <i>Psychological Medicine</i> , 2022, 52, 1875-1882.	4.5	51
28	No pain, no gain: Depressed clients' experiences of cognitive behavioural therapy. <i>British Journal of Clinical Psychology</i> , 2013, 52, 347-364.	3.5	48
29	Mirtazapine added to SSRIs or SNRIs for treatment resistant depression in primary care: phase III randomised placebo controlled trial (MIR). <i>BMJ: British Medical Journal</i> , 2018, 363, k4218.	2.3	44
30	Usefulness of PHQ-9 in primary care to determine meaningful symptoms of low mood: a qualitative study. <i>British Journal of General Practice</i> , 2016, 66, e78-e84.	1.4	33
31	Cognitive behavioural therapy as an adjunct to pharmacotherapy for treatment resistant depression in primary care: The CoBaT randomised controlled trial protocol. <i>Contemporary Clinical Trials</i> , 2012, 33, 312-319.	1.8	32
32	Exploring patients' reasons for declining contact in a cognitive behavioural therapy randomised controlled trial in primary care. <i>British Journal of General Practice</i> , 2012, 62, e371-e377.	1.4	30
33	Pharmacological interventions for treatment-resistant depression in adults. <i>The Cochrane Library</i> , 2019, 12, CD010557.	2.8	30
34	Longitudinal Associations between Adolescent Psychotic Experiences and Depressive Symptoms. <i>PLoS ONE</i> , 2014, 9, e105758.	2.5	30
35	Socioeconomic Indicators of Treatment Prognosis for Adults With Depression. <i>JAMA Psychiatry</i> , 2022, 79, 406.	11.0	30
36	Effective dose 50 method as the minimal clinically important difference: Evidence from depression trials. <i>Journal of Clinical Epidemiology</i> , 2021, 137, 200-208.	5.0	29

#	ARTICLE	IF	CITATIONS
37	A comparison of four different approaches to measuring health utility in depressed patients. <i>Health and Quality of Life Outcomes</i> , 2013, 11, 81.	2.4	28
38	The contribution of depressive disorder characteristics™ to determinations of prognosis for adults with depression: an individual patient data meta-analysis. <i>Psychological Medicine</i> , 2021, 51, 1068-1081.	4.5	28
39	Detecting Heterogeneity of Intervention Effects Using Analysis and Meta-analysis of Differences in Variance Between Trial Arms. <i>Epidemiology</i> , 2021, 32, 846-854.	2.7	28
40	Cost-effectiveness of cognitive behavioural therapy as an adjunct to pharmacotherapy for treatment-resistant depression in primary care: economic evaluation of the CoBaIT Trial. <i>British Journal of Psychiatry</i> , 2014, 204, 69-76.	2.8	27
41	A Randomized Controlled Trial of Cognitive Behavioural Therapy as an Adjunct to Pharmacotherapy in Primary Care Based Patients with Treatment Resistant Depression: A Pilot Study. <i>Behavioural and Cognitive Psychotherapy</i> , 2008, 36, 21-33.	1.2	24
42	GENetic and clinical Predictors Of treatment response in Depression: the GenPod randomised trial protocol. <i>Trials</i> , 2008, 9, 29.	1.6	23
43	Severity of depression and response to antidepressants: GENPOD randomised controlled trial. <i>British Journal of Psychiatry</i> , 2012, 200, 130-136.	2.8	23
44	Management of treatment-resistant depression in primary care: a mixed-methods study. <i>British Journal of General Practice</i> , 2018, 68, e673-e681.	1.4	23
45	Combining mirtazapine with SSRIs or SNRIs for treatment-resistant depression: the MIR RCT. <i>Health Technology Assessment</i> , 2018, 22, 1-136.	2.8	21
46	The feasibility of low-intensity psychological therapy for depression co-occurring with autism in adults: The Autism Depression Trial (ADEPT) – a pilot randomised controlled trial. <i>Autism</i> , 2020, 24, 1360-1372.	4.1	18
47	Is social support pre-treatment associated with prognosis for adults with depression in primary care?. <i>Acta Psychiatrica Scandinavica</i> , 2021, 143, 392-405.	4.5	17
48	Moderators of response to cognitive behavioural therapy as an adjunct to pharmacotherapy for treatment-resistant depression in primary care. <i>Journal of Affective Disorders</i> , 2015, 174, 272-280.	4.1	16
49	Why do GPs exclude patients from participating in research? An exploration of adherence to and divergence from trial criteria. <i>Family Practice</i> , 2014, 31, 364-370.	1.9	15
50	A randomised controlled trial assessing the severity and duration of depressive symptoms associated with a clinically significant response to sertraline versus placebo, in people presenting to primary care with depression (PANDA trial): study protocol for a randomised controlled trial. <i>Trials</i> , 2017, 18, 496.	1.6	15
51	Design Considerations for the Integrated Delivery of Cognitive Behavioral Therapy for Depression: User-Centered Design Study. <i>JMIR Mental Health</i> , 2020, 7, e15972.	3.3	15
52	Trends in the recording of anxiety in UK primary care: a multi-method approach. <i>Social Psychiatry and Psychiatric Epidemiology</i> , 2022, 57, 375-386.	3.1	13
53	Integrating the Digital and the Traditional to Deliver Therapy for Depression. , 2020, , .		13
54	Protocol for a feasibility study and randomised pilot trial of a low-intensity psychological intervention for depression in adults with autism: the Autism Depression Trial (ADEPT). <i>BMJ Open</i> , 2017, 7, e019545.	1.9	12

#	ARTICLE	IF	CITATIONS
55	Variation in recognition of happy and sad facial expressions and self-reported depressive symptom severity: A prospective cohort study. <i>Journal of Affective Disorders</i> , 2019, 257, 461-469.	4.1	12
56	A randomised controlled trial assessing the use of citalopram, sertraline, fluoxetine and mirtazapine in preventing relapse in primary care patients who are taking long-term maintenance antidepressants (ANTLER: ANTidepressants to prevent reLapse in dEpRession): study protocol for a randomised controlled trial. <i>Trials</i> , 2019, 20, 319.	1.6	12
57	Personality difficulties and response to community-based psychological treatment for anxiety and depression. <i>Journal of Affective Disorders</i> , 2021, 279, 266-273.	4.1	12
58	Guided self-help for depression in autistic adults: the ADEPT feasibility RCT. <i>Health Technology Assessment</i> , 2019, 23, 1-94.	2.8	12
59	Individuals'™ Long Term Use of Cognitive Behavioural Skills to Manage their Depression: A Qualitative Study. <i>Behavioural and Cognitive Psychotherapy</i> , 2017, 45, 46-57.	1.2	11
60	Reference-based multiple imputation for missing data sensitivity analyses in trial-based cost-effectiveness analysis. <i>Health Economics (United Kingdom)</i> , 2020, 29, 171-184.	1.7	11
61	Patients'™ experiences of participating in a large-scale trial of cognitive behavioural therapy for depression: a mixed methods study. <i>Family Practice</i> , 2013, 30, 705-711.	1.9	10
62	Mirtazapine added to selective serotonin reuptake inhibitors for treatment-resistant depression in primary care (MIR trial): study protocol for a randomised controlled trial. <i>Trials</i> , 2016, 17, 66.	1.6	10
63	A Patient Stratification Approach to Identifying the Likelihood of Continued Chronic Depression and Relapse Following Treatment for Depression. <i>Journal of Personalized Medicine</i> , 2021, 11, 1295.	2.5	10
64	Antidepressant medication to prevent depression relapse in primary care: the ANTLER RCT. <i>Health Technology Assessment</i> , 2021, 25, 1-62.	2.8	9
65	GPs'™ views and experiences of managing patients with personality disorder: a qualitative interview study. <i>BMJ Open</i> , 2019, 9, e026616.	1.9	8
66	An expert consensus on the most effective components of cognitive behavioural therapy for adults with depression: a modified Delphi study. <i>Cognitive Behaviour Therapy</i> , 2020, 49, 242-255.	3.5	8
67	PTSD as a mediator of the relationship between trauma and psychotic experiences. <i>Psychological Medicine</i> , 2022, 52, 2722-2730.	4.5	8
68	Identifying patients at risk of psychosis: a qualitative study of GP views in South West England. <i>British Journal of General Practice</i> , 2021, 71, e113-e120.	1.4	8
69	Cost-Effectiveness of Different Formats for Delivery of Cognitive Behavioral Therapy for Depression: A Systematic Review Based Economic Model. <i>Value in Health</i> , 2020, 23, 1662-1670.	0.3	7
70	Life events and treatment prognosis for depression: A systematic review and individual patient data meta-analysis. <i>Journal of Affective Disorders</i> , 2022, 299, 298-308.	4.1	7
71	Rise in prescribing for anxiety in UK primary care between 2003 and 2018: a population-based cohort study using Clinical Practice Research Datalink. <i>British Journal of General Practice</i> , 2022, 72, e511-e518.	1.4	7
72	Materials used to support cognitive behavioural therapy for depression: a survey of therapists'™ clinical practice and views. <i>Cognitive Behaviour Therapy</i> , 2019, 48, 463-481.	3.5	6

#	ARTICLE	IF	CITATIONS
73	GPs™ and patients™ views on the value of diagnosing anxiety disorders in primary care: a qualitative interview study. <i>British Journal of General Practice</i> , 2021, 71, e450-e457.	1.4	6
74	Antidepressant treatment with sertraline for adults with depressive symptoms in primary care: the PANDA research programme including RCT. <i>Programme Grants for Applied Research</i> , 2019, 7, 1-108.	1.0	6
75	Cost-Utility Analysis of Discontinuing Antidepressants in England Primary Care Patients Compared with Long-Term Maintenance: The ANTLER Study. <i>Applied Health Economics and Health Policy</i> , 2022, 20, 269-282.	2.1	6
76	Cognitive behavioural therapy for treatment-resistant depression “ Authors' reply. <i>Lancet</i> , The, 2013, 381, 1814-1815.	13.7	4
77	Sex differences in depressive symptoms and tolerability after treatment with selective serotonin reuptake inhibitor antidepressants: Secondary analyses of the GENPOD trial. <i>Journal of Psychopharmacology</i> , 2021, 35, 919-927.	4.0	4
78	Pharmacological interventions for treatment-resistant depression in adults. <i>The Cochrane Library</i> , 0, ,.	2.8	3
79	Cost-Effectiveness of Sertraline in Primary Care According to Initial Severity and Duration of Depressive Symptoms: Findings from the PANDA RCT. <i>Pharmacoeconomics - Open</i> , 2020, 4, 427-438.	1.8	3
80	Sensitivity to missing not at random dropout in clinical trials: Use and interpretation of the trimmed means estimator. <i>Statistics in Medicine</i> , 2022, 41, 1462-1481.	1.6	3
81	Are multiple physical symptoms a poor prognostic factor or just a marker of depression severity? Secondary analysis of the GenPod trial. <i>Journal of Affective Disorders</i> , 2014, 163, 40-46.	4.1	2
82	Feasibility study of eye movement desensitisation and reprocessing (EMDR) in people with an at-risk mental state (ARMS) for psychosis: study protocol. <i>BMJ Open</i> , 2020, 10, e038620.	1.9	2
83	Does anxiety moderate the effectiveness of mirtazapine in patients with treatment-resistant depression? A secondary analysis of the MIR trial. <i>Journal of Psychopharmacology</i> , 2020, 34, 1342-1349.	4.0	2
84	Republished research: Facilitated physical activity as a treatment for depressed adults: randomised controlled trial. <i>British Journal of Sports Medicine</i> , 2013, 47, 629-629.	6.7	1
85	Conditional Beliefs of Primary-Care Patients with Treatment-Resistant Depression. <i>Behavioural and Cognitive Psychotherapy</i> , 2016, 44, 513-526.	1.2	0