

# Anne-Catherine Prats

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/4924736/publications.pdf>

Version: 2024-02-01

81  
papers

5,804  
citations

71102

41  
h-index

76900

74  
g-index

90  
all docs

90  
docs citations

90  
times ranked

5618  
citing authors

#	ARTICLE	IF	CITATIONS
1	High Level of Staufen1 Expression Confers Longer Recurrence Free Survival to Non-Small Cell Lung Cancer Patients by Promoting THBS1 mRNA Degradation. <i>International Journal of Molecular Sciences</i> , 2022, 23, 215.	4.1	8
2	Pre-Conditioning Methods and Novel Approaches with Mesenchymal Stem Cells Therapy in Cardiovascular Disease. <i>Cells</i> , 2022, 11, 1620.	4.1	17
3	Sex Hormones in Lymphedema. <i>Cancers</i> , 2021, 13, 530.	3.7	11
4	Coordinating Effect of VEGFC and Oleic Acid Participates to Tumor Lymphangiogenesis. <i>Cancers</i> , 2021, 13, 2851.	3.7	4
5	Circular RNA, the Key for Translation. <i>International Journal of Molecular Sciences</i> , 2020, 21, 8591.	4.1	92
6	How are circRNAs translated by non-canonical initiation mechanisms?. <i>Biochimie</i> , 2019, 164, 45-52.	2.6	50
7	IRES Trans-Acting Factors, Key Actors of the Stress Response. <i>International Journal of Molecular Sciences</i> , 2019, 20, 924.	4.1	112
8	Paclitaxel induces lymphatic endothelial cells autophagy to promote metastasis. <i>Cell Death and Disease</i> , 2019, 10, 956.	6.3	32
9	Periprostatic Adipose Tissue Favors Prostate Cancer Cell Invasion in an Obesity-Dependent Manner: Role of Oxidative Stress. <i>Molecular Cancer Research</i> , 2019, 17, 821-835.	3.4	76
10	Vasohibin1, a new mouse cardiomyocyte IRES trans-acting factor that regulates translation in early hypoxia. <i>ELife</i> , 2019, 8, .	6.0	19
11	Lymphatic Vasculature Requires Estrogen Receptor- $\beta$ Signaling to Protect From Lymphedema. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2018, 38, 1346-1357.	2.4	47
12	Activated human primary NK cells efficiently kill colorectal cancer cells in 3D spheroid cultures irrespectively of the level of PD-L1 expression. <i>Oncolmmunology</i> , 2018, 7, e1395123.	4.6	37
13	Therapeutic Benefit and Gene Network Regulation by Combined Gene Transfer of Apelin, FGF2, and SERCA2a into Ischemic Heart. <i>Molecular Therapy</i> , 2018, 26, 902-916.	8.2	20
14	Local production of tenascin-C acts as a trigger for monocyte/macrophage recruitment that provokes cardiac dysfunction. <i>Cardiovascular Research</i> , 2018, 114, 123-137.	3.8	38
15	Therapeutic Benefits and Adverse Effects of Combined Proangiogenic Gene Therapy in Mouse Critical Leg Ischemia. <i>Annals of Vascular Surgery</i> , 2017, 40, 252-261.	0.9	12
16	Dachsous1- $\beta$ Fat4 Signaling Controls Endothelial Cell Polarization During Lymphatic Valve Morphogenesis- $\beta$ Brief Report. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2017, 37, 1732-1735.	2.4	31
17	Apelin modulates pathological remodeling of lymphatic endothelium after myocardial infarction. <i>JCI Insight</i> , 2017, 2, .	5.0	68
18	Nucleolin Promotes Heat Shock-Associated Translation of VEGF-D to Promote Tumor Lymphangiogenesis. <i>Cancer Research</i> , 2016, 76, 4394-4405.	0.9	26

#	ARTICLE	IF	CITATIONS
19	Promoter-Dependent Translation Controlled by p54nrb and hnRNPM during Myoblast Differentiation. PLoS ONE, 2015, 10, e0136466.	2.5	19
20	Role of hypoxia and vascular endothelial growth factors in lymphangiogenesis. Molecular and Cellular Oncology, 2015, 2, e1024821.	0.7	41
21	Internal ribosome entry site-based vectors for combined gene therapy. World Journal of Experimental Medicine, 2015, 5, 11.	1.7	26
22	Role of hypoxia and vascular endothelial growth factors in lymphangiogenesis. Molecular and Cellular Oncology, 2014, 1, e29907.	0.7	33
23	Hypothalamic Apelin/Reactive Oxygen Species Signaling Controls Hepatic Glucose Metabolism in the Onset of Diabetes. Antioxidants and Redox Signaling, 2014, 20, 557-573.	5.4	44
24	Hypoxia Induces VEGF-C Expression in Metastatic Tumor Cells via a HIF-1 $\alpha$ -Independent Translation-Mediated Mechanism. Cell Reports, 2014, 6, 155-167.	6.4	102
25	The p53 isoform, $\Delta 133p53$ , stimulates angiogenesis and tumour progression. Oncogene, 2013, 32, 2150-2160.	5.9	75
26	p53 Acts as a Safeguard of Translational Control by Regulating Fibrillar and rRNA Methylation in Cancer. Cancer Cell, 2013, 24, 318-330.	16.8	246
27	CXCL4L1 $\alpha$ fibroblast cooperation inhibits tumor angiogenesis, lymphangiogenesis and metastasis. Microvascular Research, 2013, 89, 25-33.	2.5	29
28	p53 mutant breast cancer patients expressing $\Delta 133p53$ have as good a prognosis as wild-type p53 breast cancer patients. Breast Cancer Research, 2011, 13, R7.	5.0	92
29	p53 directly transactivates $\Delta 133p53$ , regulating cell fate outcome in response to DNA damage. Cell Death and Differentiation, 2011, 18, 248-258.	11.2	103
30	Abstract 3482: Hypoxia induces translational regulation of lymphangiogenic growth factors. , 2011, , .		0
31	R22: Rôle de l'hypoxie dans la régulation traductionnelle des facteurs de croissance lymphangiogéniques lors du développement tumoral. Bulletin Du Cancer, 2010, 97, S24.	1.6	0
32	D133P53, directly transactivated by p53, prevents p53-mediated apoptosis without inhibiting p53-mediated cell cycle arrest. Breast Cancer Research, 2010, 12, .	5.0	1
33	Dysregulation of Ribosome Biogenesis and Translational Capacity Is Associated with Tumor Progression of Human Breast Cancer Cells. PLoS ONE, 2009, 4, e7147.	2.5	198
34	Fibroblast growth factor 1 induced during myogenesis by a transcription $\alpha$ translation coupling mechanism. Nucleic Acids Research, 2009, 37, 5267-5278.	14.5	31
35	IRES-based Vector Coexpressing FGF2 and Cyr61 Provides Synergistic and Safe Therapeutics of Lower Limb Ischemia. Molecular Therapy, 2009, 17, 2010-2019.	8.2	21
36	Low dietary inorganic phosphate affects the brain by controlling apoptosis, cell cycle and protein translation. Journal of Nutritional Biochemistry, 2008, 19, 16-25.	4.2	15

#	ARTICLE	IF	CITATIONS
37	Efficient gene transfer in skeletal muscle with AAV-derived bicistronic vector using the FGF-1 IRES. <i>Gene Therapy</i> , 2008, 15, 1090-1098.	4.5	12
38	Potent activation of FGF-2 IRES-dependent mechanism of translation during brain development. <i>Rna</i> , 2008, 14, 1852-1864.	3.5	25
39	FGF2 Translationally Induced by Hypoxia Is Involved in Negative and Positive Feedback Loops with HIF-1 $\pm$ . <i>PLoS ONE</i> , 2008, 3, e3078.	2.5	65
40	Translational Induction of VEGF Internal Ribosome Entry Site Elements During the Early Response to Ischemic Stress. <i>Circulation Research</i> , 2007, 100, 305-308.	4.5	64
41	Long term expression of bicistronic vector driven by the FGF-1 IRES in mouse muscle. <i>BMC Biotechnology</i> , 2007, 7, 74.	3.3	17
42	The use of IRES-based bicistronic vectors allows the stable expression of recombinant G-protein coupled receptors such as NPY5 and histamine 4. <i>Biochimie</i> , 2006, 88, 737-746.	2.6	11
43	IRES-dependent regulation of FGF-2 mRNA translation in pathophysiological conditions in the mouse. <i>Biochemical Society Transactions</i> , 2006, 34, 17-21.	3.4	19
44	A High Inorganic Phosphate Diet Perturbs Brain Growth, Alters Akt-ERK Signaling, and Results in Changes in Cap-Dependent Translation. <i>Toxicological Sciences</i> , 2006, 90, 221-229.	3.1	30
45	Testosterone regulates FGF $\alpha$ 2 expression during testis maturation by an IRES-dependent translational mechanism. <i>FASEB Journal</i> , 2006, 20, 476-478.	0.5	49
46	Heterogeneous Nuclear Ribonucleoprotein A1 Is a Novel Internal Ribosome Entry Site trans-Acting Factor That Modulates Alternative Initiation of Translation of the Fibroblast Growth Factor 2 mRNA. <i>Journal of Biological Chemistry</i> , 2005, 280, 4144-4153.	3.4	134
47	Hyperglycemia upregulates translation of the fibroblast growth factor 2 mRNA in mouse aorta via internal ribosome entry site. <i>FASEB Journal</i> , 2004, 18, 1583-1585.	0.5	27
48	Antiangiogenic Properties of Fibstatin, an Extracellular FGF-2-binding Polypeptide. <i>Cancer Research</i> , 2004, 64, 7507-7512.	0.9	47
49	Role of Fibroblast Growth Factor-2 Isoforms in the Effect of Estradiol on Endothelial Cell Migration and Proliferation. <i>Circulation Research</i> , 2004, 94, 1301-1309.	4.5	56
50	Internal Ribosome Entry Site Structural Motifs Conserved among Mammalian Fibroblast Growth Factor 1 Alternatively Spliced mRNAs. <i>Molecular and Cellular Biology</i> , 2004, 24, 7622-7635.	2.3	60
51	Estrogens and atherosclerosis. <i>European Journal of Endocrinology</i> , 2004, 150, 113-117.	3.7	42
52	Use and comparison of different internal ribosomal entry sites (IRES) in tricistronic retroviral vectors. <i>BMC Biotechnology</i> , 2004, 4, 16.	3.3	56
53	HIGH MW FGF2, BUT NOT LOW MW FGF2 NOR VEGF, MEDIATES THE EFFECT OF ESTRADIOL ON REENDOTHELIALIZATION. <i>Journal of Hypertension</i> , 2004, 22, S189.	0.5	0
54	Generation of protein isoform diversity by alternative initiation of translation at non-AUG codons. <i>Biology of the Cell</i> , 2003, 95, 169-178.	2.0	220

#	ARTICLE	IF	CITATIONS
55	Unr Is Required In Vivo for Efficient Initiation of Translation from the Internal Ribosome Entry Sites of both Rhinovirus and Poliovirus. <i>Journal of Virology</i> , 2003, 77, 3353-3359.	3.4	106
56	A Single Internal Ribosome Entry Site Containing a G Quartet RNA Structure Drives Fibroblast Growth Factor 2 Gene Expression at Four Alternative Translation Initiation Codons. <i>Journal of Biological Chemistry</i> , 2003, 278, 39330-39336.	3.4	151
57	Translational control of gene expression: Role of IRESs and consequences for cell transformation and angiogenesis. <i>Progress in Molecular Biology and Translational Science</i> , 2002, 72, 367-413.	1.9	51
58	Pyrimidine tract binding protein and La autoantigen interact differently with the 5' untranslated regions of lentiviruses and oncoretrovirus mRNAs. <i>FEBS Letters</i> , 2001, 490, 54-58.	2.8	23
59	Tumour suppressor p53 inhibits human fibroblast growth factor 2 expression by a post-transcriptional mechanism. <i>Oncogene</i> , 2001, 20, 1669-1677.	5.9	48
60	Translation of the human c-myc PO tricistronic mRNA involves two independent internal ribosome entry sites. <i>Oncogene</i> , 2001, 20, 4270-4280.	5.9	31
61	p53 directs conformational change and translation initiation blockade of human fibroblast growth factor 2 mRNA. <i>Oncogene</i> , 2001, 20, 4613-4620.	5.9	47
62	c- myc Internal Ribosome Entry Site Activity Is Developmentally Controlled and Subjected to a Strong Translational Repression in Adult Transgenic Mice. <i>Molecular and Cellular Biology</i> , 2001, 21, 1833-1840.	2.3	89
63	New Ways of Initiating Translation in Eukaryotes?. <i>Molecular and Cellular Biology</i> , 2001, 21, 8238-8246.	2.3	60
64	Fibroblast Growth Factor 2 Internal Ribosome Entry Site (Ires) Activity Ex Vivo and in Transgenic Mice Reveals a Stringent Tissue-Specific Regulation. <i>Journal of Cell Biology</i> , 2000, 150, 275-281.	5.2	138
65	Alternative Translation Initiation of Human Fibroblast Growth Factor 2 mRNA Controlled by Its 3'-Untranslated Region Involves a Poly(A) Switch and a Translational Enhancer. <i>Journal of Biological Chemistry</i> , 2000, 275, 19361-19367.	3.4	34
66	Expression of Human Fibroblast Growth Factor 2 mRNA Is Post-transcriptionally Controlled by a Unique Destabilizing Element Present in the 3'-Untranslated Region between Alternative Polyadenylation Sites. <i>Journal of Biological Chemistry</i> , 1999, 274, 21402-21408.	3.4	68
67	A New 34-Kilodalton Isoform of Human Fibroblast Growth Factor 2 Is Cap Dependently Synthesized by Using a Non-AUG Start Codon and Behaves as a Survival Factor. <i>Molecular and Cellular Biology</i> , 1999, 19, 505-514.	2.3	200
68	FGF-2 : la traduction bat son plein.. <i>Medecine/Sciences</i> , 1999, 15, 905.	0.2	0
69	Understanding the Translation Regulatory Mechanisms to Improve the Efficiency and the Specificity of Protein Production by the Cell Factory. <i>Cell Engineering</i> , 1999, , 1-37.	0.4	0
70	Two Independent Internal Ribosome Entry Sites Are Involved in Translation Initiation of Vascular Endothelial Growth Factor mRNA. <i>Molecular and Cellular Biology</i> , 1998, 18, 6178-6190.	2.3	276
71	Alternative Translation of the Proto-oncogene c-myc by an Internal Ribosome Entry Site. <i>Journal of Biological Chemistry</i> , 1997, 272, 32061-32066.	3.4	219
72	Translation of CUG- but not AUG-initiated forms of human fibroblast growth factor 2 is activated in transformed and stressed cells.. <i>Journal of Cell Biology</i> , 1996, 135, 1391-1402.	5.2	146

#	ARTICLE	IF	CITATIONS
73	Alternative Translation of Human Fibroblast Growth Factor 2 mRNA Occurs by Internal Entry of Ribosomes. <i>Molecular and Cellular Biology</i> , 1995, 15, 35-44.	2.3	327
74	Alternative Translation Initiation of the Moloney Murine Leukemia Virus mRNA Controlled by Internal Ribosome Entry Involving the p57/PTB Splicing Factor. <i>Journal of Biological Chemistry</i> , 1995, 270, 20376-20383.	3.4	108
75	cis-acting elements involved in the alternative translation initiation process of human basic fibroblast growth factor mRNA.. <i>Molecular and Cellular Biology</i> , 1992, 12, 4796-4805.	2.3	75
76	Viral RNA annealing activities of the nucleocapsid protein of Moloney murine leukemia virus are zinc independent. <i>Nucleic Acids Research</i> , 1991, 19, 3533-3541.	14.5	95
77	CUG initiation codon used for the synthesis of a cell surface antigen coded by the murine leukemia virus. <i>Journal of Molecular Biology</i> , 1989, 205, 363-372.	4.2	170
78	High molecular mass forms of basic fibroblast growth factor are initiated by alternative CUG codons.. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1989, 86, 1836-1840.	7.1	444
79	Replication of pSC101: effects of mutations in the E. coli DNA binding protein IHF. <i>Molecular Genetics and Genomics</i> , 1986, 204, 85-89.	2.4	107
80	A plasmid vector allowing positive selection of recombinant plasmids in <i>Streptococcus pneumoniae</i> . <i>Gene</i> , 1985, 39, 41-48.	2.2	21
81	Hyperrecombination at a specific DNA sequence in pneumococcal transformation.. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1984, 81, 5184-5188.	7.1	20