Flavia Restitutti

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/4927602/publications.pdf

Version: 2024-02-01

933447 940533 16 388 10 16 citations h-index g-index papers 16 16 16 210 docs citations times ranked citing authors all docs

#	Article	IF	CITATIONS
1	Investigation of the effects of vatinoxan on somatic and visceral antinociceptive efficacy of medetomidine in dogs. American Journal of Veterinary Research, 2020, 81, 299-308.	0.6	4
2	Cardiovascular effects of intravenous vatinoxan (MK-467) in medetomidine–tiletamine–zolazepam anaesthetised red deer (Cervus elaphus). Veterinary Anaesthesia and Analgesia, 2020, 47, 518-527.	0.6	10
3	Cardiovascular and sedation reversal effects of intramuscular administration of atipamezole in dogs treated with medetomidine hydrochloride with or without the peripheral $\hat{l}\pm 2$ -adrenoceptor antagonist vatinoxan hydrochloride. American Journal of Veterinary Research, 2019, 80, 912-922.	0.6	6
4	Plasma concentration and cardiovascular effects of intramuscular medetomidine combined with three doses of the peripheral alpha2-antagonist MK-467 in dogs. Veterinary Anaesthesia and Analgesia, 2017, 44, 417-426.	0.6	23
5	Evaluation of a transcutaneous method to assess canine liver function by indocyanine green plasma disappearance rate in healthy adult Beagle dogs. Veterinary Journal, 2016, 209, 169-173.	1.7	6
6	Dexmedetomidine or medetomidine: which should veterinary surgeons select?. Companion Animal, 2016, 21, 128-137.	0.2	3
7	Effect of aquapuncture on postoperative analgesia after ovariohysterectomy in dogs. Semina:Ciencias Agrarias, 2015, 36, 1979.	0.3	5
8	A comparison in dogs of medetomidine, with or without MK-467, and the combination acepromazine-butorphanol as premedication prior to anaesthesia induced by propofol and maintained with isoflurane. Veterinary Anaesthesia and Analgesia, 2014, 41, 163-173.	0.6	14
9	The cardiopulmonary effects of a peripheral alpha-2-adrenoceptor antagonist, MK-467, in dogs sedated with a combination of medetomidine and butorphanol. Veterinary Anaesthesia and Analgesia, 2014, 41, 567-574.	0.6	18
10	Thermographic imaging of superficial temperature in dogs sedated with medetomidine and butorphanol with and without MK-467 (L-659'066). Veterinary Anaesthesia and Analgesia, 2013, 40, 142-148.	0.6	30
11	Effect of MKâ€467 on organ blood flow parameters detected by contrastâ€enhanced ultrasound in dogs treated with dexmedetomidine. Veterinary Anaesthesia and Analgesia, 2013, 40, e48-e56.	0.6	27
12	Influence of MK-467, a Peripherally Acting $\hat{l}\pm <$ sub>2-Adrenoceptor Antagonist on the Disposition of Intravenous Dexmedetomidine in Dogs. Drug Metabolism and Disposition, 2012, 40, 445-449.	3.3	49
13	Plasma glucose, insulin, free fatty acids, lactate and cortisol concentrations in dexmedetomidine-sedated dogs with or without MK-467: A peripheral α-2 adrenoceptor antagonist. Veterinary Journal, 2012, 193, 481-485.	1.7	49
14	Effect of intraperitoneal or incisional bupivacaine on pain and the analgesic requirement after ovariohysterectomy in dogs. Veterinary Anaesthesia and Analgesia, 2012, 39, 426-430.	0.6	41
15	Effects of different doses of L-659'066 on the bispectral index and clinical sedation in dogs treated with dexmedetomidine. Veterinary Anaesthesia and Analgesia, 2011, 38, 415-422.	0.6	35
16	The effects of increasing doses of MKâ€467, a peripheral alpha ₂ â€adrenergic receptor antagonist, on the cardiopulmonary effects of intravenous dexmedetomidine in conscious dogs. Journal of Veterinary Pharmacology and Therapeutics, 2011, 34, 332-337.	1.3	68