

# Hui Lin

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/5003642/publications.pdf>

Version: 2024-02-01

39  
papers

2,426  
citations

257450

24  
h-index

302126

39  
g-index

39  
all docs

39  
docs citations

39  
times ranked

2685  
citing authors

#	ARTICLE	IF	CITATIONS
1	Plants Mitigate Nitrous Oxide Emissions from Antibiotic-Contaminated Agricultural Soils. <i>Environmental Science &amp; Technology</i> , 2022, 56, 4950-4960.	10.0	18
2	Copper exposure effects on antibiotic degradation in swine manure vary between mesophilic and thermophilic conditions. <i>Science of the Total Environment</i> , 2022, 841, 156759.	8.0	3
3	Efficient degradation of tylosin by <i>Klebsiella oxytoca</i> TYL-T1. <i>Science of the Total Environment</i> , 2022, 847, 157305.	8.0	2
4	Effect of composting on the conjugative transmission of sulfonamide resistance and sulfonamide-resistant bacterial population. <i>Journal of Cleaner Production</i> , 2021, 285, 125483.	9.3	17
5	Degradation of tetracycline antibiotics by <i>Arthrobacter nicotianae</i> OTC-16. <i>Journal of Hazardous Materials</i> , 2021, 403, 123996.	12.4	71
6	Microplastics are a hotspot for antibiotic resistance genes: Progress and perspective. <i>Science of the Total Environment</i> , 2021, 773, 145643.	8.0	130
7	Genomic insights into the antibiotic resistance pattern of the tetracycline-degrading bacterium, <i>Arthrobacter nicotianae</i> OTC-16. <i>Scientific Reports</i> , 2021, 11, 15638.	3.3	4
8	Simultaneous reductions in antibiotics and heavy metal pollution during manure composting. <i>Science of the Total Environment</i> , 2021, 788, 147830.	8.0	33
9	Soil microbial activity and community composition as influenced by application of pig biogas slurry in paddy field in southeast China. <i>Paddy and Water Environment</i> , 2020, 18, 15-25.	1.8	12
10	Acidic conditions enhance the removal of sulfonamide antibiotics and antibiotic resistance determinants in swine manure. <i>Environmental Pollution</i> , 2020, 263, 114439.	7.5	33
11	Metagenomic insights into the abundance and composition of resistance genes in aquatic environments: Influence of stratification and geography. <i>Environment International</i> , 2019, 127, 371-380.	10.0	98
12	Compost-bulking agents reduce the reservoir of antibiotics and antibiotic resistance genes in manures by modifying bacterial microbiota. <i>Science of the Total Environment</i> , 2019, 649, 396-404.	8.0	96
13	Plastics in the marine environment are reservoirs for antibiotic and metal resistance genes. <i>Environment International</i> , 2019, 123, 79-86.	10.0	305
14	Fate of tetracycline and sulfonamide resistance genes in a grassland soil amended with different organic fertilizers. <i>Ecotoxicology and Environmental Safety</i> , 2019, 170, 39-46.	6.0	38
15	Antibiotics and antibiotic resistance genes in global lakes: A review and meta-analysis. <i>Environment International</i> , 2018, 116, 60-73.	10.0	474
16	Agro-industrial waste recycling by <i>Trichosporon fermentans</i> : conversion of waste sweetpotato vines alone into lipid. <i>Environmental Science and Pollution Research</i> , 2018, 25, 8793-8799.	5.3	9
17	Antibiotic resistance genes in surface water of eutrophic urban lakes are related to heavy metals, antibiotics, lake morphology and anthropic impact. <i>Ecotoxicology</i> , 2017, 26, 831-840.	2.4	126
18	Effect of temperature on sulfonamide antibiotics degradation, and on antibiotic resistance determinants and hosts in animal manures. <i>Science of the Total Environment</i> , 2017, 607-608, 725-732.	8.0	82

#	ARTICLE	IF	CITATIONS
19	A compositional shift in the soil microbiome induced by tetracycline, sulfamonomethoxine and ciprofloxacin entering a plant-soil system. <i>Environmental Pollution</i> , 2016, 212, 440-448.	7.5	71
20	Occurrence of trace elements and antibiotics in manure-based fertilizers from the Zhejiang Province of China. <i>Science of the Total Environment</i> , 2016, 559, 174-181.	8.0	109
21	The binding, synergistic and structural characteristics of BsEXLX1 for loosening the main components of lignocellulose: Lignin, xylan, and cellulose. <i>Enzyme and Microbial Technology</i> , 2016, 92, 67-75.	3.2	6
22	Comparative genome analysis of the oleaginous yeast <i>Trichosporon fermentans</i> reveals its potential applications in lipid accumulation. <i>Microbiological Research</i> , 2016, 192, 203-210.	5.3	15
23	Antibiotics and Antibiotic Resistance Genes in Sediment of Honghu Lake and East Dongting Lake, China. <i>Microbial Ecology</i> , 2016, 72, 791-801.	2.8	73
24	Effects of manure and mineral fertilization strategies on soil antibiotic resistance gene levels and microbial community in a paddy-upland rotation system. <i>Environmental Pollution</i> , 2016, 211, 332-337.	7.5	80
25	Variations in the fate and biological effects of sulfamethoxazole, norfloxacin and doxycycline in different vegetable-soil systems following manure application. <i>Journal of Hazardous Materials</i> , 2016, 304, 49-57.	12.4	78
26	Characterization of Cellulase Secretion and Cre1-Mediated Carbon Source Repression in the Potential Lignocellulose-Degrading Strain <i>Trichoderma asperellum</i> T-1. <i>PLoS ONE</i> , 2015, 10, e0119237.	2.5	10
27	Sweetpotato vines hydrolysate promotes single cell oils production of <i>Trichosporon fermentans</i> in high-density molasses fermentation. <i>Bioresource Technology</i> , 2015, 176, 249-256.	9.6	26
28	Preparation of a new-style composite containing a key bioflocculant produced by <i>Pseudomonas aeruginosa</i> ZJU1 and its flocculating effect on harmful algal blooms. <i>Journal of Hazardous Materials</i> , 2015, 284, 215-221.	12.4	53
29	Wheat Bran Enhances the Cytotoxicity of Immobilized <i>Alcaligenes aquatilis</i> F8 against <i>Microcystis aeruginosa</i> . <i>PLoS ONE</i> , 2015, 10, e0136429.	2.5	16
30	Soil microbial systems respond differentially to tetracycline, sulfamonomethoxine, and ciprofloxacin entering soil under pot experimental conditions alone and in combination. <i>Environmental Science and Pollution Research</i> , 2014, 21, 7436-7448.	5.3	55
31	Engineering <i>Aspergillus oryzae</i> A-4 through the Chromosomal Insertion of Foreign Cellulase Expression Cassette to Improve Conversion of Cellulosic Biomass into Lipids. <i>PLoS ONE</i> , 2014, 9, e108442.	2.5	6
32	Sweetpotato vines hydrolysate induces glycerol to be an effective substrate for lipid production of <i>Trichosporon fermentans</i> . <i>Bioresource Technology</i> , 2013, 136, 725-729.	9.6	20
33	Mechanism for the disparity of the lipid production by <i>Trichosporon fermentans</i> grown on different sweetpotato vines hydrolysates. <i>Industrial Crops and Products</i> , 2013, 50, 844-851.	5.2	4
34	Potential utilization of waste sweetpotato vines hydrolysate as a new source for single cell oils production by <i>Trichosporon fermentans</i> . <i>Bioresource Technology</i> , 2013, 135, 622-629.	9.6	41
35	Genetic engineering of microorganisms for biodiesel production. <i>Bioengineered</i> , 2013, 4, 292-304.	3.2	41
36	Evaluation of Bacterial Expansin EXLX1 as a Cellulase Synergist for the Saccharification of Lignocellulosic Agro-Industrial Wastes. <i>PLoS ONE</i> , 2013, 8, e75022.	2.5	25

#	ARTICLE	IF	CITATIONS
37	Artificial construction and characterization of a fungal consortium that produces cellulolytic enzyme system with strong wheat straw saccharification. <i>Bioresource Technology</i> , 2011, 102, 10569-10576.	9.6	29
38	Isolation and characterization of <i>Rhodococcus</i> sp. NB5 capable of degrading a high concentration of nitrobenzene. <i>Journal of Basic Microbiology</i> , 2011, 51, 397-403.	3.3	5
39	Direct microbial conversion of wheat straw into lipid by a cellulolytic fungus of <i>Aspergillus oryzae</i> A-4 in solid-state fermentation. <i>Bioresource Technology</i> , 2010, 101, 7556-7562.	9.6	112