

Adam A Garsa

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/5005712/publications.pdf>

Version: 2024-02-01

16
papers

475
citations

933447

10
h-index

996975

15
g-index

16
all docs

16
docs citations

16
times ranked

860
citing authors

#	ARTICLE	IF	CITATIONS
1	Association of CYP2C8, CYP3A4, CYP3A5, and ABCB1 Polymorphisms with the Pharmacokinetics of Paclitaxel. <i>Clinical Cancer Research</i> , 2005, 11, 8097-8104.	7.0	170
2	A multi-institutional comparison of outcomes of immunosuppressed and immunocompetent patients treated with surgery and radiation therapy for cutaneous squamous cell carcinoma of the head and neck. <i>Cancer</i> , 2017, 123, 2054-2060.	4.1	115
3	Prognostic value of 18F-FDG PET metabolic parameters in oropharyngeal squamous cell carcinoma. <i>Journal of Radiation Oncology</i> , 2013, 2, 27-34.	0.7	30
4	Analysis of fat necrosis after adjuvant high-dose-rate interstitial brachytherapy for early stage breast cancer. <i>Brachytherapy</i> , 2013, 12, 99-106.	0.5	30
5	Predictors of Individual Tumor Local Control After Stereotactic Radiosurgery for Non-Small Cell Lung Cancer Brain Metastases. <i>International Journal of Radiation Oncology Biology Physics</i> , 2014, 90, 407-413.	0.8	27
6	Internal dose escalation is associated with increased local control for non-small cell lung cancer (NSCLC) brain metastases treated with stereotactic radiosurgery (SRS). <i>Advances in Radiation Oncology</i> , 2018, 3, 146-153.	1.2	22
7	Management of Chordoma and Chondrosarcoma with Fractionated Stereotactic Radiotherapy. <i>Frontiers in Surgery</i> , 2017, 4, 35.	1.4	20
8	Radiation Therapy for Brain Metastases: A Systematic Review. <i>Practical Radiation Oncology</i> , 2021, 11, 354-365.	2.1	18
9	Pretreatment metabolic tumor volume as a prognostic factor in HPV-associated oropharyngeal cancer in the context of AJCC 8th edition staging. <i>Head and Neck</i> , 2018, 40, 2280-2287.	2.0	14
10	Squamous cell carcinoma of unknown primary of the head and neck: Favorable prognostic factors comparable to those in oropharyngeal cancer. <i>Head and Neck</i> , 2018, 40, 904-916.	2.0	12
11	Influence of respiratory motion management technique on radiation pneumonitis risk with robotic stereotactic body radiation therapy. <i>Journal of Applied Clinical Medical Physics</i> , 2018, 19, 48-57.	1.9	7
12	Use of Salvage Surgery or Stereotactic Radiosurgery for Multiply Recurrent Skull Base Chordomas: A Single-Institution Experience and Review of the Literature. <i>Journal of Neurological Surgery, Part B: Skull Base</i> , 2021, 82, 161-174.	0.8	4
13	Does radiation dose matter in thyroid cancer?: Patterns of local-regional failure in recurrent and metastatic well-differentiated thyroid cancers treated with dose-painted intensity-modulated radiation therapy. <i>Journal of Medical Imaging and Radiation Oncology</i> , 2016, 60, 560-567.	1.8	3
14	Bevacizumab is more effective in nasopharyngeal carcinoma patients with lower maximum radiation dose to the temporal lobe. <i>Chinese Clinical Oncology</i> , 2019, 8, S20-S20.	1.2	2
15	Challenges in the re-irradiation of locally advanced head and neck cancers: outcomes and toxicities. <i>Journal of Radiation Oncology</i> , 2019, 8, 259-266.	0.7	1
16	Slant board immobilisation of head-and-neck radiotherapy patients who cannot tolerate a flat position. <i>Journal of Radiotherapy in Practice</i> , 2016, 15, 303-308.	0.5	0