

# Shuji Murakami

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/50120/publications.pdf>

Version: 2024-02-01

61  
papers

9,335  
citations

430874

18  
h-index

175258

52  
g-index

61  
all docs

61  
docs citations

61  
times ranked

9803  
citing authors

#	ARTICLE	IF	CITATIONS
1	Uncommon EGFR mutations conducted with osimertinib in patients with NSCLC: a study protocol of phase 2 study (UNICORN/TCOG1901). <i>Future Oncology</i> , 2022, 18, 523-531.	2.4	7
2	Predictive Markers for Immune Checkpoint Inhibitors in Non-Small Cell Lung Cancer. <i>Journal of Clinical Medicine</i> , 2022, 11, 1855.	2.4	11
3	A phase II study of durvalumab (MEDI4736) immediately after completion of chemoradiotherapy in unresectable stage III non-small cell lung cancer: TORG1937 (DATE study). <i>Journal of Clinical Oncology</i> , 2022, 40, 8536-8536.	1.6	4
4	Clinically-meaningful improvements in therapy for unresectable NSCLC. <i>Expert Review of Anticancer Therapy</i> , 2022, 22, 927-937.	2.4	6
5	Tissue surface area and tumor cell count affect the success rate of the Oncomine Dx Target Test in the analysis of biopsy tissue samples. <i>Thoracic Cancer</i> , 2021, 12, 194-200.	1.9	12
6	Suitability of Bronchoscopic Biopsy Tissue Samples for Next-Generation Sequencing. <i>Diagnostics</i> , 2021, 11, 391.	2.6	20
7	Overview of checkpoint inhibitor pneumonitis: incidence and associated risk factors. <i>Expert Opinion on Drug Safety</i> , 2021, 20, 537-547.	2.4	9
8	Pembrolizumab plus chemotherapy-induced pneumonitis in chemo-naïve patients with non-squamous non-small cell lung cancer: A multicentre, retrospective cohort study. <i>European Journal of Cancer</i> , 2021, 150, 63-72.	2.8	20
9	Readministration of Pembrolizumab after Treatment of Tuberculosis Activated by Initial Pembrolizumab Therapy. <i>Internal Medicine</i> , 2021, 60, 1743-1746.	0.7	9
10	Acute eosinophilic pneumonia after changing dosing schedule of nivolumab. <i>Japanese Journal of Clinical Oncology</i> , 2021, 51, 1766-1767.	1.3	1
11	Radiation recall pneumonitis after COVID-19 vaccination. <i>Thoracic Cancer</i> , 2021, , .	1.9	8
12	Three-Year Overall Survival with Durvalumab after Chemoradiotherapy in Stage III NSCLC—Update from PACIFIC. <i>Journal of Thoracic Oncology</i> , 2020, 15, 288-293.	1.1	328
13	Association of immune-related pneumonitis with the presence of preexisting interstitial lung disease in patients with non-small lung cancer receiving anti-programmed cell death 1 antibody. <i>Cancer Immunology, Immunotherapy</i> , 2020, 69, 15-22.	4.2	54
14	Tumor invasion in the central airway is a risk factor for early-onset checkpoint inhibitor pneumonitis in patients with non-small cell lung cancer. <i>Thoracic Cancer</i> , 2020, 11, 3576-3584.	1.9	13
15	Association between serum level soluble programmed cell death ligand 1 and prognosis in patients with non-small cell lung cancer treated with anti-PD-1 antibody. <i>Thoracic Cancer</i> , 2020, 11, 3585-3595.	1.9	32
16	Safety evaluation of durvalumab for the treatment of non-small-cell lung cancer. <i>Expert Opinion on Drug Safety</i> , 2020, 19, 653-659.	2.4	7
17	Predictive value of serum VEGF levels for elderly patients or for patients with poor performance status receiving anti-PD-1 antibody therapy for advanced non-small-cell lung cancer. <i>Cancer Immunology, Immunotherapy</i> , 2020, 69, 1229-1236.	4.2	18
18	Analysis of targeted somatic mutations in pleomorphic carcinoma of the lung using next-generation sequencing technique. <i>Thoracic Cancer</i> , 2020, 11, 2262-2269.	1.9	5

#	ARTICLE	IF	CITATIONS
19	Number of metastatic organs negatively affects the treatment sequence in patients with EGFR TKI failure. <i>Thoracic Cancer</i> , 2020, 11, 1038-1044.	1.9	0
20	Nivolumab-induced autoimmune encephalitis in an anti-neuronal autoantibody-positive patient. <i>Japanese Journal of Clinical Oncology</i> , 2019, 49, 793-794.	1.3	23
21	Tumor expression and usefulness as a biomarker of programmed death ligand 1 in advanced non-small cell lung cancer patients with preexisting interstitial lung disease. <i>Medical Oncology</i> , 2019, 36, 49.	2.5	16
22	Efficacy of subsequent docetaxel + ramucirumab and Sâ€ after nivolumab for patients with advanced nonâ€small cell lung cancer. <i>Thoracic Cancer</i> , 2019, 10, 1141-1148.	1.9	11
23	Pembrolizumab versus chemotherapy for previously untreated, PD-L1-expressing, locally advanced or metastatic non-small-cell lung cancer (KEYNOTE-042): a randomised, open-label, controlled, phase 3 trial. <i>Lancet, The</i> , 2019, 393, 1819-1830.	13.7	2,347
24	Mixed response to osimertinib and the beneficial effects of additional local therapy. <i>Thoracic Cancer</i> , 2019, 10, 738-743.	1.9	6
25	Malignant pleural effusion as a predictor of the efficacy of antiâ€PDâ€ antibody in patients with nonâ€small cell lung cancer. <i>Thoracic Cancer</i> , 2019, 10, 815-822.	1.9	20
26	Durvalumab for the treatment of non-small cell lung cancer. <i>Expert Review of Anticancer Therapy</i> , 2019, 19, 1009-1016.	2.4	20
27	Phase II study of bevacizumab, cisplatin, and pemetrexed in advanced non-squamous non-small cell lung cancer (NS-NSCLC) with EGFR wild-type. <i>Journal of Experimental Therapeutics and Oncology</i> , 2019, 13, 131-138.	0.5	4
28	Durvalumab as third-line or later treatment for advanced non-small-cell lung cancer (ATLANTIC): an open-label, single-arm, phase 2 study. <i>Lancet Oncology, The</i> , 2018, 19, 521-536.	10.7	486
29	Phase II study of nedaplatin and irinotecan as adjuvant chemotherapy for completely resected non-small cell lung cancer. <i>Cancer Chemotherapy and Pharmacology</i> , 2018, 81, 81-87.	2.3	6
30	Overall Survival with Durvalumab after Chemoradiotherapy in Stage III NSCLC. <i>New England Journal of Medicine</i> , 2018, 379, 2342-2350.	27.0	2,150
31	Does the histologic predominance of pathological stage IA lung adenocarcinoma influence the extent of resection?. <i>General Thoracic and Cardiovascular Surgery</i> , 2017, 65, 512-518.	0.9	19
32	Durvalumab after Chemoradiotherapy in Stage III Nonâ€Small-Cell Lung Cancer. <i>New England Journal of Medicine</i> , 2017, 377, 1919-1929.	27.0	3,261
33	Prognostic value of <i>EGFR</i> mutations in surgically resected pathological stage I lung adenocarcinoma. <i>Asia-Pacific Journal of Clinical Oncology</i> , 2017, 13, e204-e211.	1.1	22
34	Second predominant subtype predicts outcomes of intermediate-malignant invasive lung adenocarcinoma. <i>European Journal of Cardio-thoracic Surgery</i> , 2016, 51, ezw318.	1.4	10
35	Clonality analysis performed using human androgen receptor assay in a rare case of undifferentiated thymic carcinoma coexisting with type AB thymoma. <i>Pathology International</i> , 2016, 66, 398-403.	1.3	4
36	Progression-Free Survival, Response Rate, and Disease Control Rate as Predictors of Overall Survival in Phase III Randomized Controlled Trials Evaluating the First-Line Chemotherapy for Advanced, Locally Advanced, and Recurrent Nonâ€Small Cell Lung Carcinoma. <i>Journal of Thoracic Oncology</i> , 2016, 11, 1574-1585.	1.1	22

#	ARTICLE	IF	CITATIONS
37	Prognostic significance of vascular invasion in intermediate-grade subtype of lung adenocarcinoma. Japanese Journal of Clinical Oncology, 2016, 46, 1015-1021.	1.3	5
38	Prognostic Role of Subtype Classification in Small-Sized Pathologic N0 Invasive Lung Adenocarcinoma. Annals of Thoracic Surgery, 2016, 102, 1668-1673.	1.3	46
39	Negative prognostic influence of micropapillary pattern in stage IA lung adenocarcinoma. European Journal of Cardio-thoracic Surgery, 2016, 49, 293-299.	1.4	40
40	Phase II study of gefitinib as first-line chemotherapy in patients with advanced non-small cell lung cancer harboring EGFR mutations and poor prognostic characteristics.. Journal of Clinical Oncology, 2016, 34, e20625-e20625.	1.6	0
41	Overall survival (OS) of EGFR mutation-positive non-small cell lung cancer (NSCLC) patients: Real-world treatment patterns of 1,660 Japanese patients (pts).. Journal of Clinical Oncology, 2016, 34, e20503-e20503.	1.6	0
42	Phase II study of carboplatin and pemetrexed followed by gefitinib for patients with advanced non-small cell lung cancer harboring sensitive EGFR mutation.. Journal of Clinical Oncology, 2016, 34, e20581-e20581.	1.6	0
43	Comparison of Malignant Grade Between Pure and Partially Invasive Types of Early Lung Adenocarcinoma. Annals of Thoracic Surgery, 2015, 99, 956-960.	1.3	20
44	Prediction of lymph node status in clinical stage IA squamous cell carcinoma of the lung. European Journal of Cardio-thoracic Surgery, 2015, 47, 1022-1026.	1.4	16
45	Prognostic value of the new IASLC/ATS/ERS classification of clinical stage IA lung adenocarcinoma. Lung Cancer, 2015, 90, 199-204.	2.0	66
46	Relation Between Thin-Section Computed Tomography and Clinical Findings of Mucinous Adenocarcinoma. Annals of Thoracic Surgery, 2015, 99, 975-981.	1.3	53
47	A case of lung adenocarcinoma with multiple cavitory metastases. Japanese Journal of Clinical Oncology, 2015, 45, 504-505.	1.3	2
48	Two Cases of Stage IV Lung Adenocarcinoma That Achieved a Long-term Survival on Gefitinib. Japanese Journal of Lung Cancer, 2015, 55, 1029-1036.	0.1	2
49	Multicenter Phase II Study of Nedaplatin and Irinotecan for Patients with Squamous Cell Carcinoma of the Lung: Thoracic Oncology Research Group 0910. Anticancer Research, 2015, 35, 6705-11.	1.1	7
50	Clinical usefulness of testing for UDP glucuronosyltransferase 1 family, polypeptide A1 polymorphism prior to the initiation of irinotecan-based chemotherapy. Molecular and Clinical Oncology, 2014, 2, 737-743.	1.0	5
51	Phase II study of bevacizumab, cisplatin, and pemetrexed as first-line chemotherapy for advanced nonsquamous non-small cell lung cancer (NS-NSCLC) with EGFR wild-type.. Journal of Clinical Oncology, 2014, 32, e19125-e19125.	1.6	0
52	The clinical value and prognostic role of preoperative thin-section computed tomography findings in small-sized adenocarcinomas of the lung (10 mm or less in diameter).. Journal of Clinical Oncology, 2014, 32, e18514-e18514.	1.6	0
53	Phase I/II study of amrubicin in combination with S-1 as second-line chemotherapy for non-small-cell lung cancer without EGFR mutation. Cancer Chemotherapy and Pharmacology, 2013, 71, 705-711.	2.3	6
54	<sup>18</sup> F-fluorodeoxyglucose uptake on positron emission tomography in mucinous adenocarcinoma. European Journal of Radiology, 2013, 82, e721-e725.	2.6	13

#	ARTICLE	IF	CITATIONS
55	Periaortitis Associated with Anti-neutrophil Cytoplasmic Antibodies Induced by Bevacizumab Combination Therapy. <i>Internal Medicine</i> , 2013, 52, 589-591.	0.7	10
56	The usefulness of UGT1A1 polymorphism testing before starting irinotecan-based chemotherapy.. <i>Journal of Clinical Oncology</i> , 2013, 31, 11055-11055.	1.6	0
57	Phase II study of nedaplatin and irinotecan as adjuvant chemotherapy in patients with completely resected non-small cell lung cancer.. <i>Journal of Clinical Oncology</i> , 2013, 31, 7531-7531.	1.6	0
58	Prospective study of paclitaxel and irinotecan for elderly patients with unresectable non-small cell lung cancer. <i>Journal of Experimental Therapeutics and Oncology</i> , 2013, 10, 203-8.	0.5	0
59	Prognostic value of preoperative FDG-PET in stage IA lung adenocarcinoma. <i>European Journal of Radiology</i> , 2012, 81, 1891-1895.	2.6	13
60	Recurrent EML4-ALK-associated lung adenocarcinoma with a slow clinical course. <i>Lung Cancer</i> , 2010, 69, 361-364.	2.0	19
61	Correlation of 18F-fluorodeoxyglucose uptake on positron emission tomography with Ki-67 index and pathological invasive area in lung adenocarcinomas 30mm or less in size. <i>European Journal of Radiology</i> , 2010, 75, e62-e66.	2.6	21