

# Li-Xu Yan

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/5069230/publications.pdf>

Version: 2024-02-01

13  
papers

1,831  
citations

840776

11  
h-index

1125743

13  
g-index

13  
all docs

13  
docs citations

13  
times ranked

3496  
citing authors

#	ARTICLE	IF	CITATIONS
1	MicroRNA miR-21 overexpression in human breast cancer is associated with advanced clinical stage, lymph node metastasis and patient poor prognosis. <i>Rna</i> , 2008, 14, 2348-2360.	3.5	993
2	miR-125b Is Methylated and Functions as a Tumor Suppressor by Regulating the ETS1 Proto-oncogene in Human Invasive Breast Cancer. <i>Cancer Research</i> , 2011, 71, 3552-3562.	0.9	260
3	Eight-Signature Classifier for Prediction of Nasopharyngeal Carcinoma Survival. <i>Journal of Clinical Oncology</i> , 2011, 29, 4516-4525.	1.6	131
4	PIK3R1 targeting by miR-21 suppresses tumor cell migration and invasion by reducing PI3K/AKT signaling and reversing EMT, and predicts clinical outcome of breast cancer. <i>International Journal of Oncology</i> , 2016, 48, 471-484.	3.3	95
5	Upregulation of MiR-155 in Nasopharyngeal Carcinoma is Partly Driven by LMP1 and LMP2A and Downregulates a Negative Prognostic Marker JMJD1A. <i>PLoS ONE</i> , 2011, 6, e19137.	2.5	94
6	Epstein-Barr virus encoded latent membrane protein 1 regulates mTOR signaling pathway genes which predict poor prognosis of nasopharyngeal carcinoma. <i>Journal of Translational Medicine</i> , 2010, 8, 30.	4.4	67
7	Profiling Plasma MicroRNA in Nasopharyngeal Carcinoma with Deep Sequencing. <i>Clinical Chemistry</i> , 2014, 60, 773-782.	3.2	53
8	MYC Expression in Concert with BCL2 and BCL6 Expression Predicts Outcome in Chinese Patients with Diffuse Large B-Cell Lymphoma, Not Otherwise Specified. <i>PLoS ONE</i> , 2014, 9, e104068.	2.5	39
9	LATS2 is De-methylated and Overexpressed in Nasopharyngeal Carcinoma and Predicts Poor Prognosis. <i>BMC Cancer</i> , 2010, 10, 538.	2.6	33
10	Methylation-associated silencing of miR-200b facilitates human hepatocellular carcinoma progression by directly targeting <i>BMI1</i> . <i>Oncotarget</i> , 2016, 7, 18684-18693.	1.8	29
11	Intrahepatic cholangiocarcinoma prognostic determination using pre-operative serum C-reactive protein levels. <i>BMC Cancer</i> , 2016, 16, 792.	2.6	28
12	Clinical significance of elevated spleen tyrosine kinase expression in nasopharyngeal carcinoma. <i>Head and Neck</i> , 2012, 34, 1456-1464.	2.0	8
13	Immunophenotypic features and t(14;18) (q32;q21) translocation of Chinese follicular lymphomas helps to distinguish subgroups. <i>Diagnostic Pathology</i> , 2013, 8, 154.	2.0	1