

# Jian-Dong Li

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/5073323/publications.pdf>

Version: 2024-02-01

9  
papers

327  
citations

1307594

7  
h-index

1720034

7  
g-index

9  
all docs

9  
docs citations

9  
times ranked

449  
citing authors

#	ARTICLE	IF	CITATIONS
1	Inhibition of p38 MAPK by Glucocorticoids via Induction of MAPK Phosphatase-1 Enhances Nontypeable Haemophilus influenzae-induced Expression of Toll-like Receptor 2. <i>Journal of Biological Chemistry</i> , 2002, 277, 47444-47450.	3.4	128
2	Nontypeable Haemophilus influenzae lipoprotein P6 induces MUC5AC mucin transcription via TLR2-dependent p38 MAPK-AP1 and IKK $\beta$ -I $\kappa$ B-NF- $\kappa$ B signaling pathways. <i>Biochemical and Biophysical Research Communications</i> , 2004, 324, 1087-1094.	2.1	122
3	Phosphodiesterase 4B Mediates Extracellular Signal-regulated Kinase-dependent Up-regulation of Mucin MUC5AC Protein by Streptococcus pneumoniae by Inhibiting cAMP-protein Kinase A-dependent MKP-1 Phosphatase Pathway. <i>Journal of Biological Chemistry</i> , 2012, 287, 22799-22811.	3.4	30
4	Vinpocetine Inhibits Streptococcus pneumoniae-induced Upregulation of Mucin MUC5AC Expression via Induction of MKP-1 Phosphatase in the Pathogenesis of Otitis Media. <i>Journal of Immunology</i> , 2015, 194, 5990-5998.	0.8	16
5	Curcumin suppresses NTHi-induced CXCL5 expression via inhibition of positive IKK $\beta$ pathway and up-regulation of negative MKP-1 pathway. <i>Scientific Reports</i> , 2016, 6, 31695.	3.3	12
6	Differential regulation of Streptococcus pneumoniae-induced human MUC5AC mucin expression through distinct MAPK pathways. <i>American Journal of Translational Research (discontinued)</i> , 2009, 1, 300-11.	0.0	11
7	Curcumin Inhibits NTHi-Induced MUC5AC Mucin Overproduction in Otitis Media via Upregulation of MAPK Phosphatase MKP-1. <i>International Journal of Inflammation</i> , 2017, 2017, 1-9.	1.5	8
8	cAMP-dependent protein kinase A acts as a negative regulator for nontypeable Haemophilus influenzae-induced GM-CSF expression via inhibition of MEK-ERK signaling pathway. <i>Integrative Molecular Medicine</i> , 2016, 3, .	0.3	0
9	Receptor Interacting Protein-2 acts as negative regulator for nontypeable Haemophilus influenzae-induced mucin MUC5AC expression. <i>Integrative Molecular Medicine</i> , 2017, 4, .	0.3	0