

Leopold Eckhart

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/5098219/publications.pdf>

Version: 2024-02-01

129
papers

10,800
citations

66343

42
h-index

31849

101
g-index

131
all docs

131
docs citations

131
times ranked

21189
citing authors

#	ARTICLE	IF	CITATIONS
1	Guidelines for the use and interpretation of assays for monitoring autophagy (3rd edition). <i>Autophagy</i> , 2016, 12, 1-222.	9.1	4,701
2	Caspase-14: Analysis of Gene Structure and mRNA Expression during Keratinocyte Differentiation. <i>Biochemical and Biophysical Research Communications</i> , 2000, 277, 655-659.	2.1	393
3	Cell death by cornification. <i>Biochimica Et Biophysica Acta - Molecular Cell Research</i> , 2013, 1833, 3471-3480.	4.1	358
4	miR-17, miR-19b, miR-20a, and miR-106a are down-regulated in human aging. <i>Aging Cell</i> , 2010, 9, 291-296.	6.7	338
5	Human caspase 12 has acquired deleterious mutations. <i>Biochemical and Biophysical Research Communications</i> , 2002, 293, 722-726.	2.1	320
6	Dense sampling of bird diversity increases power of comparative genomics. <i>Nature</i> , 2020, 587, 252-257.	27.8	251
7	Knockdown of Filaggrin Impairs Diffusion Barrier Function and Increases UV Sensitivity in a Human Skin Model. <i>Journal of Investigative Dermatology</i> , 2010, 130, 2286-2294.	0.7	236
8	Terminal Differentiation of Human Keratinocytes and Stratum Corneum Formation is Associated with Caspase-14 Activation. <i>Journal of Investigative Dermatology</i> , 2000, 115, 1148-1151.	0.7	186
9	Melanin Binds Reversibly to Thermostable DNA Polymerase and Inhibits Its Activity. <i>Biochemical and Biophysical Research Communications</i> , 2000, 271, 726-730.	2.1	163
10	Histamine suppresses epidermal keratinocyte differentiation and impairs skin barrier function in a human skin model. <i>Allergy: European Journal of Allergy and Clinical Immunology</i> , 2013, 68, 37-47.	5.7	142
11	Autophagy Is Induced by UVA and Promotes Removal of Oxidized Phospholipids and Protein Aggregates in Epidermal Keratinocytes. <i>Journal of Investigative Dermatology</i> , 2013, 133, 1629-1637.	0.7	116
12	Retinoic Acid Increases the Expression of p53 and Proapoptotic Caspases and Sensitizes Keratinocytes to Apoptosis. <i>Cancer Research</i> , 2004, 64, 6542-6548.	0.9	111
13	Evolutionary Origin and Diversification of Epidermal Barrier Proteins in Amniotes. <i>Molecular Biology and Evolution</i> , 2014, 31, 3194-3205.	8.9	109
14	Increased Sensitivity of Histidinemic Mice to UVB Radiation Suggests a Crucial Role of Endogenous Urocanic Acid in Photoprotection. <i>Journal of Investigative Dermatology</i> , 2011, 131, 188-194.	0.7	108
15	Identification of reptilian genes encoding hair keratin-like proteins suggests a new scenario for the evolutionary origin of hair. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2008, 105, 18419-18423.	7.1	104
16	Caspase-14 Expression by Epidermal Keratinocytes is Regulated by Retinoids in a Differentiation-associated Manner. <i>Journal of Investigative Dermatology</i> , 2002, 119, 1150-1155.	0.7	102
17	Identification of Novel Mammalian Caspases Reveals an Important Role of Gene Loss in Shaping the Human Caspase Repertoire. <i>Molecular Biology and Evolution</i> , 2008, 25, 831-841.	8.9	95
18	Suppression of Autophagy Dysregulates the Antioxidant Response and Causes Premature Senescence of Melanocytes. <i>Journal of Investigative Dermatology</i> , 2015, 135, 1348-1357.	0.7	88

#	ARTICLE	IF	CITATIONS
19	Identification of a Homozygous PSTPIP1 Mutation in a Patient With a PAPA-Like Syndrome Responding to Canakinumab Treatment. <i>JAMA Dermatology</i> , 2013, 149, 209.	4.1	82
20	Cell aging and cellular senescence in skin aging – Recent advances in fibroblast and keratinocyte biology. <i>Experimental Gerontology</i> , 2020, 130, 110780.	2.8	81
21	Epidermal cornification is preceded by the expression of a keratinocyte-specific set of pyroptosis-related genes. <i>Scientific Reports</i> , 2017, 7, 17446.	3.3	78
22	Flagellin is the principal inducer of the antimicrobial peptide S100A7c (psoriasin) in human epidermal keratinocytes exposed to <i>Escherichia coli</i> . <i>FASEB Journal</i> , 2008, 22, 2168-2176.	0.5	72
23	The Antimicrobial Heterodimer S100A8/S100A9 (Calprotectin) Is Upregulated by Bacterial Flagellin in Human Epidermal Keratinocytes. <i>Journal of Investigative Dermatology</i> , 2010, 130, 2423-2430.	0.7	67
24	Holocrine Secretion of Sebum Is a Unique DNase2-Dependent Mode of Programmed Cell Death. <i>Journal of Investigative Dermatology</i> , 2017, 137, 587-594.	0.7	67
25	DNase1L2 Degrades Nuclear DNA during Corneocyte Formation. <i>Journal of Investigative Dermatology</i> , 2007, 127, 24-30.	0.7	65
26	Trichohyalin-Like Proteins Have Evolutionarily Conserved Roles in the Morphogenesis of Skin Appendages. <i>Journal of Investigative Dermatology</i> , 2014, 134, 2685-2692.	0.7	62
27	Convergent evolution of cysteine-rich proteins in feathers and hair. <i>BMC Evolutionary Biology</i> , 2015, 15, 82.	3.2	60
28	Essential Role of the Keratinocyte-Specific Endonuclease DNase1L2 in the Removal of Nuclear DNA from Hair and Nails. <i>Journal of Investigative Dermatology</i> , 2011, 131, 1208-1215.	0.7	59
29	Epidermal keratinocytes form a functional skin barrier in the absence of Atg7 dependent autophagy. <i>Journal of Dermatological Science</i> , 2013, 71, 67-75.	1.9	59
30	Is the Filaggrin-Histidine-Urocanic Acid Pathway Essential for Stratum Corneum Acidification?. <i>Journal of Investigative Dermatology</i> , 2010, 130, 2141-2144.	0.7	56
31	Impact of filaggrin mutations on Raman spectra and biophysical properties of the stratum corneum in mild to moderate atopic dermatitis. <i>Journal of the European Academy of Dermatology and Venereology</i> , 2012, 26, 983-990.	2.4	53
32	The molecular organization of the beta-sheet region in Corneous beta-proteins (beta-keratins) of sauropsids explains its stability and polymerization into filaments. <i>Journal of Structural Biology</i> , 2016, 194, 282-291.	2.8	53
33	Evidence That Caspase-13 Is Not a Human but a Bovine Gene. <i>Biochemical and Biophysical Research Communications</i> , 2001, 285, 1150-1154.	2.1	52
34	Autophagic Control of Skin Aging. <i>Frontiers in Cell and Developmental Biology</i> , 2019, 7, 143.	3.7	52
35	Differential Evolution of the Epidermal Keratin Cytoskeleton in Terrestrial and Aquatic Mammals. <i>Molecular Biology and Evolution</i> , 2019, 36, 328-340.	8.9	51
36	Stratum corneum-derived caspase-14 is catalytically active. <i>FEBS Letters</i> , 2004, 577, 446-450.	2.8	50

#	ARTICLE	IF	CITATIONS
37	Identification and Characterization of a Novel Mammalian Caspase with Proapoptotic Activity. <i>Journal of Biological Chemistry</i> , 2005, 280, 35077-35080.	3.4	50
38	Review: Evolution and diversification of corneous beta-keratins, the characteristic epidermal proteins of reptiles and birds. <i>Journal of Experimental Zoology Part B: Molecular and Developmental Evolution</i> , 2018, 330, 438-453.	1.3	48
39	Autophagy in the Thymic Epithelium Is Dispensable for the Development of Self-Tolerance in a Novel Mouse Model. <i>PLoS ONE</i> , 2012, 7, e38933.	2.5	47
40	Comparative Genomics Identifies Epidermal Proteins Associated with the Evolution of the Turtle Shell. <i>Molecular Biology and Evolution</i> , 2016, 33, 726-737.	8.9	46
41	The skin barrier: Epidermis vs environment. <i>Experimental Dermatology</i> , 2018, 27, 805-806.	2.9	46
42	Degradation by Stratum Corneum Proteases Prevents Endogenous RNase Inhibitor from Blocking Antimicrobial Activities of RNase 5 and RNase 7. <i>Journal of Investigative Dermatology</i> , 2009, 129, 2193-2201.	0.7	45
43	Pangolins Lack IFIH1/MDA5, a Cytoplasmic RNA Sensor That Initiates Innate Immune Defense Upon Coronavirus Infection. <i>Frontiers in Immunology</i> , 2020, 11, 939.	4.8	45
44	Psoriasin (S100A7) is a major <i>Escherichia coli</i> -cidal factor of the female genital tract. <i>Mucosal Immunology</i> , 2010, 3, 602-609.	6.0	42
45	DNase 2 Is the Main DNA-Degrading Enzyme of the Stratum Corneum. <i>PLoS ONE</i> , 2011, 6, e17581.	2.5	42
46	Caspase-14 but not caspase-3 is processed during the development of fetal mouse epidermis. <i>Differentiation</i> , 2005, 73, 406-413.	1.9	41
47	Targeted deletion of Atg5 reveals differential roles of autophagy in keratin K5-expressing epithelia. <i>Biochemical and Biophysical Research Communications</i> , 2013, 430, 689-694.	2.1	41
48	Reverse Transcription-Polymerase Chain Reaction Products of Alternatively Spliced mRNAs Form DNA Heteroduplexes and Heteroduplex Complexes. <i>Journal of Biological Chemistry</i> , 1999, 274, 2613-2615.	3.4	39
49	Hepatocyte Growth Factor/Scatter Factor Inhibits UVB-induced Apoptosis of Human Keratinocytes but Not of Keratinocyte-derived Cell Lines via the Phosphatidylinositol 3-Kinase/AKT Pathway. <i>Journal of Biological Chemistry</i> , 2002, 277, 14146-14152.	3.4	36
50	Ultrastructural Localization of Caspase-14 in Human Epidermis. <i>Journal of Histochemistry and Cytochemistry</i> , 2004, 52, 1561-1574.	2.5	36
51	Against the Rules: Human Keratin K80. <i>Journal of Biological Chemistry</i> , 2010, 285, 36909-36921.	3.4	36
52	Comparative genomics reveals conservation of filaggrin and loss of caspase-14 in dolphins. <i>Experimental Dermatology</i> , 2015, 24, 365-369.	2.9	35
53	Convergent Evolution of Cysteine-Rich Keratins in Hard Skin Appendages of Terrestrial Vertebrates. <i>Molecular Biology and Evolution</i> , 2020, 37, 982-993.	8.9	33
54	Loss of Keratin K2 Expression Causes Aberrant Aggregation of K10, Hyperkeratosis, and Inflammation. <i>Journal of Investigative Dermatology</i> , 2014, 134, 2579-2588.	0.7	31

#	ARTICLE	IF	CITATIONS
55	Terminal differentiation of nail matrix keratinocytes involves up-regulation of DNase1L2 but is independent of caspase-14 expression. <i>Differentiation</i> , 2007, 75, 939-946.	1.9	29
56	Identification and comparative analysis of the epidermal differentiation complex in snakes. <i>Scientific Reports</i> , 2017, 7, 45338.	3.3	29
57	Identification of a Human cDNA Encoding a Novel Bcl-x Isoform. <i>Biochemical and Biophysical Research Communications</i> , 1998, 248, 147-152.	2.1	28
58	Filaggrin Expression and Processing Deficiencies Impair Corneocyte Surface Texture and Stiffness in Mice. <i>Journal of Investigative Dermatology</i> , 2020, 140, 615-623.e5.	0.7	28
59	Histidase expression in human epidermal keratinocytes: Regulation by differentiation status and all-trans retinoic acid. <i>Journal of Dermatological Science</i> , 2008, 50, 209-215.	1.9	27
60	In situ labeling of DNA reveals interindividual variation in nuclear DNA breakdown in hair and may be useful to predict success of forensic genotyping of hair. <i>International Journal of Legal Medicine</i> , 2012, 126, 63-70.	2.2	27
61	Inactivation of DNase1L2 and DNase2 in keratinocytes suppresses DNA degradation during epidermal cornification and results in constitutive parakeratosis. <i>Scientific Reports</i> , 2017, 7, 6433.	3.3	27
62	Inactivation of autophagy leads to changes in sebaceous gland morphology and function. <i>Experimental Dermatology</i> , 2018, 27, 1142-1151.	2.9	27
63	Alternative Splicing of Caspase-8 mRNA during Differentiation of Human Leukocytes. <i>Biochemical and Biophysical Research Communications</i> , 2001, 289, 777-781.	2.1	26
64	The anatomy and development of the claws of <i>Xenopus laevis</i> (Lissamphibia: Anura) reveal alternate pathways of structural evolution in the integument of tetrapods. <i>Journal of Anatomy</i> , 2009, 214, 607-619.	1.5	26
65	Mechanisms and emerging functions of DNA degradation in the epidermis. <i>Frontiers in Bioscience - Landmark</i> , 2012, 17, 2461.	3.0	26
66	Comparative Analysis of Epidermal Differentiation Genes of Crocodylians Suggests New Models for the Evolutionary Origin of Avian Feather Proteins. <i>Genome Biology and Evolution</i> , 2018, 10, 694-704.	2.5	26
67	Distribution of caspase-14 in epidermis and hair follicles is evolutionarily conserved among mammals. <i>The Anatomical Record Part A: Discoveries in Molecular, Cellular, and Evolutionary Biology</i> , 2005, 286A, 962-973.	2.0	22
68	Immunolocalization of a Histidine-Rich Epidermal Differentiation Protein in the Chicken Supports the Hypothesis of an Evolutionary Developmental Link between the Embryonic Subperiderm and Feather Barbs and Barbules. <i>PLoS ONE</i> , 2016, 11, e0167789.	2.5	22
69	Deleterious Mutations of a Claw Keratin in Multiple Taxa of Reptiles. <i>Journal of Molecular Evolution</i> , 2011, 72, 265-273.	1.8	21
70	ATG7 is essential for secretion of iron from ameloblasts and normal growth of murine incisors during aging. <i>Autophagy</i> , 2020, 16, 1851-1857.	9.1	20
71	2,3,7,8-Tetrachlorodibenzo-p-Dioxin Impairs Differentiation of Normal Human Epidermal Keratinocytes in a Skin Equivalent Model. <i>Journal of Investigative Dermatology</i> , 2005, 124, 275-277.	0.7	19
72	Keratins K2 and K10 are essential for the epidermal integrity of plantar skin. <i>Journal of Dermatological Science</i> , 2016, 81, 10-16.	1.9	19

#	ARTICLE	IF	CITATIONS
73	Evolution of Trichocyte Keratins. <i>Advances in Experimental Medicine and Biology</i> , 2018, 1054, 33-45.	1.6	19
74	The Differentiation-Associated Keratinocyte Protein Cornifelin Contributes to Cell-Cell Adhesion of Epidermal and Mucosal Keratinocytes. <i>Journal of Investigative Dermatology</i> , 2019, 139, 2292-2301.e9.	0.7	19
75	Phylogenetic profiling and gene expression studies implicate a primary role of <scp>PSORS</scp>1C2 in terminal differentiation of keratinocytes. <i>Experimental Dermatology</i> , 2017, 26, 352-358.	2.9	18
76	Cornification of nail keratinocytes requires autophagy for bulk degradation of intracellular proteins while sparing components of the cytoskeleton. <i>Apoptosis: an International Journal on Programmed Cell Death</i> , 2019, 24, 62-73.	4.9	18
77	TINCR is not a non-coding RNA but encodes a protein component of cornified epidermal keratinocytes. <i>Experimental Dermatology</i> , 2020, 29, 376-379.	2.9	18
78	Autophagy in epithelial homeostasis and defense. <i>Frontiers in Bioscience - Elite</i> , 2013, E5, 1000-1010.	1.8	17
79	Aldehyde dehydrogenase 1A3 is transcriptionally activated by all-trans-retinoic acid in human epidermal keratinocytes. <i>Biochemical and Biophysical Research Communications</i> , 2010, 400, 207-211.	2.1	16
80	Ultrastructural localization of hair keratin homologs in the claw of the lizard <i>Anolis carolinensis</i>. <i>Journal of Morphology</i> , 2011, 272, 363-370.	1.2	16
81	Cytosolic DNA sensing through cGAS and STING is inactivated by gene mutations in pangolins. <i>Apoptosis: an International Journal on Programmed Cell Death</i> , 2020, 25, 474-480.	4.9	16
82	Phylogenomics of caspase-activated DNA fragmentation factor. <i>Biochemical and Biophysical Research Communications</i> , 2007, 356, 293-299.	2.1	15
83	Control of cell death-associated danger signals during cornification prevents autoinflammation of the skin. <i>Experimental Dermatology</i> , 2018, 27, 884-891.	2.9	15
84	Immunolocalization and phylogenetic profiling of the feather protein with the highest cysteine content. <i>Protoplasma</i> , 2019, 256, 1257-1265.	2.1	15
85	Genome Sequence Comparison Reveals Independent Inactivation of the Caspase-15 Gene in Different Evolutionary Lineages of Mammals. <i>Molecular Biology and Evolution</i> , 2006, 23, 2081-2089.	8.9	14
86	Transcription of the caspase-14 gene in human epidermal keratinocytes requires AP-1 and NF- κ B. <i>Biochemical and Biophysical Research Communications</i> , 2008, 371, 261-266.	2.1	14
87	Cuts by Caspase-14 Control the Proteolysis of Filaggrin. <i>Journal of Investigative Dermatology</i> , 2011, 131, 2173-2175.	0.7	14
88	The Expression of the Endogenous mTORC1 Inhibitor Sestrin 2 Is Induced by UVB and Balanced with the Expression Level of Sestrin 1. <i>PLoS ONE</i> , 2016, 11, e0166832.	2.5	14
89	Filaggrin has evolved from an "S100 fused" type protein-(<scp>SFTP</scp>) gene present in a common ancestor of amphibians and mammals. <i>Experimental Dermatology</i> , 2017, 26, 955-957.	2.9	14
90	Double deficiency of Trex2 and DNase1L2 nucleases leads to accumulation of DNA in lingual cornifying keratinocytes without activating inflammatory responses. <i>Scientific Reports</i> , 2017, 7, 11902.	3.3	14

#	ARTICLE	IF	CITATIONS
91	Multifaceted role of TREX2 in the skin defense against UV-induced skin carcinogenesis. <i>Oncotarget</i> , 2015, 6, 22375-22396.	1.8	14
92	New facets of keratin K77: interspecies variations of expression and different intracellular location in embryonic and adult skin of humans and mice. <i>Cell and Tissue Research</i> , 2013, 354, 793-812.	2.9	13
93	Immunolocalization of Scaffoldin, a Trichohyalin-Like Protein, in the Epidermis of the Chicken Embryo. <i>Anatomical Record</i> , 2015, 298, 479-487.	1.4	13
94	Tyrosinase-Cre-Mediated Deletion of the Autophagy Gene Atg7 Leads to Accumulation of the RPE65 Variant M450 in the Retinal Pigment Epithelium of C57BL/6 Mice. <i>PLoS ONE</i> , 2016, 11, e0161640.	2.5	13
95	Filamentous Aggregation of Sequestosome-1/p62 in Brain Neurons and Neuroepithelial Cells upon Tyr-Cre-Mediated Deletion of the Autophagy Gene Atg7. <i>Molecular Neurobiology</i> , 2018, 55, 8425-8437.	4.0	13
96	MCPIP1 contributes to the inflammatory response of UVB-treated keratinocytes. <i>Journal of Dermatological Science</i> , 2017, 87, 10-18.	1.9	12
97	Identification of epidermal differentiation genes of the tuatara provides insights into the early evolution of lepidosaurian skin. <i>Scientific Reports</i> , 2020, 10, 12844.	3.3	12
98	Assessing Autophagy in Archived Tissue or How to Capture Autophagic Flux from a Tissue Snapshot. <i>Biology</i> , 2020, 9, 59.	2.8	12
99	The Whey Acidic Protein WFDC12 Is Specifically Expressed in Terminally Differentiated Keratinocytes and Regulates Epidermal Serine Protease Activity. <i>Journal of Investigative Dermatology</i> , 2021, 141, 1198-1206.e13.	0.7	12
100	Gene duplications and gene loss in the epidermal differentiation complex during the evolutionary land-to-water transition of cetaceans. <i>Scientific Reports</i> , 2021, 11, 12334.	3.3	12
101	Localisation of keratin K78 in the basal layer and first suprabasal layers of stratified epithelia completes expression catalogue of type II keratins and provides new insights into sequential keratin expression. <i>Cell and Tissue Research</i> , 2016, 363, 735-750.	2.9	11
102	Duplication of the caspase-12 prodomain and inactivation of NLR4/IPAF in the dog. <i>Biochemical and Biophysical Research Communications</i> , 2009, 384, 226-230.	2.1	10
103	Suppression of autophagy perturbs turnover of sequestosome-1/p62 in Merkel cells but not in keratinocytes. <i>Journal of Dermatological Science</i> , 2018, 90, 209-211.	1.9	10
104	Suppression of Epithelial Autophagy Compromises the Homeostasis of Sweat Glands during Aging. <i>Journal of Investigative Dermatology</i> , 2018, 138, 2061-2063.	0.7	10
105	Comparative genomics suggests loss of keratin K24 in three evolutionary lineages of mammals. <i>Scientific Reports</i> , 2019, 9, 10924.	3.3	10
106	A Stress Response Program at the Origin of Evolutionary Innovation in the Skin. <i>Evolutionary Bioinformatics</i> , 2019, 15, 117693431986224.	1.2	10
107	The caspase-1 inhibitor CARD18 is specifically expressed during late differentiation of keratinocytes and its expression is lost in lichen planus. <i>Journal of Dermatological Science</i> , 2017, 87, 176-182.	1.9	8
108	PIWIL-2 and piRNAs are regularly expressed in epithelia of the skin and their expression is related to differentiation. <i>Archives of Dermatological Research</i> , 2020, 312, 705-714.	1.9	8

#	ARTICLE	IF	CITATIONS
109	Urocanic Acid: An Endogenous Regulator of Langerhans Cells. <i>Journal of Investigative Dermatology</i> , 2016, 136, 1735-1737.	0.7	7
110	Autophagy - how to control your intracellular diet. <i>British Journal of Dermatology</i> , 2017, 176, 1417-1419.	1.5	7
111	Identification of a novel exon encoding the amino-terminus of the predominant caspase-5 variants. <i>Biochemical and Biophysical Research Communications</i> , 2006, 348, 682-688.	2.1	6
112	The tail domains of keratins contain conserved amino acid sequence motifs. <i>Journal of Dermatological Science</i> , 2009, 54, 208-209.	1.9	6
113	An In Vitro Model of Avian Skin Reveals Evolutionarily Conserved Transcriptional Regulation of Epidermal Barrier Formation. <i>Journal of Investigative Dermatology</i> , 2021, 141, 2829-2837.	0.7	6
114	Autophagy protects murine preputial glands against premature aging, and controls their sebum phospholipid and pheromone profile. <i>Autophagy</i> , 2022, 18, 1005-1019.	9.1	6
115	A basement membrane-like matrix formed by cell-released proteins at the medium/air interface supports growth of keratinocytes. <i>European Journal of Cell Biology</i> , 2003, 82, 549-555.	3.6	5
116	The Trichohyalin-Like Protein Scaffoldin Is Expressed in the Multilayered Periderm during Development of Avian Beak and Egg Tooth. <i>Genes</i> , 2021, 12, 248.	2.4	5
117	Immunolocalization of epidermal differentiation complex proteins reveals distinct molecular compositions of cells that control structure and mechanical properties of avian skin appendages. <i>Journal of Morphology</i> , 2021, 282, 917-933.	1.2	5
118	Identification of Chicken Transglutaminase 1 and In Situ Localization of Transglutaminase Activity in Avian Skin and Esophagus. <i>Genes</i> , 2021, 12, 1565.	2.4	4
119	Single-cell transcriptomics defines keratinocyte differentiation in avian scutate scales. <i>Scientific Reports</i> , 2022, 12, 126.	3.3	4
120	Ultrastructural characterization of an artificial basement membrane produced by cultured keratinocytes. <i>Journal of Biomedical Materials Research - Part A</i> , 2005, 73A, 158-164.	4.0	3
121	Caspase-5 is autoprocessed at two sites that contain an aspartate residue in the C^{pro} domain. <i>Biochemical and Biophysical Research Communications</i> , 2006, 348, 682-688.	2.1	3
122	Don't be so overprotective! <i>EMBO Molecular Medicine</i> , 2012, 4, 362-363.	6.9	3
123	Immunolocalization of loricrin in the maturing layer of normal and regenerating epidermis of the lizard <i>Anolis carolinensis</i> . <i>Journal of Experimental Zoology Part B: Molecular and Developmental Evolution</i> , 2015, 324, 159-167.	1.3	3
124	Identification of New Biological Pathways Involved in Skin Aging From the Analysis of French Women Genome-Wide Data. <i>Frontiers in Genetics</i> , 2022, 13, 836581.	2.3	3
125	NOD2 and reproduction-associated NOD-like receptors have been lost during the evolution of pangolins. <i>Immunogenetics</i> , 2021, , 1.	2.4	2
126	Comparative genomics reveals evolutionary loss of epiplakin in cetaceans. <i>Scientific Reports</i> , 2022, 12, 1112.	3.3	2

#	ARTICLE	IF	CITATIONS
127	Urocanic Acid and Skin Photodamage: New Light on an Old Chromophore. , 2016, , 79-99.		1
128	Cerebellar Degeneration-related Antigen 1 Is Ubiquitously Expressed in Human Epidermis and Dermis. Current Medical Science, 2020, 40, 570-573.	1.8	1
129	Roles of Autophagy in the Thymic Epithelium. , 2015, , 231-240.		0