

# Fei Ye

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/5101191/publications.pdf>

Version: 2024-02-01

57  
papers

1,716  
citations

471509

17  
h-index

289244

40  
g-index

59  
all docs

59  
docs citations

59  
times ranked

1713  
citing authors

#	ARTICLE	IF	CITATIONS
1	Intravascular Ultrasound Versus Angiography-Guided Drug-Eluting Stent Implantation. <i>Journal of the American College of Cardiology</i> , 2018, 72, 3126-3137.	2.8	392
2	3-Year Outcomes of the ULTIMATE Trial Comparing Intravascular Ultrasound Versus Angiography-Guided Drug-Eluting Stent Implantation. <i>JACC: Cardiovascular Interventions</i> , 2021, 14, 247-257.	2.9	149
3	Clinical Outcome After DK Crush Versus Culotte Stenting of Distal Left Main Bifurcation Lesions. <i>JACC: Cardiovascular Interventions</i> , 2015, 8, 1335-1342.	2.9	137
4	Multicentre, randomized comparison of two-stent and provisional stenting techniques in patients with complex coronary bifurcation lesions: the DEFINITION II trial. <i>European Heart Journal</i> , 2020, 41, 2523-2536.	2.2	124
5	Cutoff Value and Long-Term Prediction of Clinical Events by FFR Measured Immediately After Implantation of a Drug-Eluting Stent in Patients With Coronary Artery Disease. <i>JACC: Cardiovascular Interventions</i> , 2017, 10, 986-995.	2.9	111
6	Randomized Comparison of FFR-Guided and Angiography-Guided Provisional Stenting of True Coronary Bifurcation Lesions. <i>JACC: Cardiovascular Interventions</i> , 2015, 8, 536-546.	2.9	101
7	Clinical Outcome of Double Kissing Crush Versus Provisional Stenting of Coronary Artery Bifurcation Lesions. <i>Circulation: Cardiovascular Interventions</i> , 2017, 10, .	3.9	81
8	Clinical implications of three-vessel fractional flow reserve measurement in patients with coronary artery disease. <i>European Heart Journal</i> , 2018, 39, 945-951.	2.2	68
9	Incidence and Clinical Outcomes of Stent Fractures on the Basis of 6,555 Patients and 16,482 Drug-Eluting Stents From 4 Centers. <i>JACC: Cardiovascular Interventions</i> , 2016, 9, 1115-1123.	2.9	66
10	Prognostic Implication of Functional Incomplete Revascularization and Residual Functional SYNTAX Score in Patients With Coronary Artery Disease. <i>JACC: Cardiovascular Interventions</i> , 2018, 11, 237-245.	2.9	51
11	Comparison of one-year clinical outcomes between intravascular ultrasound-guided versus angiography-guided implantation of drug-eluting stents for left main lesions: a single-center analysis of a 1,016-patient cohort. <i>Patient Preference and Adherence</i> , 2014, 8, 1299.	1.8	43
12	Role of Post-Stent Physiological Assessment in a Risk Prediction Model After Coronary Stent Implantation. <i>JACC: Cardiovascular Interventions</i> , 2020, 13, 1639-1650.	2.9	36
13	Intravascular ultrasound-guided drug-eluting stent implantation is associated with improved clinical outcomes in patients with unstable angina and complex coronary artery true bifurcation lesions. <i>International Journal of Cardiovascular Imaging</i> , 2018, 34, 1685-1696.	1.5	34
14	The Anatomic- and Clinical-Based NERS (New Risk Stratification) Score II to Predict Clinical Outcomes After Stenting Unprotected Left Main Coronary Artery Disease. <i>JACC: Cardiovascular Interventions</i> , 2013, 6, 1233-1241.	2.9	24
15	The role of HYAL2 in LSS-induced glycocalyx impairment and the PKA-mediated decrease in eNOS Ser-633 phosphorylation and nitric oxide production. <i>Molecular Biology of the Cell</i> , 2016, 27, 3972-3979.	2.1	24
16	Spontaneous Coronary Artery Dissection as a Cause of Acute Myocardial Infarction in Young Female Population. <i>Chinese Medical Journal</i> , 2017, 130, 1534-1539.	2.3	21
17	The Acute Changes of Fractional Flow Reserve in DK (Double Kissing), Crush, and Stent Technique for True Bifurcation Lesions. <i>Journal of Interventional Cardiology</i> , 2010, 23, 341-345.	1.2	19
18	Prognostic Impact of Residual Anatomic Disease Burden After Functionally Complete Revascularization. <i>Circulation: Cardiovascular Interventions</i> , 2020, 13, e009232.	3.9	16

#	ARTICLE	IF	CITATIONS
19	The impact of everolimus versus other rapamycin derivative-eluting stents on clinical outcomes in patients with coronary artery disease: A meta-analysis of 16 randomized trials. <i>Journal of Cardiology</i> , 2014, 64, 185-193.	1.9	15
20	High platelet reactivity affects the clinical outcomes of patients undergoing percutaneous coronary intervention. <i>BMC Cardiovascular Disorders</i> , 2016, 16, 240.	1.7	15
21	Clinical Outcomes of Deferred Lesions With Angiographically Insignificant Stenosis But Low Fractional Flow Reserve. <i>Journal of the American Heart Association</i> , 2017, 6, .	3.7	14
22	Comparison of intravascular ultrasound-guided with angiography-guided double kissing crush stenting for patients with complex coronary bifurcation lesions: Rationale and design of a prospective, randomized, and multicenter DKCRUSH VIII trial. <i>American Heart Journal</i> , 2021, 234, 101-110.	2.7	12
23	Angioscopy study from a large patient population comparing sirolimus-eluting stent with biodegradable versus durable polymer. <i>Catheterization and Cardiovascular Interventions</i> , 2012, 80, 420-428.	1.7	10
24	Ivabradine promotes angiogenesis and reduces cardiac hypertrophy in mice with myocardial infarction. <i>Anatolian Journal of Cardiology</i> , 2018, 20, 266-272.	0.9	10
25	Clinical Relevance of Functionally Insignificant Moderate Coronary Artery Stenosis Assessed by 3â€Vessel Fractional Flow Reserve Measurement. <i>Journal of the American Heart Association</i> , 2018, 7, .	3.7	9
26	Ginkgolide A protects adverse cardiac remodeling through enhancing antioxidation and nitric oxide utilization in mice with pressure overload. <i>Die Pharmazie</i> , 2019, 74, 698-702.	0.5	9
27	Advantages of Transradial Rotational Atherectomy versus Transfemoral Approach in Elderly Patients with Hard-Handling Calcified Coronary Lesions - A Single Center Experience. <i>Acta Cardiologica Sinica</i> , 2018, 34, 464-471.	0.2	9
28	Treatment effects of systematic two-stent and provisional stenting techniques in patients with complex coronary bifurcation lesions: rationale and design of a prospective, randomised and multicentre DEFINITION II trial. <i>BMJ Open</i> , 2018, 8, e020019.	1.9	7
29	Prognostic Value of Prevascularization Fractional Flow Reserve Mediated by the Postrevascularization Level. <i>JAMA Network Open</i> , 2020, 3, e2018162.	5.9	7
30	Comparison in Prevalence, Predictors, and Clinical Outcome of VSR Versus FWR after Acute Myocardial Infarction: The Prospective, Multicenter Registry MOODY Trial-Heart Rupture Analysis. <i>Cardiovascular Revascularization Medicine</i> , 2019, 20, 1158-1164.	0.8	6
31	Ticagrelor versus clopidogrel for prevention of subclinical stent thrombosis detected by optical coherence tomography in patients with drug-eluting stent implantationâ€”a multicenter and randomized study. <i>Platelets</i> , 2021, 32, 404-412.	2.3	6
32	Comparison between twoâ€dimensional and threeâ€dimensional quantitative coronary angiography for the prediction of functional severity in true bifurcation lesions: Insights from the randomized DKâ€CRUSH II, III, and IV trials. <i>Catheterization and Cardiovascular Interventions</i> , 2016, 87, 589-598.	1.7	5
33	Stent fracture is associated with a higher mortality in patients with type-2 diabetes treated by implantation of a second-generation drug-eluting stent. <i>International Journal of Cardiovascular Imaging</i> , 2017, 33, 1873-1881.	1.5	5
34	Correlation between preâ€procedural plaque morphology and patterns of inâ€stent neointimal hyperplasia at 1â€year followâ€up in patients treated with newâ€generation drugâ€eluting stents: An optical coherence tomography based analysis. <i>Journal of Interventional Cardiology</i> , 2018, 31, 737-746.	1.2	5
35	Zinc finger protein 91 loss induces cardiac hypertrophy through adenosine A1 receptor downâ€regulation under pressure overload status. <i>Journal of Cellular and Molecular Medicine</i> , 2020, 24, 10189-10201.	3.6	5
36	Optical coherence tomography predictors of target vessel myocardial infarction after provisional stenting in patients with coronary bifurcation disease. <i>Catheterization and Cardiovascular Interventions</i> , 2021, 97, 1331-1340.	1.7	5

#	ARTICLE	IF	CITATIONS
37	Impact of biodegradable versus durable polymer drug-eluting stents on clinical outcomes in patients with coronary artery disease: a meta-analysis of 15 randomized trials. <i>Chinese Medical Journal</i> , 2014, 127, 2159-66.	2.3	5
38	Cardamonin protects against adverse cardiac remodeling through mTORC1 inhibition in mice with myocardial infarction. <i>Die Pharmazie</i> , 2018, 73, 508-512.	0.5	5
39	Rare Solitary Fibrous Tumor in the Pulmonary Artery Mimicking Pulmonary Embolism. <i>Circulation: Cardiovascular Imaging</i> , 2017, 10, .	2.6	4
40	Improvement of clinical outcome in patients with STâ€elevation myocardial infarction between 1999 and 2016 in China: The Prospective, Multicentre Registry MOODY study. <i>European Journal of Clinical Investigation</i> , 2020, 50, e13197.	3.4	4
41	Cut-off values of lesion and vessel quantitative flow ratio in de novo coronary lesion post-drug-coated balloon therapy predicting vessel restenosis at mid-term follow-up. <i>Chinese Medical Journal</i> , 2021, 134, 1450-1456.	2.3	4
42	Amelioratory effect of neoandrographolide on myocardial ischemicâ€ereperfusion injury by its antiâ€enflammatory and antiâ€eapoptotic activities. <i>Environmental Toxicology</i> , 2021, 36, 2367-2379.	4.0	4
43	Association of periâ€eprocedural myocardial infarction with mortality after stenting true coronary bifurcation lesions: A pooled individual participant data analysis from four randomized controlled trials. <i>Catheterization and Cardiovascular Interventions</i> , 2021, , .	1.7	4
44	Retrospective analysis of OCT on MB characteristics and 1-year follow-up of the ISR incidence after the DES implantation in patients with MB. <i>Scientific Reports</i> , 2022, 12, 534.	3.3	4
45	Rationale and design for comparison of non-compliant balloon with drug-coating balloon angioplasty for side branch after provisional stenting for patients with true coronary bifurcation lesions: a prospective, multicentre and randomised DCB-BIF trial. <i>BMJ Open</i> , 2022, 12, e052788.	1.9	4
46	Intensive statin versus low-dose statin + ezetimibe treatment for fibrous cap thickness of coronary vulnerable plaques. <i>Chinese Medical Journal</i> , 2020, 133, 2415-2421.	2.3	3
47	[18F]FEDAC translocator protein positron emission tomographyâ€ecomputed tomography for early detection of mitochondrial dysfunction secondary to myocardial ischemia. <i>Annals of Nuclear Medicine</i> , 2021, 35, 927-936.	2.2	3
48	Differential Prognostic Implications of Pre- and Post-Stent Fractional Flow Reserve in Patients Undergoing Percutaneous Coronary Intervention. <i>Korean Circulation Journal</i> , 2022, 52, 47.	1.9	3
49	Comparison of clinical outcomes of Chinese men and women after coronary stenting for coronary artery disease: a multi-center retrospective analysis of 4,334 patients. <i>Journal of Biomedical Research</i> , 2014, 28, 368.	1.6	3
50	Postâ€ePCI quantitative flow ratio predicts 3â€eyear outcome after rotational atherectomy in patients with heavily calcified lesions. <i>Clinical Cardiology</i> , 2022, , .	1.8	3
51	NOBORIâ„¢ biodegradable-polymer biolimus-eluting stent versus durable-polymer drug-eluting stents: A meta-analysis. <i>International Journal of Cardiology</i> , 2014, 174, 151-153.	1.7	2
52	Serum biomarker analysis at the protein level on pulmonary hypertension secondary to old anterior myocardial infarction. <i>Pulmonary Circulation</i> , 2020, 10, 1-13.	1.7	2
53	The association of diabetes mellitus with neointimal formation following deployment of second-generation drug-eluting stents: an optical coherence tomographic study. <i>Coronary Artery Disease</i> , 2021, 32, 105-111.	0.7	2
54	Effect of Coronary Disease Characteristics on Prognostic Relevance of Residual Ischemia After Stent Implantation. <i>Frontiers in Cardiovascular Medicine</i> , 2021, 8, 696756.	2.4	2

#	ARTICLE	IF	CITATIONS
55	Outcomes in Patients Fully Covered With Coronary Artery Aneurysm and Stenosis Lesion by Second Generation Drug-Eluting Stents After 1 Year. <i>Angiology</i> , 2020, 71, 942-947.	1.8	1
56	Relationship between neointimal strut bridge and jailed side-branch ostial area. <i>Herz</i> , 2021, 46, 178-187.	1.1	1
57	Apnea Hypopnea Index is an Independent Predictor of Coronary Microcirculatory Dysfunction in Stable Angina Pectoris Patients with a Single Borderline Lesion. <i>Acta Cardiologica Sinica</i> , 2020, 36, 207-215.	0.2	0