
List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/5185946/publications.pdf Version: 2024-02-01



| # | Article | IF | CITATIONS |
|----|--|-----|-----------|
| 1 | Duration of SARS-CoV-2 viremia and its correlation to mortality and inflammatory parameters in patients hospitalized for COVID-19: a cohort study. Diagnostic Microbiology and Infectious Disease, 2022, 102, 115595. | 1.8 | 28 |
| 2 | Risk factors for death in adult COVID-19 patients: Frailty predicts fatal outcome in older patients. International Journal of Infectious Diseases, 2021, 102, 415-421. | 3.3 | 85 |
| 3 | Brace versus cast following surgical treatment of distal radial fracture: a prospective randomised study comparing quality of recovery. F1000Research, 2021, 10, 336. | 1.6 | 0 |
| 4 | Postâ€operative hypertension during early recovery following liver tumour ablation: A retrospective study. Acta Anaesthesiologica Scandinavica, 2021, 65, 1248-1253. | 1.6 | 1 |
| 5 | Emergencies in freestanding ambulatory surgery centre. Current Opinion in Anaesthesiology, 2021, Publish Ahead of Print, 690-694. | 2.0 | 3 |
| 6 | Supraclavicular block with Mepivacaine vs Ropivacaine, their impact on postoperative pain: a prospective randomised study. BMC Anesthesiology, 2021, 21, 273. | 1.8 | 4 |
| 7 | Workplace exposure to carbon dioxide during routine laparoscopy – is it safe?. F1000Research, 2020, 9, 571. | 1.6 | Ο |
| 8 | Workplace exposure to carbon dioxide during routine laparoscopy – is it safe?. F1000Research, 2020, 9, 571. | 1.6 | 2 |
| 9 | Can simple perioperative measures improve quality of recovery following ambulatory laparoscopic surgery in females? An open prospective randomised cohort study, comparing nutritional preoperative drink and chewing gum during recovery to standard care. Annals of Medicine and Surgery, 2019, 44, 20-25. | 1.1 | 8 |
| 10 | European Society of Anaesthesiology Task Force on Nitrous Oxide: a narrative review of its role in clinical practice. British Journal of Anaesthesia, 2019, 122, 587-604. | 3.4 | 66 |
| 11 | Recovery and discharge criteria after ambulatory anesthesia: can we improve them?. Current Opinion in Anaesthesiology, 2019, 32, 698-702. | 2.0 | 15 |
| 12 | High Frequency Jet Ventilation during stereotactic ablation of liver tumours: an observational study on blood gas analysis as a measure of lung function during general anaesthesia. F1000Research, 2019, 8, 386. | 1.6 | 11 |
| 13 | Acute hip fracture surgery anaesthetic technique and 30-day mortality in Sweden 2016 and 2017: A retrospective register study. F1000Research, 2018, 7, 1009. | 1.6 | 13 |
| 14 | In-hospital cardiac arrest resuscitation performed by the hospital emergency team: A 6-year retrospective register analysis at Danderyd University Hospital, Sweden. F1000Research, 2018, 7, 1013. | 1.6 | 9 |
| 15 | Continuous surgical multi-level extrapleural block for video-assisted thoracoscopic surgery: a retrospective study assessing its efficacy as pain relief following lobectomy and wedge resection. F1000Research, 2018, 7, 1783. | 1.6 | 8 |
| 16 | Acute hip fracture surgery anaesthetic technique and 30-day mortality in Sweden 2016 and 2017: A retrospective register study. F1000Research, 2018, 7, 1009. | 1.6 | 11 |
| 17 | Pulmonary emboli cardiac arrest with CPR complication: Liver laceration and massive abdominal bleed, a case report. International Journal of Surgery Case Reports, 2017, 31, 24-26. | 0.6 | 4 |
| 18 | Peer review report 1 on "Liposomal bupivacaine infiltration versus femoral nerve block for pain control in total knee arthroplasty: a systematic review and meta-analysis― International Journal of Surgery, 2017, 37, 57. | 2.7 | 0 |

| # | Article | IF | CITATIONS |
|----|--|-----|-----------|
| 19 | Peer review report 3 on "Comparison of femoral nerve block and fascia iliaca block for pain management in total hip arthroplasty: A meta-analysis― International Journal of Surgery, 2017, 37, 488. | 2.7 | 0 |
| 20 | Peer review report 3 on "Liposomal bupivacaine versus interscalene nerve block for pain control after total shoulder arthroplasty: A systematic review and meta-analysis― International Journal of Surgery, 2017, 37, 503. | 2.7 | 0 |
| 21 | High frequency jet ventilation for motion management during ablation procedures, a narrative review. Acta Anaesthesiologica Scandinavica, 2017, 61, 1066-1074. | 1.6 | 28 |
| 22 | The use of high-frequency ventilation during general anaesthesia: an update. F1000Research, 2017, 6, 756. | 1.6 | 26 |
| 23 | Neuraxial opioids as analgesia in labour, caesarean section and hysterectomy: A questionnaire survey in Sweden. F1000Research, 2017, 6, 133. | 1.6 | 10 |
| 24 | Wash-in and wash-out of sevoflurane in a test-lung model: A comparison between Aisys and FLOW-i. F1000Research, 2017, 6, 389. | 1.6 | 12 |
| 25 | Monitoring respiration and oxygen saturation in patients during the first night after elective bariatric surgery: A cohort study. F1000Research, 2017, 6, 735. | 1.6 | 8 |
| 26 | Monitoring respiration and oxygen saturation in patients during the first night after elective bariatric surgery: A cohort study. F1000Research, 2017, 6, 735. | 1.6 | 6 |
| 27 | Anaesthesia for open wrist fracture surgery in adults/elderly. F1000Research, 2017, 6, 1996. | 1.6 | 8 |
| 28 | The decision to delivery interval in emergency caesarean sections: Impact of anaesthetic technique and work shift. F1000Research, 2017, 6, 1977. | 1.6 | 18 |
| 29 | The decision to delivery interval in emergency caesarean sections: Impact of anaesthetic technique and work shift. F1000Research, 2017, 6, 1977. | 1.6 | 13 |
| 30 | The impact of fresh gas flow on wash-in, wash-out time and gas consumption for sevoflurane and desflurane, comparing two anaesthesia machines, a test-lung study F1000Research, 2017, 6, 1997. | 1.6 | 11 |
| 31 | The impact of fresh gas flow on wash-in, wash-out time and gas consumption for sevoflurane and desflurane, comparing two anaesthesia machines, a test-lung study F1000Research, 2017, 6, 1997. | 1.6 | 10 |
| 32 | Pelvic organ prolapse surgery: changes in perioperative management improving hospital pathway. Minerva Obstetrics and Gynecology, 2017, 69, 18-22. | 1.0 | 2 |
| 33 | Neuraxial opioids as analgesia in labour and postoperative after caesarean section and hysterectomy: A questionnaire survey in Sweden. F1000Research, 2017, 6, 133. | 1.6 | 8 |
| 34 | Wash-in and wash-out of sevoflurane in a test-lung model: A comparison between Aisys and FLOW-i. F1000Research, 2017, 6, 389. | 1.6 | 7 |
| 35 | The choice between surgical scrubbing and sterile covering before or after induction of anaesthesia: A prospective study. F1000Research, 2017, 6, 1019. | 1.6 | 2 |
| 36 | The choice between surgical scrubbing and sterile covering before or after induction of anaesthesia: A prospective study. F1000Research, 2017, 6, 1019. | 1.6 | 2 |

| # | Article | IF | CITATIONS |
|----|--|-----|-----------|
| 37 | Portable respiratory polygraphy monitoring of obese mothers the first night after caesarean section with bupivacaine/morphine/fentanyl spinal anaesthesia. F1000Research, 2017, 6, 2062. | 1.6 | 0 |
| 38 | Portable respiratory polygraphy monitoring of obese mothers the first night after caesarean section with bupivacaine/morphine/fentanyl spinal anaesthesia. F1000Research, 2017, 6, 2062. | 1.6 | 0 |
| 39 | Perioperative regional anaesthesia and postoperative longer-term outcomes. F1000Research, 2016, 5, 2501. | 1.6 | 22 |
| 40 | Circulatory collapse following epidural bolus for Caesarean section a profound vasovagal reaction? A case report. International Journal of Surgery Case Reports, 2016, 23, 74-76. | 0.6 | 3 |
| 41 | Pneumothorax following shoulder arthroscopy under combined regional and general anaesthesia—A case report. International Journal of Surgery Case Reports, 2016, 24, 73-76. | 0.6 | 13 |
| 42 | Suspected total spinal in patient having emergent Caesarean section, a case report and literature review. International Journal of Surgery Case Reports, 2016, 28, 173-175. | 0.6 | 2 |
| 43 | Peer review report 2 on "Multimodal analgesia: A systematic review of local NSAIDs for non-ophthalmologic postoperative pain management― International Journal of Surgery, 2016, 25, 314. | 2.7 | 0 |
| 44 | Peer review report 1 on "Intrathecal morphine verse femoral nerve block for pain control in total knee arthroplasty: A meta-analysis from randomized control trials― International Journal of Surgery, 2016, 25, 324. | 2.7 | 0 |
| 45 | Anaesthetics and analgesics; neurocognitive effects, organ protection and cancer reoccurrence an update. International Journal of Surgery, 2016, 34, 41-46. | 2.7 | 11 |
| 46 | Upper extremity nerve block: how can benefit, duration, and safety be improved? An update. F1000Research, 2016, 5, 907. | 1.6 | 29 |
| 47 | Evaluation of the Postoperative Quality of Recovery Scale test and re-test in Swedish among healthy volunteers. F1000Research, 2016, 5, 2549. | 1.6 | 5 |
| 48 | Desflurane and sevoflurane use during low- and minimal-flow anesthesia at fixed vaporizer settings. Minerva Anestesiologica, 2016, 82, 180-5. | 1.0 | 19 |
| 49 | Cognitive testing with the <scp>P</scp> ostâ€Operative <scp>Q</scp> uality of <scp>R</scp> ecovery <scp>S</scp> cale in preâ€surgery cancer patients – a controlled study. Acta Anaesthesiologica Scandinavica, 2015, 59, 763-772. | 1.6 | 8 |
| 50 | Transversus abdominal plane (TAP) block for postoperative pain management: a review. F1000Research, 2015, 4, 1359. | 1.6 | 33 |
| 51 | Transient Left-Sided Paralysis following Robotic-Assisted Laparoscopic Uteropexy. Case Reports in Anesthesiology, 2015, 2015, 1-3. | 0.4 | 1 |
| 52 | Post-operative nausea and vomiting: Update on predicting the probability and ways to minimize its occurrence, with focus on ambulatory surgery. International Journal of Surgery, 2015, 15, 100-106. | 2.7 | 74 |
| 53 | Preoperative platelet count and volume could not help predict PONV in women undergoing breast cancer surgery: A prospective cohort study. International Journal of Surgery, 2015, 18, 128-131. | 2.7 | 3 |
| 54 | Peer review report 2 on "ASA class is a reliable independent predictor of medical complications and mortality following surgery – An observational study― International Journal of Surgery, 2015, 13, S44. | 2.7 | 1 |

| # | Article | IF | CITATIONS |
|----|--|-----|-----------|
| 55 | Subacute complications during recovery from severe traumatic brain injury: frequency and associations with outcome. BMJ Open, 2015, 5, e007208-e007208. | 1.9 | 40 |
| 56 | Adjunct High Frequency Transcutaneous Electric Stimulation (TENS) for Postoperative Pain Management during Weaning from Epidural Analgesia Following Colon Surgery: Results from a Controlled Pilot Study. Pain Management Nursing, 2015, 16, 944-950. | 0.9 | 6 |
| 57 | Perioperative management in order to minimise postoperative delirium and postoperative cognitive dysfunction: Results from a Swedish web-based survey. Annals of Medicine and Surgery, 2014, 3, 100-107. | 1.1 | 27 |
| 58 | Routines for reducing the occurrence of emergence agitation during awakening in children, a national survey. SpringerPlus, 2014, 3, 572. | 1.2 | 11 |
| 59 | Suspected Transfusion Related Acute Lung Injury Improving following Administration of Tranexamic Acid: A Case Report. Case Reports in Anesthesiology, 2014, 2014, 1-3. | 0.4 | 3 |
| 60 | Pain Management in Ambulatory Surgery—A Review. Pharmaceuticals, 2014, 7, 850-865. | 3.8 | 29 |
| 61 | Cognitive recovery after ambulatory anaesthesia based on desflurane or propofol: a prospective randomised study. Acta Anaesthesiologica Scandinavica, 2014, 58, 1111-1120. | 1.6 | 17 |
| 62 | Outcomes, Measures and Recovery After Ambulatory Surgery and Anaesthesia: A review. Current Anesthesiology Reports, 2014, 4, 334-341. | 2.0 | 6 |
| 63 | AAI-guided anaesthesia is associated with lower incidence of 24-hÂMMSEÂ<Â25 and may impact the IL-6 response. International Journal of Surgery, 2014, 12, 290-295. | 2.7 | 6 |
| 64 | Preoperative centre improving patients' satisfaction?. International Journal of Surgery, 2013, 11, 430-431. | 2.7 | 0 |
| 65 | Day surgery, variations in routines and practices a questionnaire survey. International Journal of Surgery, 2013, 11, 178-182. | 2.7 | 20 |
| 66 | Post-operative Quality of Recovery Scale: the impact of assessment method on cognitive recovery. Acta Anaesthesiologica Scandinavica, 2013, 57, 1308-1312. | 1.6 | 10 |
| 67 | Lornoxicam: pharmacology and usefulness to treat acute postoperative and musculoskeletal pain a narrative review. Expert Opinion on Pharmacotherapy, 2013, 14, 1679-1694. | 1.8 | 18 |
| 68 | Non-opioid analgesics for pain management following ambulatory surgery: a review. Minerva Anestesiologica, 2013, 79, 1077-87. | 1.0 | 21 |
| 69 | Brief review: Theory and practice of minimal fresh gas flow anesthesia. Canadian Journal of Anaesthesia, 2012, 59, 785-797. | 1.6 | 56 |
| 70 | Assessing recovery after ambulatory anaesthesia, measures of resumption of activities of daily living. Current Opinion in Anaesthesiology, 2011, 24, 601-604. | 2.0 | 16 |
| 71 | Preoperative single-dose intravenous dexamethasone during ambulatory surgery: update around the benefit versus risk. Current Opinion in Anaesthesiology, 2010, 23, 682-686. | 2.0 | 23 |
| 72 | Anaesthetic gas monitoring comparison between two side-stream monitors. Journal of Clinical Monitoring and Computing, 2010, 24, 169-172. | 1.6 | 1 |

| # | Article | IF | CITATIONS |
|----|--|-----|-----------|
| 73 | Clinical Performance of a Novel Main-Stream Anaesthetic End-Tidal Gas Monitors during Routine Low Flow Anaesthesia. Journal of Anesthesia & Clinical Research, 2010, 1, . | 0.1 | 1 |
| 74 | The effects of fresh gas flow on the amount of sevoflurane vaporized during 1 minimum alveolar concentration anaesthesia for day surgery: a clinical study. Acta Anaesthesiologica Scandinavica, 2007, 51, 290-293. | 1.6 | 23 |
| 75 | Desflurane vs. sevoflurane as the main inhaled anaesthetic for spontaneous breathing via a laryngeal mask for varicose vein day surgery: a prospective randomized study. Acta Anaesthesiologica Scandinavica, 2006, 50, 549-552. | 1.6 | 34 |
| 76 | Operating room nitrous oxide trace concentrations: a clinical study in ambulatory surgery. Ambulatory Surgery, 2005, 12, 23-26. | 0.1 | 3 |
| 77 | Apnea and bradypnea in patients receiving epidural bupivacaine-morphine for postoperative pain relief as assessed by a new monitoring method. Journal of Clinical Anesthesia, 2002, 14, 129-134. | 1.6 | 13 |
| 78 | Intrathecal Sufentanil, Fentanyl, or Placebo Added to Bupivacaine for Cesarean Section. Anesthesia and Analgesia, 1997, 85, 1288-1293. | 2.2 | 92 |
| 79 | Laparoscopic cholecystectomy perioperative management: an update. Ambulatory Anesthesia, 0, , 53. | 0.0 | 1 |
| 80 | Brace versus cast following surgical treatment of distal radial fracture: a prospective randomised study comparing quality of recovery. F1000Research, 0, 10, 336. | 1.6 | 0 |