Guorong Fan

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/521123/publications.pdf

Version: 2024-02-01

1163117 1058476 14 268 8 14 citations h-index g-index papers 14 14 14 286 docs citations times ranked citing authors all docs

#	Article	IF	CITATIONS
1	Separation and determination of coumarins in Fructus cnidii extracts by pressurized capillary electrochromatography using a packed column with a monolithic outlet frit. Journal of Pharmaceutical and Biomedical Analysis, 2009, 50, 695-702.	2.8	54
2	Separation and quantitative analysis of coumarin compounds from Angelica dahurica (Fisch. ex) Tj ETQq0 0 0 rgBT and Biomedical Analysis, 2006, 41, 105-116.	Overlock	10 Tf 50 70 52
3	Fingerprint analysis of the fruits of Cnidium monnieri extract by high-performance liquid chromatography–diode array detection–electrospray ionization tandem mass spectrometry. Journal of Pharmaceutical and Biomedical Analysis, 2007, 43, 926-936.	2.8	49
4	Optimal extraction and fingerprint analysis of Cnidii fructus by accelerated solvent extraction and high performance liquid chromatographic analysis with photodiode array and mass spectrometry detections. Food Chemistry, 2013, 141, 1962-1971.	8.2	35
5	Microdialysis combined with liquid chromatography–tandem mass spectrometry for the determination of levo-tetrahydropalmatine in the rat striatum. Journal of Pharmaceutical and Biomedical Analysis, 2012, 64-65, 1-7.	2.8	14
6	Development and validation of a rapid HPLC method for the determination of cefdinir in beagle dog plasma integrated with an automatic on-line solid-phase extraction following protein precipitation in the 96-well plate format. Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences, 2012, 895-896, 83-88.	2.3	13
7	Determination of ulifloxacin, the active metabolite of prulifloxacin, in human plasma by a 96-well format solid-phase extraction and capillary zone electrophoresis. Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences, 2008, 872, 172-176.	2.3	11
8	Sodium desoxycholateâ€assisted capillary electrochromatography with methacrylate esterâ€based monolithic column on fast separation and determination of coumarin analogs in <i><scp>A</scp>ngelica dahurica</i>	2.4	11
9	High-Throughput Determination of Sodium Danshensu in Beagle Dogs by the LCMS/MS Method, Employing Liquid-Liquid Extraction Based on 96-Well Format Plates. Molecules, 2017, 22, 667.	3.8	9
10	Application and optimization of organic–inorganic hybrid monolithic capillary electrochromatography for in vivo cefdinir determination with microdialysis. Journal of Separation Science, 2016, 39, 440-449.	2.5	7
11	Simultaneous determination of the potent anti-tuberculosis regimen—Pyrazinamide, ethambutol, protionamide, clofazimine in beagle dog plasma using LC–MS/MS method coupled with 96-well format plate. Journal of Pharmaceutical and Biomedical Analysis, 2019, 168, 44-54.	2.8	6
12	Fast and direct determination of catechol-3, 6-bis(methyleiminodiacetic acid) prototype in beagle dog plasma using liquid chromatography tandem mass spectrometry: A simplified and high throughput in-vivo method for the metal chelator. Journal of Chromatography A, 2019, 1596, 84-95.	3.7	3
13	Identification of 3′,4′-Dimethoxy Flavonol-3-β-d-Glucopyranoside Metabolites in Rats by Liquid Chromatography-Electrospray Ionization Ion Trap Mass Spectrometry. Molecules, 2016, 21, 470.	3.8	2
14	A fast and accurate method for the pharmacokinetic research of four coumarin analogs in <i>Fructus cnidii</i> using capillary electroâ€chromatography with a methacrylate esterâ€based monolithic column. Electrophoresis, 2017, 38, 3036-3047.	2.4	2