

Guorong Fan

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/521123/publications.pdf>

Version: 2024-02-01

14
papers

268
citations

1163117

8
h-index

1058476

14
g-index

14
all docs

14
docs citations

14
times ranked

286
citing authors

#	ARTICLE	IF	CITATIONS
1	Fast and direct determination of catechol-3, 6-bis(methyleiminodiacetic acid) prototype in beagle dog plasma using liquid chromatography tandem mass spectrometry: A simplified and high throughput in-vivo method for the metal chelator. <i>Journal of Chromatography A</i> , 2019, 1596, 84-95.	3.7	3
2	Simultaneous determination of the potent anti-tuberculosis regimenâ€”Pyrazinamide, ethambutol, protionamide, clofazimine in beagle dog plasma using LCâ€”MS/MS method coupled with 96-well format plate. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , 2019, 168, 44-54.	2.8	6
3	A fast and accurate method for the pharmacokinetic research of four coumarin analogs in <i>Fructus cnidii</i> using capillary electrochromatography with a methacrylate esterâ€”based monolithic column. <i>Electrophoresis</i> , 2017, 38, 3036-3047.	2.4	2
4	High-Throughput Determination of Sodium Danshensu in Beagle Dogs by the LCMS/MS Method, Employing Liquid-Liquid Extraction Based on 96-Well Format Plates. <i>Molecules</i> , 2017, 22, 667.	3.8	9
5	Identification of 3,4-Dimethoxy Flavonol-3- β -D-Glucopyranoside Metabolites in Rats by Liquid Chromatography-Electrospray Ionization Ion Trap Mass Spectrometry. <i>Molecules</i> , 2016, 21, 470.	3.8	2
6	Application and optimization of organicâ€”inorganic hybrid monolithic capillary electrochromatography for in vivo cefdinir determination with microdialysis. <i>Journal of Separation Science</i> , 2016, 39, 440-449.	2.5	7
7	Optimal extraction and fingerprint analysis of <i>Cnidii fructus</i> by accelerated solvent extraction and high performance liquid chromatographic analysis with photodiode array and mass spectrometry detections. <i>Food Chemistry</i> , 2013, 141, 1962-1971.	8.2	35
8	Sodium desoxycholateâ€”assisted capillary electrochromatography with methacrylate esterâ€”based monolithic column on fast separation and determination of coumarin analogs in <i>Angelica dahurica</i> extract. <i>Electrophoresis</i> , 2012, 33, 2884-2891.	2.4	11
9	Development and validation of a rapid HPLC method for the determination of cefdinir in beagle dog plasma integrated with an automatic on-line solid-phase extraction following protein precipitation in the 96-well plate format. <i>Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences</i> , 2012, 895-896, 83-88.	2.3	13
10	Microdialysis combined with liquid chromatographyâ€”tandem mass spectrometry for the determination of levo-tetrahydropalmatine in the rat striatum. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , 2012, 64-65, 1-7.	2.8	14
11	Separation and determination of coumarins in <i>Fructus cnidii</i> extracts by pressurized capillary electrochromatography using a packed column with a monolithic outlet frit. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , 2009, 50, 695-702.	2.8	54
12	Determination of ulifloxacin, the active metabolite of prulifloxacin, in human plasma by a 96-well format solid-phase extraction and capillary zone electrophoresis. <i>Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences</i> , 2008, 872, 172-176.	2.3	11
13	Fingerprint analysis of the fruits of <i>Cnidium monnieri</i> extract by high-performance liquid chromatographyâ€”diode array detectionâ€”electrospray ionization tandem mass spectrometry. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , 2007, 43, 926-936.	2.8	49
14	Separation and quantitative analysis of coumarin compounds from <i>Angelica dahurica</i> (Fisch. ex) Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50 22 and <i>Biomedical Analysis</i> , 2006, 41, 105-116.	2.8	52