## Hongbing Mei

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/5271866/publications.pdf

Version: 2024-02-01

687363 713466 25 477 13 21 h-index citations g-index papers 25 25 25 734 docs citations times ranked citing authors all docs

#	Article	IF	CITATIONS
1	Over-expression of long noncoding RNA BANCR inhibits malignant phenotypes of human bladder cancer. Journal of Experimental and Clinical Cancer Research, 2016, 35, 125.	8.6	64
2	Decreased expression of LncRNA MIR31HG in human bladder cancer. Cancer Biomarkers, 2016, 17, 231-236.	1.7	44
3	Overexpression of CRNDE promotes the progression of bladder cancer. Biomedicine and Pharmacotherapy, 2018, 99, 638-644.	5.6	33
4	Role of long noncoding RNA UCA1 as a common molecular marker for lymph node metastasis and prognosis in various cancers: a meta-analysis. Oncotarget, 2017, 8, 1937-1943.	1.8	33
5	CMTM3 Inhibits Human Testicular Cancer Cell Growth through Inducing Cell-Cycle Arrest and Apoptosis. PLoS ONE, 2014, 9, e88965.	2.5	31
6	Potential Role of IncRNA H19 as a Cancer Biomarker in Human Cancers Detection and Diagnosis: A Pooled Analysis Based on 1585 Subjects. BioMed Research International, 2019, 2019, 1-11.	1.9	30
7	rs11614913 polymorphism in miRNA-196a2 and cancer risk: an updated meta-analysis. OncoTargets and Therapy, 2018, Volume 11, 1121-1139.	2.0	25
8	Patientâ€derived renal cell carcinoma organoids for personalized cancer therapy. Clinical and Translational Medicine, 2022, 12, .	4.0	24
9	CRISPR-Cas13-Mediated Knockdown of IncRNA-GACAT3 Inhibited Cell Proliferation and Motility, and Induced Apoptosis by Increasing p21, Bax, and E-Cadherin Expression in Bladder Cancer. Frontiers in Molecular Biosciences, 2020, 7, 627774.	3.5	22
10	<p>Loss of <em>KCNJ15</em> expression promotes malignant phenotypes and correlates with poor prognosis in renal carcinoma</p> . Cancer Management and Research, 2019, Volume 11, 1211-1220.	1.9	21
11	YTHDF2 is a Potential Biomarker and Associated with Immune Infiltration in Kidney Renal Clear Cell Carcinoma. Frontiers in Pharmacology, 2021, 12, 709548.	3.5	21
12	Association of polymorphisms in interleukin-8 gene with cancer risk: a meta-analysis of 22 case–control studies. OncoTargets and Therapy, 2016, 9, 3727.	2.0	19
13	Targeted Methylation of the LncRNA NEAT1 Suppresses Malignancy of Renal Cell Carcinoma. Frontiers in Cell and Developmental Biology, 2021, 9, 777349.	3.7	16
14	NDNF inhibits the migration and invasion of human renal cancer cells through epithelial‑mesenchymal transition. Oncology Letters, 2019, 17, 2969-2975.	1.8	15
15	Long noncoding RNA MALAT1 acts as a potential biomarker in cancer diagnosis and detection: a meta-analysis. Biomarkers in Medicine, 2019, 13, 45-54.	1.4	15
16	Patientâ€Derived Upper Tract Urothelial Carcinoma Organoids as a Platform for Drug Screening. Advanced Science, 2022, 9, e2103999.	11.2	12
17	Knockdown of long noncoding RNA FGFR3- AS1 induces cell proliferation inhibition, apoptosis and motility reduction in bladder cancer. Cancer Biomarkers, 2018, 21, 277-285.	1.7	11
18	Current Advances on the Important Roles of Enhancer RNAs in Gene Regulation and Cancer. BioMed Research International, 2018, 2018, 1-6.	1.9	10

#	Article	lF	CITATIONS
19	Association between two interleukin-2 gene polymorphisms and cancer susceptibility: a meta-analysis. OncoTargets and Therapy, 2016, 9, 2181.	2.0	8
20	High expression of enhancer RNA MARC1 or its activation by DHT is associated with the malignant behavior in bladder cancer. Experimental Cell Research, 2018, 370, 303-311.	2.6	7
21	A synthetic targeted RNA demethylation system based on CRISPRâ€Cas13b inhibits bladder cancer progression. Clinical and Translational Medicine, 2022, 12, e734.	4.0	5
22	Lentivirus-mediated shRNA targeting <em>MUTYH</em> inhibits malignant phenotypes of bladder cancer SW780 cells. OncoTargets and Therapy, 2018, Volume 11, 6101-6109.	2.0	4
23	Association of Polymorphisms in Intercellular Adhesion Molecule 1 (ICAM-1) Gene with Cancer Susceptibility: A Meta-Analysis of 14 Case-Control Studies. Medical Science Monitor, 2016, 22, 569-579.	1.1	3
24	Long non-coding RNA BRAF activated non-coding RNA as a potential novel molecular biomarker for metastasis and prognosis in various cancers. Minerva Medica, 2017, 108, 293-298.	0.9	3
25	The Effect of Single-port Transvesical Laparoscopic Radical Prostatectomy on Erectile Function and Urinary Continence Compared to Intrafascial Endoscopic Extraperitoneal Radical Prostatectomy. Urology Journal, 2020, 17, 592-596.	0.4	1