

Gregory H Tesch

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/5332977/publications.pdf>

Version: 2024-02-01

93
papers

6,786
citations

53794

45
h-index

60623

81
g-index

95
all docs

95
docs citations

95
times ranked

7287
citing authors

#	ARTICLE	IF	CITATIONS
1	WNT1-inducible signaling pathway protein 1 regulates kidney inflammation through the NF- κ B pathway. <i>Clinical Science</i> , 2022, 136, 29-44.	4.3	4
2	Mice with Established Diabetes Show Increased Susceptibility to Renal Ischemia/Reperfusion Injury. <i>American Journal of Pathology</i> , 2022, 192, 441-453.	3.8	2
3	ASK1 is a novel molecular target for preventing aminoglycoside-induced hair cell death. <i>Journal of Molecular Medicine</i> , 2022, 100, 797-813.	3.9	3
4	Review article: Have emergency department time-based targets influenced patient care? A systematic review of qualitative literature. <i>EMA - Emergency Medicine Australasia</i> , 2021, 33, 202-213.	1.1	4
5	c-Jun Amino Terminal Kinase Signaling Promotes Aristolochic Acid-Induced Acute Kidney Injury. <i>Frontiers in Physiology</i> , 2021, 12, 599114.	2.8	6
6	WNT1-inducible signaling pathway protein 1 regulates the development of kidney fibrosis through the TGF- β 1 pathway. <i>FASEB Journal</i> , 2020, 34, 14507-14520.	0.5	9
7	Targeting apoptosis signaling-regulating kinase 1 in acute and chronic kidney disease. <i>Anatomical Record</i> , 2020, 303, 2553-2560.	1.4	8
8	Novel mineralocorticoid receptor mechanisms regulate cardiac tissue inflammation in male mice. <i>Journal of Endocrinology</i> , 2020, 246, 123-134.	2.6	6
9	Combined inhibition of CCR2 and ACE provides added protection against progression of diabetic nephropathy in <i>Nos3</i> -deficient mice. <i>American Journal of Physiology - Renal Physiology</i> , 2019, 317, F1439-F1449.	2.7	8
10	Establishing equivalent diabetes in male and female <i>Nos3</i> -deficient mice results in a comparable onset of diabetic kidney injury. <i>Physiological Reports</i> , 2019, 7, e14197.	1.7	9
11	Pharmacological inhibition of protease-activated receptor-2 reduces crescent formation in rat nephrotoxic serum nephritis. <i>Clinical and Experimental Pharmacology and Physiology</i> , 2019, 46, 456-464.	1.9	8
12	Proximal tubular epithelial cells preferentially endocytose covalently modified albumin compared to native albumin. <i>Nephrology</i> , 2019, 24, 121-126.	1.6	0
13	Matrix metalloproteinase-12 deficiency attenuates experimental crescentic anti-glomerular basement membrane glomerulonephritis. <i>Nephrology</i> , 2018, 23, 183-189.	1.6	13
14	Cyclophilin D promotes tubular cell damage and the development of interstitial fibrosis in the obstructed kidney. <i>Clinical and Experimental Pharmacology and Physiology</i> , 2018, 45, 250-260.	1.9	18
15	Reduced tubular degradation of glomerular filtered plasma albumin is a common feature in acute and chronic kidney disease. <i>Clinical and Experimental Pharmacology and Physiology</i> , 2018, 45, 241-249.	1.9	5
16	ASK1 inhibitor treatment suppresses p38/JNK signalling with reduced kidney inflammation and fibrosis in rat crescentic glomerulonephritis. <i>Journal of Cellular and Molecular Medicine</i> , 2018, 22, 4522-4533.	3.6	47
17	miR-378 reduces mesangial hypertrophy and kidney tubular fibrosis via MAPK signalling. <i>Clinical Science</i> , 2017, 131, 411-423.	4.3	27
18	Diabetic nephropathy "is this an immune disorder?". <i>Clinical Science</i> , 2017, 131, 2183-2199.	4.3	182

#	ARTICLE	IF	CITATIONS
19	Inhibition of Spleen Tyrosine Kinase Reduces Renal Allograft Injury in a Rat Model of Acute Antibody-Mediated Rejection in Sensitized Recipients. <i>Transplantation</i> , 2017, 101, e240-e248.	1.0	10
20	Mineralocorticoid Receptor Signaling as a Therapeutic Target for Renal and Cardiac Fibrosis. <i>Frontiers in Pharmacology</i> , 2017, 8, 313.	3.5	74
21	ASK1: a new therapeutic target for kidney disease. <i>American Journal of Physiology - Renal Physiology</i> , 2016, 311, F373-F381.	2.7	53
22	Myeloid cell-mediated renal injury in rapidly progressive glomerulonephritis depends upon spleen tyrosine kinase. <i>Journal of Pathology</i> , 2016, 238, 10-20.	4.5	19
23	Cardiac Tissue Injury and Remodeling Is Dependent Upon MR Regulation of Activation Pathways in Cardiac Tissue Macrophages. <i>Endocrinology</i> , 2016, 157, 3213-3223.	2.8	47
24	Earlier onset of diabetes-induced adverse cardiac remodeling in female compared to male mice. <i>Obesity</i> , 2015, 23, 1166-1177.	3.0	19
25	ASK1 Inhibitor Halts Progression of Diabetic Nephropathy in <i>Nos3</i> -Deficient Mice. <i>Diabetes</i> , 2015, 64, 3903-3913.	0.6	76
26	Spleen tyrosine kinase contributes to acute renal allograft rejection in the rat. <i>International Journal of Experimental Pathology</i> , 2015, 96, 54-62.	1.3	7
27	Suppression of Rapidly Progressive Mouse Glomerulonephritis with the Non-Steroidal Mineralocorticoid Receptor Antagonist BR-4628. <i>PLoS ONE</i> , 2015, 10, e0145666.	2.5	12
28	Myeloid Mineralocorticoid Receptor Activation Contributes to Progressive Kidney Disease. <i>Journal of the American Society of Nephrology: JASN</i> , 2014, 25, 2231-2240.	6.1	60
29	ASK1/p38 signaling in renal tubular epithelial cells promotes renal fibrosis in the mouse obstructed kidney. <i>American Journal of Physiology - Renal Physiology</i> , 2014, 307, F1263-F1273.	2.7	87
30	Deletion of bone-marrow-derived receptor for AGEs (RAGE) improves renal function in an experimental mouse model of diabetes. <i>Diabetologia</i> , 2014, 57, 1977-1985.	6.3	26
31	Role of macrophages in the fibrotic phase of rat crescentic glomerulonephritis. <i>American Journal of Physiology - Renal Physiology</i> , 2013, 304, F1043-F1053.	2.7	63
32	Design and pharmacology of a highly specific dual FMS and KIT kinase inhibitor. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2013, 110, 5689-5694.	7.1	82
33	Inflammation in Diabetic Nephropathy. <i>Mediators of Inflammation</i> , 2012, 2012, 1-12.	3.0	330
34	Macrophage Mineralocorticoid Receptor Signaling Plays a Key Role in Aldosterone-Independent Cardiac Fibrosis. <i>Endocrinology</i> , 2012, 153, 3416-3425.	2.8	102
35	Aldosterone Induces Kidney Fibroblast Proliferation via Activation of Growth Factor Receptors and PI3K/MAPK Signalling. <i>Nephron Experimental Nephrology</i> , 2012, 120, e115-e122.	2.2	43
36	TGF- β 1-activated kinase-1 regulates inflammation and fibrosis in the obstructed kidney. <i>American Journal of Physiology - Renal Physiology</i> , 2011, 300, F1410-F1421.	2.7	92

#	ARTICLE	IF	CITATIONS
37	c-fms blockade reverses glomerular macrophage infiltration and halts development of crescentic anti-GBM glomerulonephritis in the rat. <i>Laboratory Investigation</i> , 2011, 91, 978-991.	3.7	54
38	Recent insights into diabetic renal injury from the <i>db/db</i> mouse model of type 2 diabetic nephropathy. <i>American Journal of Physiology - Renal Physiology</i> , 2011, 300, F301-F310.	2.7	120
39	Evaluation of JNK Blockade as an Early Intervention Treatment for Type 1 Diabetic Nephropathy in Hypertensive Rats. <i>American Journal of Nephrology</i> , 2011, 34, 337-346.	3.1	34
40	Successes Achieved and Challenges Ahead in Translating Biomarkers into Clinical Applications. <i>AAPS Journal</i> , 2010, 12, 243-253.	4.4	26
41	Lymphocytes promote albuminuria, but not renal dysfunction or histological damage in a mouse model of diabetic renal injury. <i>Diabetologia</i> , 2010, 53, 1772-1782.	6.3	61
42	Review: Serum and urine biomarkers of kidney disease: A pathophysiological perspective. <i>Nephrology</i> , 2010, 15, 609-616.	1.6	107
43	Macrophages and Diabetic Nephropathy. <i>Seminars in Nephrology</i> , 2010, 30, 290-301.	1.6	119
44	Lefty antagonises TGF- β 1 induced epithelial-mesenchymal transition in tubular epithelial cells. <i>Biochemical and Biophysical Research Communications</i> , 2010, 393, 855-859.	2.1	17
45	Deletion of Mineralocorticoid Receptors From Macrophages Protects Against Deoxycorticosterone/Salt-Induced Cardiac Fibrosis and Increased Blood Pressure. <i>Hypertension</i> , 2009, 54, 537-543.	2.7	272
46	Role of MKK3-p38 MAPK signalling in the development of type 2 diabetes and renal injury in obese <i>db/db</i> mice. <i>Diabetologia</i> , 2009, 52, 347-358.	6.3	100
47	Antibody blockade of c-fms suppresses the progression of inflammation and injury in early diabetic nephropathy in obese <i>db/db</i> mice. <i>Diabetologia</i> , 2009, 52, 1669-1679.	6.3	85
48	Blockade of the c-Jun amino terminal kinase prevents crescent formation and halts established anti-GBM glomerulonephritis in the rat. <i>Laboratory Investigation</i> , 2009, 89, 470-484.	3.7	58
49	MKK3 signalling plays an essential role in leukocyte-mediated pancreatic injury in the multiple low-dose streptozotocin model. <i>Laboratory Investigation</i> , 2008, 88, 398-407.	3.7	20
50	c-Jun amino terminal kinase 1 deficient mice are protected from streptozotocin-induced islet injury. <i>Biochemical and Biophysical Research Communications</i> , 2008, 366, 710-716.	2.1	17
51	In vivo visualization of albumin degradation in the proximal tubule. <i>Kidney International</i> , 2008, 74, 1480-1486.	5.2	33
52	A Pathogenic Role for c-Jun Amino-Terminal Kinase Signaling in Renal Fibrosis and Tubular Cell Apoptosis. <i>Journal of the American Society of Nephrology: JASN</i> , 2007, 18, 472-484.	6.1	152
53	MIF in the Pathogenesis of Kidney Disease. , 2007, , 153-168.		0
54	A pathogenic role for JNK signaling in experimental anti-GBM glomerulonephritis. <i>Kidney International</i> , 2007, 72, 698-708.	5.2	61

#	ARTICLE	IF	CITATIONS
55	MKK3-p38 signaling promotes apoptosis and the early inflammatory response in the obstructed mouse kidney. <i>American Journal of Physiology - Renal Physiology</i> , 2007, 293, F1556-F1563.	2.7	51
56	ROLE OF MACROPHAGES IN COMPLICATIONS OF TYPE 2 DIABETES. <i>Clinical and Experimental Pharmacology and Physiology</i> , 2007, 34, 1016-1019.	1.9	116
57	Human peritoneal mesothelial cells isolated from spent dialysate fluid maintain contaminating macrophages via production of macrophage colony stimulating factor. <i>Nephrology</i> , 2007, 12, 160-165.	1.6	3
58	Rodent models of streptozotocin-induced diabetic nephropathy (Methods in Renal Research). <i>Nephrology</i> , 2007, 12, 261-266.	1.6	386
59	Quantification of renal pathology by image analysis (Methods in Renal Research). <i>Nephrology</i> , 2007, 12, 553-558.	1.6	148
60	Monocyte chemoattractant protein-1-induced tissue inflammation is critical for the development of renal injury but not type 2 diabetes in obese db/db mice. <i>Diabetologia</i> , 2007, 50, 471-480.	6.3	222
61	Monocyte chemoattractant protein-1 has prosclerotic effects both in a mouse model of experimental diabetes and in vitro in human mesangial cells. <i>Diabetologia</i> , 2007, 51, 198-207.	6.3	73
62	Treatment of Tissue Sections for <i>In Situ</i> Hybridization. , 2006, 326, 1-8.		8
63	Monocyte chemoattractant protein-1 promotes the development of diabetic renal injury in streptozotocin-treated mice. <i>Kidney International</i> , 2006, 69, 73-80.	5.2	378
64	Recent Insights into Experimental Mouse Models of Diabetic Nephropathy. <i>Nephron Experimental Nephrology</i> , 2006, 104, e57-e62.	2.2	32
65	Intercellular Adhesion Molecule-1 Deficiency Is Protective against Nephropathy in Type 2 Diabetic db/db Mice. <i>Journal of the American Society of Nephrology: JASN</i> , 2005, 16, 1711-1722.	6.1	247
66	Kidney expression of glutathione peroxidase-1 is not protective against streptozotocin-induced diabetic nephropathy. <i>American Journal of Physiology - Renal Physiology</i> , 2005, 289, F544-F551.	2.7	60
67	The Role of p38 ^{Î±} Mitogen-Activated Protein Kinase Activation in Renal Fibrosis. <i>Journal of the American Society of Nephrology: JASN</i> , 2004, 15, 370-379.	6.1	184
68	Macrophages in streptozotocin-induced diabetic nephropathy: potential role in renal fibrosis. <i>Nephrology Dialysis Transplantation</i> , 2004, 19, 2987-2996.	0.7	171
69	Macrophage accumulation and renal fibrosis are independent of macrophage migration inhibitory factor in mouse obstructive nephropathy. <i>Nephrology</i> , 2004, 9, 278-287.	1.6	10
70	Macrophages in mouse type 2 diabetic nephropathy: Correlation with diabetic state and progressive renal injury. <i>Kidney International</i> , 2004, 65, 116-128.	5.2	461
71	Abnormal p38 mitogen-activated protein kinase signalling in human and experimental diabetic nephropathy. <i>Diabetologia</i> , 2004, 47, 1210-1222.	6.3	181
72	Induction of MIF synthesis and secretion by tubular epithelial cells: A novel action of angiotensin II. <i>Kidney International</i> , 2003, 63, 1265-1275.	5.2	49

#	ARTICLE	IF	CITATIONS
73	Role of interleukin-10 in rat mesangioproliferative glomerulonephritis. <i>Nephrology</i> , 2003, 8, 33-41.	1.6	5
74	Interferon-gamma induces macrophage migration inhibitory factor synthesis and secretion by tubular epithelial cells. <i>Nephrology</i> , 2003, 8, 156-161.	1.6	19
75	Blockade of p38 β MAPK Ameliorates Acute Inflammatory Renal Injury in Rat Anti-GBM Glomerulonephritis. <i>Journal of the American Society of Nephrology: JASN</i> , 2003, 14, 338-351.	6.1	101
76	Heterogeneity of antigen expression explains controversy over glomerular macrophage accumulation in mouse glomerulonephritis. <i>Nephrology Dialysis Transplantation</i> , 2003, 18, 178-181.	0.7	38
77	Long-term anti-glomerular basement membrane disease in the rat: a model of chronic glomerulonephritis with nephrosis, hypertension and progressive renal failure. <i>Nephrology</i> , 2002, 7, 145-154.	1.6	1
78	Macrophage accumulation at a site of renal inflammation is dependent on the M-CSF/c-fms pathway. <i>Journal of Leukocyte Biology</i> , 2002, 72, 530-7.	3.3	54
79	LF15-0195 prevents the induction and inhibits the progression of rat anti-GBM disease. <i>Kidney International</i> , 2001, 60, 1354-1365.	5.2	14
80	Combined interleukin 1 and tumour necrosis factor alpha blockade in rat crescentic anti-glomerular basement membrane glomerulonephritis. <i>Nephrology</i> , 2001, 6, 214-220.	1.6	2
81	Costimulation by B7-1 and B7-2 Is Required for Autoimmune Disease in MRL-Fas lpr Mice. <i>Journal of Immunology</i> , 2000, 164, 6046-6056.	0.8	75
82	Up-regulation of the tumour-associated marker CD44V6 in experimental kidney disease. <i>Clinical and Experimental Immunology</i> , 2000, 121, 523-532.	2.6	8
83	CD44-mediated neutrophil apoptosis in the rat. <i>Kidney International</i> , 2000, 58, 1920-1930.	5.2	40
84	Monocyte Chemoattractant Protein 1-Dependent Leukocytic Infiltrates Are Responsible for Autoimmune Disease in Mrl-Fas lpr Mice. <i>Journal of Experimental Medicine</i> , 1999, 190, 1813-1824.	8.5	287
85	Authors' reply:. <i>American Journal of Kidney Diseases</i> , 1999, 34, 765-767.	1.9	2
86	Monocyte chemoattractant protein-1 promotes macrophage-mediated tubular injury, but not glomerular injury, in nephrotoxic serum nephritis. <i>Journal of Clinical Investigation</i> , 1999, 103, 73-80.	8.2	238
87	Interleukin-10 differentially modulates MHC class II expression by mesangial cells and macrophages in vitro and in vivo. <i>Immunology</i> , 1998, 94, 72-78.	4.4	42
88	Do macrophages participate in mesangial cell proliferation?. <i>Nephrology</i> , 1997, 3, 501-507.	1.6	2
89	Effect of interleukin-10 treatment on crescentic glomerulonephritis in rats. <i>Kidney International</i> , 1997, 51, 1809-1817.	5.2	29
90	A novel method of microwave treatment for detection of cytoplasmic and nuclear antigens by flow cytometry. <i>Journal of Immunological Methods</i> , 1996, 190, 1-10.	1.4	37

#	ARTICLE	IF	CITATIONS
91	EGF and EGF-receptor expression in rat anti-Thy-1 mesangial proliferative nephritis. <i>Nephrology</i> , 1995, 1, 83-93.	1.6	4
92	DEOXYSPERGUALIN SUPPRESSES LOCAL MACROPHAGE PROLIFERATION IN RAT RENAL ALLOGRAFT REJECTION. <i>Transplantation</i> , 1994, 58, 596-601.	1.0	58
93	Effects of free and bound insulin-like growth factors on proteoglycan metabolism in articular cartilage explants. <i>Journal of Orthopaedic Research</i> , 1992, 10, 14-22.	2.3	53