

Saaid Safieddine

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/5343381/publications.pdf>

Version: 2024-02-01

33
papers

3,243
citations

218677

26
h-index

345221

36
g-index

37
all docs

37
docs citations

37
times ranked

2846
citing authors

#	ARTICLE	IF	CITATIONS
1	Viral transfer of mini-otoferlins partially restores the fast component of exocytosis and uncovers ultrafast endocytosis in auditory hair cells of otoferlin knock-out mice. <i>Journal of Neuroscience</i> , 2019, 39, 1550-18.	3.6	28
2	Dual AAV-mediated gene therapy restores hearing in a DFNB9 mouse model. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2019, 116, 4496-4501.	7.1	162
3	Hearing Protection, Restoration, and Regeneration: An Overview of Emerging Therapeutics for Inner Ear and Central Hearing Disorders. <i>Otology and Neurotology</i> , 2019, 40, 559-570.	1.3	68
4	Hair Cell Afferent Synapses: Function and Dysfunction. <i>Cold Spring Harbor Perspectives in Medicine</i> , 2019, 9, a033175.	6.2	20
5	Mechanotransduction is required for establishing and maintaining mature inner hair cells and regulating efferent innervation. <i>Nature Communications</i> , 2018, 9, 4015.	12.8	54
6	Clarin-1 gene transfer rescues auditory synaptopathy in model of Usher syndrome. <i>Journal of Clinical Investigation</i> , 2018, 128, 3382-3401.	8.2	97
7	Different Ca _v 1.3 Channel Isoforms Control Distinct Components of the Synaptic Vesicle Cycle in Auditory Inner Hair Cells. <i>Journal of Neuroscience</i> , 2017, 37, 2960-2975.	3.6	34
8	Local gene therapy durably restores vestibular function in a mouse model of Usher syndrome type 1G. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2017, 114, 9695-9700.	7.1	101
9	Otoferlin acts as a Ca ²⁺ sensor for vesicle fusion and vesicle pool replenishment at auditory hair cell ribbon synapses. <i>ELife</i> , 2017, 6, .	6.0	108
10	Hypervulnerability to Sound Exposure through Impaired Adaptive Proliferation of Peroxisomes. <i>Cell</i> , 2015, 163, 894-906.	28.9	158
11	Exocytotic Machineries of Vestibular Type I and Cochlear Ribbon Synapses Display Similar Intrinsic Otoferlin-Dependent Ca ²⁺ Sensitivity But a Different Coupling to Ca ²⁺ Channels. <i>Journal of Neuroscience</i> , 2014, 34, 10853-10869.	3.6	50
12	The Auditory Hair Cell Ribbon Synapse: From Assembly to Function. <i>Annual Review of Neuroscience</i> , 2012, 35, 509-528.	10.7	158
13	Control of Exocytosis by Synaptotagmins and Otoferlin in Auditory Hair Cells. <i>Journal of Neuroscience</i> , 2010, 30, 13281-13290.	3.6	106
14	Otoferlin Is Critical for a Highly Sensitive and Linear Calcium-Dependent Exocytosis at Vestibular Hair Cell Ribbon Synapses. <i>Journal of Neuroscience</i> , 2009, 29, 10474-10487.	3.6	113
15	Myosin VI is required for the proper maturation and function of inner hair cell ribbon synapses. <i>Human Molecular Genetics</i> , 2009, 18, 4615-4628.	2.9	81
16	Calcium- and Otoferlin-Dependent Exocytosis by Immature Outer Hair Cells. <i>Journal of Neuroscience</i> , 2008, 28, 1798-1803.	3.6	80
17	Chapter 8 Mouse Models for Human Hereditary Deafness. <i>Current Topics in Developmental Biology</i> , 2008, 84, 385-429.	2.2	68
18	â±V spectrin bridges the plasma membrane and cortical lattice in the lateral wall of the auditory outer hair cells. <i>Journal of Cell Science</i> , 2008, 121, 3347-3356.	2.0	62

#	ARTICLE	IF	CITATIONS
19	Otoferlin, Defective in a Human Deafness Form, Is Essential for Exocytosis at the Auditory Ribbon Synapse. <i>Cell</i> , 2006, 127, 277-289.	28.9	554
20	Ocsyn, a Novel Syntaxin-Interacting Protein Enriched in the Subapical Region of Inner Hair Cells. <i>Molecular and Cellular Neurosciences</i> , 2002, 20, 343-353.	2.2	15
21	Vesicle Targeting in Hair Cells. <i>Audiology and Neuro-Otology</i> , 2002, 7, 45-48.	1.3	7
22	Identification of a novel SNAP25 interacting protein (SIP30). <i>Journal of Neurochemistry</i> , 2002, 81, 1338-1347.	3.9	20
23	Cysteine-string protein in inner hair cells of the organ of Corti: synaptic expression and upregulation at the onset of hearing. <i>European Journal of Neuroscience</i> , 2002, 15, 1409-1420.	2.6	30
24	Vezeatin, a novel transmembrane protein, bridges myosin VIIA to the cadherin-catenins complex. <i>EMBO Journal</i> , 2000, 19, 6020-6029.	7.8	205
25	KCNQ4, a K ⁺ channel mutated in a form of dominant deafness, is expressed in the inner ear and the central auditory pathway. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2000, 97, 4333-4338.	7.1	382
26	Unconventional Myosin VIIA Is a Novel A-kinase-anchoring Protein. <i>Journal of Biological Chemistry</i> , 2000, 275, 29654-29659.	3.4	55
27	SNARE complex at the ribbon synapses of cochlear hair cells: analysis of synaptic vesicle- and synaptic membrane-associated proteins. <i>European Journal of Neuroscience</i> , 1999, 11, 803-812.	2.6	147
28	The Glutamate Receptor Subunit $\gamma 1$ Is Highly Expressed in Hair Cells of the Auditory and Vestibular Systems. <i>Journal of Neuroscience</i> , 1997, 17, 7523-7531.	3.6	59
29	Molecular Analysis of Excitatory Amino Acid Receptor Expression in the Cochlea. <i>Audiology and Neuro-Otology</i> , 1997, 2, 79-91.	1.3	36
30	Choline Acetyltransferase, Glutamate Decarboxylase, Tyrosine Hydroxylase, Calcitonin Gene-related Peptide and Opioid Peptides Coexist in Lateral Efferent Neurons of Rat and Guinea-pig. <i>European Journal of Neuroscience</i> , 1997, 9, 356-367.	2.6	80
31	Pre- and postsynaptic M3 muscarinic receptor mRNAs in the rodent peripheral auditory system. <i>Molecular Brain Research</i> , 1996, 40, 127-135.	2.3	24
32	Pre- and postsynaptic M3 muscarinic receptor mRNAs in the rodent peripheral auditory system. <i>Molecular Brain Research</i> , 1996, 40, 127-135.	2.3	3
33	Triple Immunofluorescence Evidence for the Coexistence of Acetylcholine, Enkephalins and Calcitonin Gene-related Peptide Within Efferent (Olivocochlear) Neurons of Rats and Guinea-pigs. <i>European Journal of Neuroscience</i> , 1992, 4, 981-992.	2.6	72