

Emanuela D'angelo

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/5350187/publications.pdf>

Version: 2024-02-01

29
papers

2,247
citations

361413

20
h-index

477307

29
g-index

31
all docs

31
docs citations

31
times ranked

2939
citing authors

#	ARTICLE	IF	CITATIONS
1	Uterine sarcomas: A review. <i>Gynecologic Oncology</i> , 2010, 116, 131-139.	1.4	659
2	Small cell carcinoma of the ovary, hypercalcemic type, displays frequent inactivating germline and somatic mutations in SMARCA4. <i>Nature Genetics</i> , 2014, 46, 427-429.	21.4	298
3	Ovarian carcinomas: at least five different diseases with distinct histological features and molecular genetics. <i>Human Pathology</i> , 2018, 80, 11-27.	2.0	170
4	Dual loss of the <sc>SWI</sc>/<sc>SNF</sc> complex <sc>ATPases SMARCA4</sc>/<sc>BRG1</sc> and <sc>SMARCA2</sc>/<sc>BRM</sc> is highly sensitive and specific for small cell carcinoma of the ovary, hypercalcaemic type. <i>Journal of Pathology</i> , 2016, 238, 389-400.	4.5	169
5	Comparative clinicopathologic and immunohistochemical analysis of uterine sarcomas diagnosed using the World Health Organization classification system. <i>Human Pathology</i> , 2009, 40, 1571-1585.	2.0	95
6	Prognostic significance of FOXL2 mutation and mRNA expression in adult and juvenile granulosa cell tumors of the ovary. <i>Modern Pathology</i> , 2011, 24, 1360-1367.	5.5	89
7	Endometrial Stromal Sarcomas With Sex Cord Differentiation Are Associated With PHF1 Rearrangement. <i>American Journal of Surgical Pathology</i> , 2013, 37, 514-521.	3.7	79
8	Undifferentiated and Dedifferentiated Endometrial Carcinomas With POLE Exonuclease Domain Mutations Have a Favorable Prognosis. <i>American Journal of Surgical Pathology</i> , 2017, 41, 1121-1128.	3.7	73
9	Uterine leiomyosarcomas: Tumor size, mitotic index, and biomarkers Ki67, and Bcl-2 identify two groups with different prognosis. <i>Gynecologic Oncology</i> , 2011, 121, 328-333.	1.4	69
10	Myometrial Invasion and Lymph Node Metastasis in Endometrioid Carcinomas: Tumor-associated Macrophages, Microvessel Density, and HIF1A Have a Crucial Role. <i>American Journal of Surgical Pathology</i> , 2010, 34, 1708-1714.	3.7	67
11	Pathology of mixed MÃ¼llerian tumours. <i>Best Practice and Research in Clinical Obstetrics and Gynaecology</i> , 2011, 25, 705-718.	2.8	60
12	Molecular analyses of juvenile granulosa cell tumors bearing <i>AKT1</i> mutations provide insights into tumor biology and therapeutic leads. <i>Human Molecular Genetics</i> , 2015, 24, 6687-6698.	2.9	51
13	Germ cell tumour growth patterns originating from clear cell carcinomas of the ovary and endometrium: a comparative immunohistochemical study favouring their origin from somatic stem cells. <i>Histopathology</i> , 2018, 72, 634-647.	2.9	48
14	Expression profiling of 22 genes involved in the PI3Kâ€“AKT pathway identifies two subgroups of high-grade endometrial carcinomas with different molecular alterations. <i>Modern Pathology</i> , 2010, 23, 694-702.	5.5	47
15	Simultaneous Carcinomas of the Breast and Ovary. <i>International Journal of Gynecological Pathology</i> , 2015, 34, 257-265.	1.4	47
16	Mixed and Ambiguous Endometrial Carcinomas. <i>American Journal of Surgical Pathology</i> , 2016, 40, 972-981.	3.7	29
17	Dedifferentiated endometrial carcinomas with neuroendocrine features: a clinicopathologic, immunohistochemical, and molecular genetic study. <i>Human Pathology</i> , 2018, 72, 100-106.	2.0	28
18	Loss-of-function variants in myocardin cause congenital megabladder in humans and mice. <i>Journal of Clinical Investigation</i> , 2019, 129, 5374-5380.	8.2	27

#	ARTICLE	IF	CITATIONS
19	Gene expression analysis identifies two groups of ovarian high-grade serous carcinomas with different prognosis. <i>Modern Pathology</i> , 2011, 24, 846-854.	5.5	26
20	The FOXO1-miR27 tandem regulates myometrial invasion in endometrioid endometrial adenocarcinoma. <i>Human Pathology</i> , 2014, 45, 942-951.	2.0	21
21	CD8 down-regulation on cytotoxic T lymphocytes of patients with endometrioid endometrial carcinomas. <i>Human Pathology</i> , 2016, 56, 180-188.	2.0	20
22	Endometrioid endometrial carcinomas with microcystic, elongated, and fragmented (MELF) type of myoinvasion: role of immunohistochemistry in the detection of occult lymph node metastases and their clinical significance. <i>Human Pathology</i> , 2017, 70, 6-13.	2.0	18
23	Squamous Cell Carcinoma of the Ovary Arising From a Mucinous Cystic Tumor of Endocervical (Müllerian) Type. <i>International Journal of Gynecological Pathology</i> , 2010, 29, 529-532.	1.4	15
24	Diagnostic use of immunohistochemistry in uterine mesenchymal tumors. <i>Seminars in Diagnostic Pathology</i> , 2014, 31, 216-222.	1.5	14
25	Classification of ovarian carcinomas based on pathology and molecular genetics. <i>Clinical and Translational Oncology</i> , 2010, 12, 783-787.	2.4	9
26	Mixed endometrial carcinomas with a "low-grade serous"-like component: a clinicopathologic, immunohistochemical, and molecular genetic study. <i>Human Pathology</i> , 2018, 71, 65-73.	2.0	9
27	Central pathology review of early-stage ovarian carcinoma: Description and correlation with follow-up—A study by the Spanish Group for Ovarian Cancer Research (GEICO).. <i>Journal of Clinical Oncology</i> , 2014, 32, 5583-5583.	1.6	2
28	A surprising evolution from oral human papillomavirus 16 infection to lymph node metastasis of tonsillar squamous cell carcinoma in an HIV-infected patient. <i>Aids</i> , 2012, 26, 1044-1045.	2.2	0
29	Clinicopathological Significance of Taz and VGLL1 Expression in Early Stage Ovarian Cancer. A study by Spanish Group for Ovarian Cancer Research (GEICO).. <i>Journal of Clinical Oncology</i> , 2015, 33, e16583-e16583.	1.6	0