## Hiroshi Yatsuya

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/5373054/publications.pdf

Version: 2024-02-01

238 papers 22,625 citations

47 h-index

47006

145 g-index

240 all docs

240 docs citations

240 times ranked 41733 citing authors

#	Article	IF	CITATIONS
1	Global, regional, and national prevalence of overweight and obesity in children and adults during 1980–2013: a systematic analysis for the Global Burden of Disease Study 2013. Lancet, The, 2014, 384, 766-781.	13.7	9,122
2	Global, regional, and national age–sex specific all-cause and cause-specific mortality for 240 causes of death, 1990–2013: a systematic analysis for the Global Burden of Disease Study 2013. Lancet, The, 2015, 385, 117-171.	13.7	5,847
3	Community Prevalence of Ideal Cardiovascular Health, by the American Heart Association Definition, and Relationship With Cardiovascular Disease Incidence. Journal of the American College of Cardiology, 2011, 57, 1690-1696.	2.8	614
4	Eating Fast Leads to Obesity: Findings Based on Self-administered Questionnaires among Middle-aged Japanese Men and Women. Journal of Epidemiology, 2006, 16, 117-124.	2.4	221
5	Global Trend in Overweight and Obesity and Its Association With Cardiovascular Disease Incidence. Circulation Journal, 2014, 78, 2807-2818.	1.6	177
6	Retinal Microvascular Abnormalities and Risk of Lacunar Stroke. Stroke, 2010, 41, 1349-1355.	2.0	172
7	Comparison of Circulating Adiponectin and Proinflammatory Markers Regarding Their Association With Metabolic Syndrome in Japanese Men. Arteriosclerosis, Thrombosis, and Vascular Biology, 2006, 26, 871-876.	2.4	160
8	The Impact of Green Tea and Coffee Consumption on the Reduced Risk of Stroke Incidence in Japanese Population. Stroke, 2013, 44, 1369-1374.	2.0	123
9	Neoadjuvant Oxaliplatin and Capecitabine and Bevacizumab without Radiotherapy for Poor-risk Rectal Cancer: N-SOG 03 Phase II Trial. Japanese Journal of Clinical Oncology, 2013, 43, 964-971.	1.3	119
10	Eating fast leads to insulin resistance: Findings in middle-aged Japanese men and women. Preventive Medicine, 2008, 46, 154-159.	3.4	118
11	Prospective study of screening for stomach cancer in Japan. International Journal of Cancer, 2003, 106, 103-107.	5.1	113
12	Thermodynamic instability of siRNA duplex is a prerequisite for dependable prediction of siRNA activities. Nucleic Acids Research, 2007, 35, e123.	14.5	109
13	BMI and Allâ $\in$ cause Mortality Among Japanese Older Adults: Findings From the Japan Collaborative Cohort Study. Obesity, 2010, 18, 362-369.	3.0	106
14	Differences by sex in the prevalence of diabetes mellitus, impaired fasting glycaemia and impaired glucose tolerance in sub-Saharan Africa: a systematic review and meta-analysis. Bulletin of the World Health Organization, 2013, 91, 671-682D.	3.3	102
15	Association of extremely high levels of high-density lipoprotein cholesterol with cardiovascular mortality in a pooled analysis of 9 cohort studies including 43,407 individuals: The EPOCH–JAPAN study. Journal of Clinical Lipidology, 2018, 12, 674-684.e5.	1.5	101
16	Dietary intakes of fat and fatty acids and risk of breast cancer: A prospective study in Japan. Cancer Science, 2005, 96, 590-599.	3.9	97
17	Leptin Is Associated with an Increased Female Colorectal Cancer Risk: A Nested Case-Control Study in Japan. Oncology, 2005, 68, 454-461.	1.9	94
18	Rate of Decline of Forced Vital Capacity Predicts Future Arterial Hypertension. Hypertension, 2012, 59, 219-225.	2.7	91

#	Article	IF	CITATIONS
19	Postural Changes in Blood Pressure and Incidence of Ischemic Stroke Subtypes. Hypertension, 2011, 57, 167-173.	2.7	89
20	Healthy lifestyle behaviours and cardiovascular mortality among Japanese men and women: the Japan collaborative cohort study. European Heart Journal, 2012, 33, 467-477.	2.2	88
21	Dietary intake of saturated fatty acids and mortality from cardiovascular disease in Japanese: the Japan Collaborative Cohort Study for Evaluation of Cancer Risk (JACC) Study. American Journal of Clinical Nutrition, 2010, 92, 759-765.	4.7	87
22	Associations of Protein, Fat, and Carbohydrate Intakes With Insomnia Symptoms Among Middle-aged Japanese Workers. Journal of Epidemiology, 2013, 23, 132-138.	2.4	83
23	Bronchus sign on thin-section computed tomography is a powerful predictive factor for successful transbronchial biopsy using endobronchial ultrasound with a guide sheath for small peripheral lung lesions: a retrospective observational study. BMC Medical Imaging, 2015, 15, 21.	2.7	83
24	Race- and Sex-Specific Associations of Obesity Measures With Ischemic Stroke Incidence in the Atherosclerosis Risk in Communities (ARIC) Study. Stroke, 2010, 41, 417-425.	2.0	82
25	Daytime napping and mortality, with a special reference to cardiovascular disease: the JACC study. International Journal of Epidemiology, 2010, 39, 233-243.	1.9	80
26	Association of White Blood Cell Count and Clustered Components of Metabolic Syndrome in Japanese Men. Circulation Journal, 2004, 68, 892-897.	1.6	77
27	Risk of Cardiovascular Disease from Cumulative Cigarette Use and the Impact of Smoking Intensity. Epidemiology, 2016, 27, 395-404.	2.7	74
28	Determinants of self-rated health: Could health status explain the association between self-rated health and mortality?. Archives of Gerontology and Geriatrics, 2006, 43, 369-380.	3.0	69
29	Long-Term Body Weight Fluctuation is Associated With Metabolic Syndrome Independent of Current Body Mass Index Among Japanese Men. Circulation Journal, 2005, 69, 13-18.	1.6	66
30	Dietary intake of saturated fatty acids and incident stroke and coronary heart disease in Japanese communities: the JPHC Study. European Heart Journal, 2013, 34, 1225-1232.	2.2	66
31	Association of Kidney Disease Measures With Ischemic Versus Hemorrhagic Strokes. Stroke, 2014, 45, 1925-1931.	2.0	66
32	Risk of Incident Cardiovascular Disease Among Users of Smokeless Tobacco in the Atherosclerosis Risk in Communities (ARIC) Study. American Journal of Epidemiology, 2010, 172, 600-605.	3.4	64
33	Body Mass Index and Risk of Stroke and Myocardial Infarction in a Relatively Lean Population. Circulation: Cardiovascular Quality and Outcomes, 2010, 3, 498-505.	2.2	62
34	Consumption of soy foods and the risk of breast cancer: findings from the Japan Collaborative Cohort (JACC) Study. Cancer Causes and Control, 2007, 18, 801-808.	1.8	59
35	Reduction of Inner Retinal Thickness in Patients with Autosomal Dominant Optic Atrophy Associated with OPA1 Mutations., 2007, 48, 4079.		57
36	Inverse association between adiponectin and C-reactive protein in substantially healthy Japanese men. Atherosclerosis, 2006, 188, 184-189.	0.8	56

#	Article	IF	Citations
37	Adiponectin Level and Left Ventricular Hypertrophy in Japanese Men. Hypertension, 2007, 49, 1448-1454.	2.7	55
38	White Blood Cell Count and Risk of All-Cause and Cardiovascular Mortality in Nationwide Sample of Japanese. Circulation Journal, 2007, 71, 479-485.	1.6	55
39	Factors Associated With Life Space Among Community-Living Rural Elders in Japan. Public Health Nursing, 2006, 23, 324-331.	1.5	54
40	Role of the Anterolateral Thigh Flap in Head and Neck Reconstruction: Advantages of Moderate Skin and Subcutaneous Thickness. Journal of Reconstructive Microsurgery, 2002, 18, 141-146.	1.8	53
41	Breakfast Skipping is Positively Associated With Incidence of Type 2 Diabetes Mellitus: Evidence From the Aichi Workers' Cohort Study. Journal of Epidemiology, 2015, 25, 351-358.	2.4	53
42	Self-reported medical history was generally accurate among Japanese workplace population. Journal of Clinical Epidemiology, 2009, 62, 306-313.	5.0	52
43	Impact of Age at Smoking Initiation, Dosage, and Time Since Quitting on Cardiovascular Disease in African Americans and Whites: The Atherosclerosis Risk in Communities Study. American Journal of Epidemiology, 2012, 175, 816-826.	3.4	52
44	Differences in Urinary Arsenic Metabolites between Diabetic and Non-Diabetic Subjects in Bangladesh. International Journal of Environmental Research and Public Health, 2013, 10, 1006-1019.	2.6	52
45	Dietary Habits and Stomach Cancer Risk in the JACC Study. Journal of Epidemiology, 2005, 15, S98-S108.	2.4	51
46	Group Education with Personal Rehabilitation for Idiopathic Parkinson's Disease. Canadian Journal of Neurological Sciences, 2009, 36, 51-59.	0.5	50
47	Association between weight fluctuation and fasting insulin concentration in Japanese men. International Journal of Obesity, 2003, 27, 478-483.	3.4	49
48	Active Smoking, Passive Smoking, and Breast Cancer Risk: Findings from the Japan Collaborative Cohort Study for Evaluation of Cancer Risk. Journal of Epidemiology, 2008, 18, 77-83.	2.4	49
49	Early age at menarche associated with increased all-cause mortality. European Journal of Epidemiology, 2011, 26, 771-778.	5.7	49
50	Population profile and residential environment of an urban poor community in Dhaka, Bangladesh. Environmental Health and Preventive Medicine, 2017, 22, 1.	3.4	49
51	Relationship of Estimated GFR and Albuminuria to Concurrent Laboratory Abnormalities: An Individual Participant Data Meta-analysis in a Global Consortium. American Journal of Kidney Diseases, 2019, 73, 206-217.	1.9	49
52	Family history and the risk of stomach cancer death in Japan: Differences by age and gender. International Journal of Cancer, 2002, 97, 688-694.	5.1	48
53	Uric Acid and Left Ventricular Hypertrophy in Japanese Men. Circulation Journal, 2009, 73, 667-672.	1.6	48
54	Association between mortality and incidence rates of coronary heart disease and stroke: The Japan Public Health Center-based prospective (JPHC) study. International Journal of Cardiology, 2016, 222, 281-286.	1.7	47

#	Article	IF	CITATIONS
55	Perceived Psychological Stress and Serum Leptin Concentrations in Japanese Men. Obesity, 2006, 14, 1832-1838.	3.0	46
56	Associations of Obesity Measures with Subtypes of Ischemic Stroke in the ARIC Study. Journal of Epidemiology, 2010, 20, 347-354.	2.4	46
57	Risk of intraparenchymal hemorrhage with magnetic resonance imagingâ€defined leukoaraiosis and brain infarcts. Annals of Neurology, 2012, 71, 552-559.	5.3	46
58	Prevalence of non-communicable disease risk factors among poor shantytown residents in Dhaka, Bangladesh: a community-based cross-sectional survey. BMJ Open, 2017, 7, e014710.	1.9	46
59	Impact of menstrual and reproductive factors on breast cancer risk in Japan: Results of the JACC study. Cancer Science, 2005, 96, 57-62.	3.9	45
60	Smoking status and adiponectin in healthy Japanese men and women. Preventive Medicine, 2007, 45, 471-475.	3.4	45
61	Development of a Point-based Prediction Model for the Incidence of Total Stroke. Stroke, 2013, 44, 1295-1302.	2.0	44
62	Serum Phospholipid Transfer Protein Mass as a Possible Protective Factor for Coronary Heart Diseases. Circulation Journal, 2004, 68, 11-16.	1.6	43
63	Association of high-density lipoprotein cholesterol concentration with different types of stroke and coronary heart disease: The Japan Public Health Center-based prospective (JPHC) study. Atherosclerosis, 2017, 265, 147-154.	0.8	43
64	Higher dietary intake of alpha-linolenic acid is associated with lower insulin resistance in middle-aged Japanese. Preventive Medicine, 2010, 50, 272-276.	3.4	42
65	Comparing different definitions of prediabetes with subsequent risk of diabetes: an individual participant data meta-analysis involving 76 513 individuals and 8208 cases of incident diabetes. BMJ Open Diabetes Research and Care, 2019, 7, e000794.	2.8	42
66	Rice consumption is not associated with risk of cardiovascular disease morbidity or mortality in Japanese men and women: a large population-based, prospective cohort study. American Journal of Clinical Nutrition, 2014, 100, 199-207.	4.7	40
67	Past Decline Versus Current eGFR and Subsequent Mortality Risk. Journal of the American Society of Nephrology: JASN, 2016, 27, 2456-2466.	6.1	40
68	Effect of Physical Activity on Breast Cancer Risk: Findings of the Japan Collaborative Cohort Study. Cancer Epidemiology Biomarkers and Prevention, 2008, 17, 3396-3401.	2.5	38
69	Milk Drinking and Mortality: Findings From the Japan Collaborative Cohort Study. Journal of Epidemiology, 2015, 25, 66-73.	2.4	38
70	Development of a Risk Equation for the Incidence of Coronary Artery Disease and Ischemic Stroke for Middle-Aged Japanese – Japan Public Health Center-Based Prospective Study –. Circulation Journal, 2016, 80, 1386-1395.	1.6	37
71	Non-communicable disease risk factor profile among public employees in a regional city in northern Ethiopia. Scientific Reports, 2018, 8, 9298.	3.3	37
72	Effects of Social Relationships on Mortality among the Elderly in a Japanese Rural Area: An 88-month Follow-up Study. Journal of Epidemiology, 2005, 15, 78-84.	2.4	36

#	Article	IF	CITATIONS
73	Association between Physical Activity and Risk of Stroke Subtypes: The Atherosclerosis Risk in Communities Study. Neuroepidemiology, 2013, 40, 109-116.	2.3	36
74	Individual and joint impact of family history and Helicobacter pylori infection on the risk of stomach cancer: a nested case–control study. British Journal of Cancer, 2004, 91, 929-934.	6.4	35
75	Incidence of metabolic syndrome according to combinations of lifestyle factors among middle-aged Japanese male workers. Preventive Medicine, 2010, 51, 118-122.	3.4	35
76	Dietary magnesium intake and risk of incident coronary heart disease in men: A prospective cohort study. Clinical Nutrition, 2018, 37, 1602-1608.	5.0	35
77	Interleukin-8 gene polymorphism associated with susceptibility to non-cardia gastric carcinoma with microsatellite instability. Journal of Gastroenterology and Hepatology (Australia), 2006, 21, 1129-1135.	2.8	32
78	High-Sensitivity C-Reactive Protein is Quite Low in Japanese Men at High Coronary Risk. Circulation Journal, 2007, 71, 820-825.	1.6	32
79	Watching Television and Risk of Mortality From Pulmonary Embolism Among Japanese Men and Women. Circulation, 2016, 134, 355-357.	1.6	32
80	Association between parental histories of hypertension, diabetes and dyslipidemia and the clustering of these disorders in offspring. Preventive Medicine, 2006, 42, 358-363.	3.4	31
81	Psychological factors and insomnia among male civil servants in Japan. Sleep Medicine, 2007, 8, 209-214.	1.6	31
82	Positive association between highâ€sensitivity Câ€reactive protein and incidence of type 2 diabetes mellitus in Japanese workers: 6â€year followâ€up. Diabetes/Metabolism Research and Reviews, 2013, 29, 398-405.	4.0	30
83	Lifetime Risk of Stroke and Coronary Heart Disease Deaths According to Blood Pressure Level. Hypertension, 2019, 73, 52-59.	2.7	30
84	Non-High-Density Lipoprotein Cholesterol and Risk of Stroke Subtypes and Coronary Heart Disease: The Japan Public Health Center-Based Prospective (JPHC) Study. Journal of Atherosclerosis and Thrombosis, 2020, 27, 363-374.	2.0	30
85	Smoking and Diabetes: Is the Association Mediated by Adiponectin, Leptin, or C-reactive Protein?. Journal of Epidemiology, 2015, 25, 99-109.	2.4	29
86	Validation of the Japanese Version of the Yale Food Addiction Scale 2.0 (J-YFAS 2.0). Nutrients, 2019, 11, 687.	4.1	29
87	Alcohol consumption and risk of stroke and coronary heart disease among Japanese women: The Japan Public Health Center-based prospective study. Preventive Medicine, 2013, 57, 505-510.	3.4	28
88	Working Hours and Risk of Acute Myocardial Infarction and Stroke Among Middle-Aged Japanese Men ― The Japan Public Health Center-Based Prospective Study Cohort II ―. Circulation Journal, 2019, 83, 1072-1079.	1.6	28
89	Th1/Th2 Immune Response in Lung Fibroblasts in Interstitial Lung Disease. Archives of Medical Research, 2008, 39, 503-510.	3.3	27
90	Vaginal Douching in Cambodian Women: Its Prevalence and Association With Vaginal Candidiasis. Journal of Epidemiology, 2010, 20, 70-76.	2.4	27

#	Article	IF	Citations
91	Hemoglobin A1c Levels and the Risk of Cardiovascular Disease in People Without Known Diabetes. Medicine (United States), 2015, 94, e785.	1.0	27
92	Green Tea and Stomach Cancer – A Short Review of Prospective Studies. Journal of Epidemiology, 2005, 15, S109-S112.	2.4	26
93	Birth Weight and Adult Hypertension Cross-Sectional Study in a Japanese Workplace Population. Circulation Journal, 2006, 70, 262-267.	1.6	26
94	Upper airway morphology in patients with obstructive sleep apnea syndrome: Effects of lateral positioning. Auris Nasus Larynx, 2009, 36, 305-309.	1.2	26
95	Association between serum leptin concentration and white blood cell count in middle-aged Japanese men and women. Diabetes/Metabolism Research and Reviews, 2005, 21, 441-447.	4.0	25
96	Psychological attitudes and risk of breast cancer in Japan: a prospective study. Cancer Causes and Control, 2007, 18, 259-267.	1.8	25
97	Multicenter feasibility study of bowel preparation with castor oil for colon capsule endoscopy.  Digestive Endoscopy, 2019, 31, 164-172.	2.3	25
98	Smoking Behavior and Lung Cancer in a Biracial Cohort. American Journal of Preventive Medicine, 2014, 46, 624-632.	3.0	24
99	Effect of laughter yoga on salivary cortisol and dehydroepiandrosterone among healthy university students: A randomized controlled trial. Complementary Therapies in Clinical Practice, 2018, 32, 6-11.	1.7	24
100	Cigarette Smoking and Mortality due to Stomach Cancer: Findings from the JACC Study Journal of Epidemiology, 2005, 15, S113-S119.	2.4	23
101	The transition to menopause reinforces adiponectin production and its contribution to improvement of insulin-resistant state. Clinical Endocrinology, 2006, 66, 061109020454003-???.	2.4	23
102	HIV prevalence and factors associated with HIV infection among male injection drug users under 30: a cross-sectional study in Long An, Vietnam. BMC Public Health, 2006, 6, 248.	2.9	23
103	Body Mass Index and Risks of Incident Ischemic Stroke Subtypes: The Japan Public Health Center-Based Prospective (JPHC) Study. Journal of Epidemiology, 2019, 29, 325-333.	2.4	23
104	Trends in the Mortality (1950–1997) and Incidence (1975–1993) of Malignant Ovarian Neoplasm among Japanese Women: Analyses by Age, Time, and Birth Cohort. Gynecologic Oncology, 2001, 83, 64-71.	1.4	22
105	Relationship between body mass index and the risk of ovarian cancer in the Japanese population: Findings from the Japanese Collaborate Cohort (JACC) study. Journal of Obstetrics and Gynaecology Research, 2005, 31, 452-458.	1.3	22
106	Prospective study of alcohol consumption and breast cancer risk in Japanese women. International Journal of Cancer, 2005, 116, 779-783.	5.1	22
107	A nested case-control study of stomach cancer and serum insulin-like growth factor (IGF)-1, IGF-2 and IGF-binding protein (IGFBP)-3. European Journal of Cancer, 2007, 43, 1611-1616.	2.8	22
108	Effect of the Interaction between Mental Stress and Eating Pattern on Body Mass Index Gain in Healthy Japanese Male Workers. Journal of Epidemiology, 2009, 19, 88-93.	2.4	22

#	Article	IF	Citations
109	Association of gammaâ€glutamyl transferase and alanine aminotransferase with typeÂ2 diabetes mellitus incidence in middleâ€aged Japanese men: 12â€year follow up. Journal of Diabetes Investigation, 2019, 10, 837-845.	2.4	22
110	Reproducibility and Validity of a Simple Checklist-type Questionnaire for Food Intake and Dietary Behavior. Journal of Epidemiology, 2003, 13, 235-245.	2.4	21
111	Serum Pepsinogen Values and Helicobacter pylori Status among Control Subjects of a Nested Case-Control Study in the JACC study. Journal of Epidemiology, 2005, 15, S126-S133.	2.4	21
112	The Effort-reward Imbalance work-stress model and daytime salivary cortisol and dehydroepiandrosterone (DHEA) among Japanese women. Scientific Reports, 2014, 4, 6402.	3.3	21
113	Similarities and differences between coronary heart disease and stroke in the associations with cardiovascular risk factors: The Japan Collaborative Cohort Study. Atherosclerosis, 2017, 261, 124-130.	0.8	21
114	Changes in C-reactive protein during weight loss and the association with changes in anthropometric variables in men and women: LIFE Study. International Journal of Obesity, 2011, 35, 684-691.	3.4	20
115	Prospective study of seaweed consumption and thyroid cancer incidence in women. European Journal of Cancer Prevention, 2016, 25, 239-245.	1.3	20
116	Aggregation of stomach cancer history in parents and offspring in comparison with other sites. International Journal of Epidemiology, 2003, 32, 579-583.	1.9	19
117	Postoperative DAVâ€IFNâ€Î² therapy does not improve survival rates of stage II and stage III melanoma patients significantly. Journal of the European Academy of Dermatology and Venereology, 2013, 27, 1514-1520.	2.4	19
118	Descriptive epidemiological study of food intake among Japanese adults: analyses by age, time and birth cohort model. BMC Public Health, 2014, 14, 328.	2.9	19
119	Association between parental history of diabetes and the incidence of type 2 diabetes mellitus differs according to the sex of the parent and offspring's body weight: A finding from a Japanese worksite-based cohort study. Preventive Medicine, 2015, 81, 49-53.	3.4	19
120	A prospective study on the possible association between having children and colon cancer risk: Findings from the JACC Study. Cancer Science, 2004, 95, 243-247.	3.9	18
121	A prospective study of reproductive and menstrual factors and colon cancer risk in Japanese women: Findings from the JACC study. Cancer Science, 2004, 95, 602-607.	3.9	18
122	Association Between Average Daily Television Viewing Time and Chronic Obstructive Pulmonary Disease-Related Mortality: Findings From the Japan Collaborative Cohort Study. Journal of Epidemiology, 2015, 25, 431-436.	2.4	18
123	Long-term weight-change slope, weight fluctuation and risk of type 2 diabetes mellitus in middle-aged Japanese men and women: findings of Aichi Workers' Cohort Study. Nutrition and Diabetes, 2017, 7, e252-e252.	3.2	18
124	Habitual tub bathing and risks of incident coronary heart disease and stroke. Heart, 2020, 106, 732-737.	2.9	18
125	Prioritization and sequential exclusion of articles in systematic reviews. Campbell Systematic Reviews, 2022, $18$ , .	3.0	18
126	Modification of the Excess Risk of Coronary Heart Disease Due to Smoking by Seafood/Fish Intake. American Journal of Epidemiology, 2014, 179, 1173-1181.	3.4	17

#	Article	IF	Citations
127	MRI and FDG-PET for Assessment of Response to Neoadjuvant Chemotherapy in Locally Advanced Rectal Cancer. Annals of Surgical Oncology, 2014, 21, 1801-1808.	1.5	17
128	Passive smoking and chronic obstructive pulmonary disease mortality: findings from the Japan collaborative cohort study. International Journal of Public Health, 2017, 62, 489-494.	2.3	17
129	Smoking cessation and COPD mortality among Japanese men and women: The JACC study. Preventive Medicine, 2012, 55, 639-643.	3.4	16
130	Low Birth Weight Is Associated With Reduced Adiponectin Concentration in Adult. Annals of Epidemiology, 2006, 16, 669-674.	1.9	15
131	Factors predicting growth of vestibular schwannoma in neurofibromatosis type 2. Neurosurgical Review, 2009, 32, 425-433.	2.4	15
132	Strengthening community participation at health centers in rural Cambodia: role of local non-governmental organizations (NGOs). Critical Public Health, 2010, 20, 97-115.	2.4	15
133	Maintenance-tailored therapy vs. standard behavior therapy for 30-month maintenance of weight loss. Preventive Medicine, 2010, 51, 457-459.	3.4	15
134	Adiposity and risk of cardiovascular diseases in Japan: secular trend, individual level associations and causal pathway $\hat{a} \in \mathbb{C}^m$ implications for the prevention of cardiovascular diseases in societies with rapid economic development. EPMA Journal, 2011, 2, 65-73.	6.1	15
135	No modifying effect of education level on the association between lifestyle behaviors and cardiovascular mortality: the Japan Collaborative Cohort Study. Scientific Reports, 2017, 7, 39820.	3.3	15
136	Risk Factors of Infectious Complications After Endobronchial Ultrasound-Guided Transbronchial Biopsy. Chest, 2020, 158, 797-807.	0.8	15
137	Further inflammatory information on metabolic syndrome by adiponectin evaluation. International Journal of Cardiology, 2008, 124, 339-344.	1.7	14
138	Contribution of adipocytokines to low-grade inflammatory state as expressed by circulating C-reactive protein in Japanese men: Comparison of leptin and adiponectin. International Journal of Cardiology, 2008, 130, 159-164.	1.7	14
139	Height and Risk of Incident Intraparenchymal Hemorrhage: Atherosclerosis Risk in Communities and Cardiovascular Health Study Cohorts. Journal of Stroke and Cerebrovascular Diseases, 2013, 22, 323-328.	1.6	14
140	Independent association of liver fat accumulation with insulin resistance. Obesity Research and Clinical Practice, 2014, 8, e350-e355.	1.8	14
141	Weight change during middle age and risk of stroke and coronary heart disease: The Japan Public Health Center–based Prospective Study. Atherosclerosis, 2021, 322, 67-73.	0.8	14
142	Gender and Age Differences in Lifestyle Factors Related to Hypertension in Middle-Aged Civil Service Employees Journal of Epidemiology, 2003, 13, 38-47.	2.4	13
143	Lung cancer mortality and body mass index in a Japanese cohort: findings from the Japan Collaborative Cohort Study (JACC Study). Cancer Causes and Control, 2007, 18, 229-234.	1.8	13
144	Sexâ€Specific HDL Cholesterol Changes With Weight Loss and Their Association With Anthropometric Variables: The LIFE Study. Obesity, 2011, 19, 429-435.	3.0	13

#	Article	IF	Citations
145	Body Mass Index and Weight Change During Adulthood Are Associated With Increased Mortality From Liver Cancer: The JACC Study. Journal of Epidemiology, 2013, 23, 219-226.	2.4	13
146	Serum $\hat{I}^3$ -glutamyltransferase and Mortality due to Cardiovascular Disease in Japanese Men and Women. Journal of Atherosclerosis and Thrombosis, 2016, 23, 792-799.	2.0	13
147	Synergistic and Non-synergistic Associations for Cigarette Smoking and Non-tobacco Risk Factors for Cardiovascular Disease Incidence in the Atherosclerosis Risk In Communities (ARIC) Study. Nicotine and Tobacco Research, 2016, 19, ntw235.	2.6	13
148	Recent Status and Methodological Quality of Return-to-Work Rates of Cancer Patients Reported in Japan: A Systematic Review. International Journal of Environmental Research and Public Health, 2019, 16, 1461.	2.6	13
149	Associations of Daily Walking Time With Pneumonia Mortality Among Elderly Individuals With or Without a Medical History of Myocardial Infarction or Stroke: Findings From the Japan Collaborative Cohort Study. Journal of Epidemiology, 2019, 29, 233-237.	2.4	13
150	Perceptions and behavior related to noncommunicable diseases among slum dwellers in a rapidly urbanizing city, Dhaka, Bangladesh: a qualitative study. Nagoya Journal of Medical Science, 2018, 80, 559-569.	0.3	13
151	Serum Levels of Insulin-like Growth Factor I, II, and Binding Protein 3, Transforming Growth Factor $\hat{l}^2$ -1, Soluble Fas Ligand and Superoxide Dismutase Activity in Stomach Cancer Cases and Their Controls in the JACC Study. Journal of Epidemiology, 2005, 15, S120-S125.	2.4	12
152	Cigarette smoking and the risk of ovarian cancer in the Japanese population: Findings from the Japanese Collaborate Cohort study. Journal of Obstetrics and Gynaecology Research, 2005, 31, 144-151.	1.3	12
153	A varicella outbreak in B-cell lymphoma patients receiving rituximab-containing chemotherapy. Journal of Infection and Chemotherapy, 2014, 20, 774-777.	1.7	12
154	Profile of Non-communicable Disease Risk Factors Among Young People in Palau. Journal of Epidemiology, 2015, 25, 392-397.	2.4	12
155	Relationships among Socioeconomic Factors and Self-rated Health in Japanese Adults: NIPPON DATA2010. Journal of Epidemiology, 2018, 28, S66-S72.	2.4	12
156	A Point System for Predicting 10-Year Risk of Developing Type 2 Diabetes Mellitus in Japanese Men: Aichi Workers' Cohort Study. Journal of Epidemiology, 2018, 28, 347-352.	2.4	12
157	Low leptin but high insulin resistance of smokers in Japanese men. Diabetes Research and Clinical Practice, 2008, 81, 358-364.	2.8	11
158	Inverse relationship of serum adiponectin concentration with type 2 diabetes mellitus incidence in middleâ€aged Japanese workers: sixâ€year followâ€up. Diabetes/Metabolism Research and Reviews, 2012, 28, 349-356.	4.0	11
159	Alcohol consumption and mortality from aortic disease among Japanese men: The Japan Collaborative Cohort study. Atherosclerosis, 2017, 266, 64-68.	0.8	11
160	The association between objective measures of residence and worksite neighborhood environment, and self-reported leisure-time physical activities: The Aichi Workers' Cohort Study. Preventive Medicine Reports, 2018, 11, 282-289.	1.8	11
161	Association of Vegetable, Fruit, and Okinawan Vegetable Consumption With Incident Stroke and Coronary Heart Disease. Journal of Epidemiology, 2020, 30, 37-45.	2.4	11
162	Profile of non-communicable disease risk factors among adults in the Republic of Palau: findings of a national STEPS survey. Nagoya Journal of Medical Science, 2015, 77, 609-19.	0.3	11

#	Article	IF	CITATIONS
163	Impact of Body Mass Index on Obesity-Related Cancer and Cardiovascular Disease Mortality; The Japan Collaborative Cohort Study. Journal of Atherosclerosis and Thrombosis, 2022, 29, 1547-1562.	2.0	11
164	Short- and Long-term Reliability of Information on Previous Illness and Family History as Compared with that on Smoking and Drinking Habits in Questionnaire Surveys Journal of Epidemiology, 2002, 12, 120-125.	2.4	10
165	An influence of Interferon-1 <sup>3</sup> gene polymorphisms on treatment response to tuberculosis in Japanese population. Journal of Infection, 2009, 58, 467-469.	3.3	10
166	Carotid Intima-Media Thickness, Electrocardiographic Left Ventricular Hypertrophy, and Incidence of Intracerebral Hemorrhage. Stroke, 2011, 42, 3075-3079.	2.0	10
167	Risk Factor of Cardiovascular Disease Among Older Individuals. Journal of Atherosclerosis and Thrombosis, 2017, 24, 258-261.	2.0	10
168	Association between average daily television viewing time and the incidence of ovarian cancer: findings from the Japan Collaborative Cohort Study. Cancer Causes and Control, 2018, 29, 213-219.	1.8	10
169	Cardiovascular disease mortality in relation to physical activity during adolescence and adulthood in Japan: Does school-based sport club participation matter?. Preventive Medicine, 2018, 113, 102-108.	3.4	10
170	Estimation of 10-Year Risk of Death from Coronary Heart Disease, Stroke, and Cardiovascular Disease in a Pooled Analysis of Japanese Cohorts: EPOCH-JAPAN. Journal of Atherosclerosis and Thrombosis, 2021, 28, 816-825.	2.0	10
171	Prevalence and predictors of prediabetes and diabetes among adults in Palau: population-based national STEPS survey. Nagoya Journal of Medical Science, 2016, 78, 475-483.	0.3	10
172	Plasma Fibrinogen Levels and Cardiovascular Risk Factors in Japanese Schoolchildren. Journal of Epidemiology, 2006, 16, 64-70.	2.4	9
173	A Positive Association between Leptin and Blood Pressure of Normal Range in Japanese Men. Hypertension Research, 2006, 29, 485-492.	2.7	9
174	Demographic and Cardiovascular Risk Factors Modify Association of Fasting Insulin With Incident Coronary Heart Disease and Ischemic Stroke (from the Atherosclerosis Risk In Communities Study). American Journal of Cardiology, 2010, 105, 1420-1425.	1.6	9
175	Reliability and Validity of the Japanese Translated Version of the Swedish Demand-Control-Support Questionnaire. Industrial Health, 2012, 50, 467-475.	1.0	9
176	Fiber Intake and Plasminogen Activator Inhibitor-1 in Type 2 Diabetes: Look AHEAD (Action for Health in) Tj ETQq0	0 0 rgBT 0.8	/Overlock 10 9
177	Phase II multicenter study of adjuvant S-1 for colorectal liver metastasis: survival analysis of N-SOG 01 trial. Cancer Chemotherapy and Pharmacology, 2015, 75, 1281-1288.	2.3	9
178	Calibration between the Estimated Probability of the Risk Assessment Chart of Japan Atherosclerosis Society and Actual Mortality Using External Population: Evidence for Cardiovascular Prevention from Observational Cohorts in Japan (EPOCH-JAPAN). Journal of Atherosclerosis and Thrombosis, 2016, 23, 176-195.	2.0	9
179	Working cancer survivors' physical and mental characteristics compared to cancer-free workers in Japan: a nationwide general population-based study. Journal of Cancer Survivorship, 2021, 15, 912-921.	2.9	9
180	Increased airway hyperresponsiveness to adenosine in patients with aspirin intolerant asthma. Allergology International, 2017, 66, 360-362.	3.3	8

#	Article	IF	CITATIONS
181	Clinical course of human herpesvirus 6 infection in pediatric living donor liver transplantation. Pediatric Transplantation, 2018, 22, e13239.	1.0	8
182	<p>Smoking results in accumulation of ectopic fat in the liver</p> . Diabetes, Metabolic Syndrome and Obesity: Targets and Therapy, 2019, Volume 12, 1075-1080.	2.4	8
183	Risk and population attributable fraction of metabolic syndrome and impaired fasting glucose for the incidence of typeÂ2 diabetes mellitus among middleâ€aged Japanese individuals: Aichi Worker's Cohort Study. Journal of Diabetes Investigation, 2020, 11, 1163-1169.	2.4	8
184	Mortality and Incidence Rates of Stomach Cancer in the JACC Study. Journal of Epidemiology, 2005, 15, S89-S97.	2.4	7
185	Pathophysiologic Mechanisms of Obesity and Related Metabolic Disorders: An Epidemiologic Study using Questionnaire and Serologic Biomarkers. Journal of Epidemiology, 2007, 17, 141-146.	2.4	7
186	Is the association between healthy lifestyle behaviors and cardiovascular mortality modified by overweight status? The Japan Collaborative Cohort Study. Preventive Medicine, 2014, 62, 142-147.	3.4	7
187	What constitutes healthiness of Washoku or Japanese diet?. European Journal of Clinical Nutrition, 2021, 75, 863-864.	2.9	7
188	Work and Family Conflicts, Depression, and "lkigai― A Mediation Analysis in a Cross-cultural Study Between Japanese and Egyptian Civil Workers. Journal of Epidemiology, 2023, 33, 360-366.	2.4	7
189	Lateâ€phase human herpesvirus 6B reactivation in hematopoietic stem cell transplant recipients. Transplant Infectious Disease, 2018, 20, e12916.	1.7	6
190	Milk Intake and Stroke Mortality in the Japan Collaborative Cohort Study—A Bayesian Survival Analysis. Nutrients, 2020, 12, 2743.	4.1	6
191	Having More Healthy Practice was Associated with Low White Blood Cell Counts in Middle-aged Japanese Male and Female Workers. Industrial Health, 2008, 46, 341-347.	1.0	6
192	Increased serum anti-CYP2E1 IgG autoantibody levels may be involved in the pathogenesis of occupational trichloroethylene hypersensitivity syndrome: a case–control study. Archives of Toxicology, 2022, 96, 2785-2797.	4.2	6
193	Association of maternal whole blood fatty acid status during the prenatal period with term birth dimensions: a cross-sectional study. Journal of Perinatal Medicine, 2015, 43, 565-75.	1.4	5
194	Increased risk of occupational trichloroethylene hypersensitivity syndrome at exposure levels higher than 15Âmg/L of urinary trichloroacetic acid, regardless of whether the patients had the HLA-B*13:01 allele. Environmental Research, 2020, 191, 109972.	7.5	5
195	Peanut Consumption and Risk of Stroke and Ischemic Heart Disease in Japanese Men and Women: The JPHC Study. Stroke, 2021, 52, 3543-3550.	2.0	5
196	The association of public trust with the utilization of digital contact tracing for COVID-19 in Japan. Public Health in Practice, 2022, 4, 100279.	1.5	5
197	Body Mass Index-Modified Relationship of Chronic Mental Stress With Resting Blood Pressure During 5 Years in Japanese Middle-Aged Male Workers. Circulation Journal, 2014, 78, 1379-1386.	1.6	4
198	Psychological job strain, social support at work and daytime secretion of dehydroepiandrosterone (DHEA) in healthy female employees: cross-sectional analyses. Scientific Reports, 2015, 5, 15844.	3.3	4

#	Article	IF	CITATIONS
199	Positive Association of Physical Activity with Both Objective and Perceived Measures of the Neighborhood Environment among Older Adults: The Aichi Workers' Cohort Study. International Journal of Environmental Research and Public Health, 2020, 17, 7971.	2.6	4
200	Determinants of alcohol consumption and marijuana use among young adults in the Republic of Palau. Environmental Health and Preventive Medicine, 2021, 26, 12.	3.4	4
201	Perception and practice of 'healthy' diet in relation to noncommunicable diseases among the urban and rural people in northern Ethiopia: a community-based qualitative study. Nagoya Journal of Medical Science, 2018, 80, 451-464.	0.3	4
202	Prediction of Lifetime Risk of Cardiovascular Disease Deaths Stratified by Sex in the Japanese Population. Journal of the American Heart Association, 2021, 10, e021753.	3.7	4
203	Association of Low-Density Lipoprotein Cholesterol with Risk of Coronary Heart Disease and Stroke among Middle-Aged Japanese Workers: An Analysis using Inverse Probability Weighting. Journal of Atherosclerosis and Thrombosis, 2023, 30, 455-466.	2.0	4
204	U-Shaped Association Between White Blood Cell Count and Fasting Plasma Glucose Level. Diabetes Care, 2003, 26, 950-950.	8.6	3
205	Association Between Low Birth Weight and Elevated White Blood Cell Count in Adulthood Within a Japanese Population. Circulation Journal, 2008, 72, 757-763.	1.6	3
206	Descriptive Epidemiology of Hypertension and Its Association With Obesity: Based on the WHO STEPwise Approach to Surveillance in Palau. Asia-Pacific Journal of Public Health, 2017, 29, 278-287.	1.0	3
207	Prevalence and Correlates of Dyslipidemia Among Men and Women in Palau: Findings of the Palau STEPS Survey 2011–2013. Journal of Epidemiology, 2019, 29, 97-103.	2.4	3
208	Apolipoprotein A2 Isoforms in Relation to the Risk of Myocardial Infarction: A Nested Case-Control Analysis in the JPHC Study. Journal of Atherosclerosis and Thrombosis, 2021, 28, 483-490.	2.0	3
209	Pilot peer health education for noncommunicable disease prevention in Bangladesh, Ethiopia, and Palau. Journal of Global Health Reports, 0, 2, .	1.0	3
210	A nested case-control study of stomach cancer incidence and serum superoxide dismutase activity in the Japan Collaborative Cohort study in Japan. Cancer Detection and Prevention, 2007, 31, 431-435.	2.1	2
211	A16358 Hypertension detection, treatment and control rates in urban slum population in Bangladesh. Journal of Hypertension, 2018, 36, e337-e338.	0.5	2
212	Avoid clinical inertia: Importance of asking and advising patients with diabetes who smoke about quitting. Journal of Diabetes Investigation, 2021, 12, 317-319.	2.4	2
213	The association of work-related stress according to the demand-control model with aggravation of pre-existing disease during the first state of COVID-19 emergency in Japan. Journal of Epidemiology, 2021, 31, 642-647.	2.4	2
214	Association of anthropometric indices of obesity with hypertension among public employees in northern Ethiopia: findings from a cross-sectional survey. BMJ Open, 2021, 11, e050969.	1.9	2
215	Relationship Between Fasting Blood Glucose Levels in Middle Age and Cognitive Function in Later Life: The Aichi Workers' Cohort Study. Journal of Epidemiology, 2023, 33, 76-81.	2.4	2
216	Cross-sectional association between working and depression prevalence in cancer survivors: a literature review. Environmental and Occupational Health Practice, 2020, 2, n/a.	0.5	2

#	Article	IF	Citations
217	Work-related factors among people with diabetes and the risk of cardiovascular diseases: A systematic review. Journal of Occupational Health, 2021, 63, e12278.	2.1	2
218	Everyday wishes of older people living with dementia in care planning: a qualitative study. BMC Health Services Research, 2022, 22, 184.	2.2	2
219	We are on the Way to Finding the Cutoff Point for High-Sensitivity C-Reactive Protein in Japanese. Circulation Journal, 2007, 71, 1502.	1.6	1
220	Possible effect of periodic limb movements during sleep on the sleepiness of patients with sleep disordered breathing. Sleep and Biological Rhythms, 2016, 14, 205-210.	1.0	1
221	Glycated haemoglobin (HbA1c) as a reliable option for detecting diabetes among the urban poor population in Bangladesh. European Journal of Public Health, 2020, 30, 839-841.	0.3	1
222	Factors Associated With Prediabetes and Diabetes Among Public Employees in Northern Ethiopia. Asia-Pacific Journal of Public Health, 2021, 33, 242-250.	1.0	1
223	Health of university students under job and financial insecurity during COVID-19 pandemic. Journal of Occupational Health, 2021, 63, e12223.	2.1	1
224	Predictive risk factors for pneumothorax after transbronchial biopsy using endobronchial ultrasonography with a guide sheath. BMC Pulmonary Medicine, 2021, 21, 181.	2.0	1
225	1503The association of work-related stress with aggravation of pre-existing disease during COVID-19 emergency in Japan. International Journal of Epidemiology, 2021, 50, .	1.9	1
226	Having hobbies and the risk of cardiovascular disease incidence: A Japan public health center-based study. Atherosclerosis, 2021, 335, 1-7.	0.8	1
227	Pilot peer health education for noncommunicable disease prevention in Bangladesh, Ethiopia, and Palau. Journal of Global Health Reports, 0, , .	1.0	1
228	Diagnostic contribution of cytological examination to endobronchial ultrasound-guided transbronchial biopsy for lung malignancies. Nagoya Journal of Medical Science, 2019, 81, 613-620.	0.3	1
229	A new method of blood sampling reduces pain for newborn infants: A prospective, randomized controlled clinical trial. Early Human Development, 2007, 83, 389-394.	1.8	0
230	Clinical trial of integrated intervention for Parkinson disease. Neuroscience Research, 2010, 68, e308.	1.9	0
231	Type 2 Diabetes in Well-Controlled Hypertension. Circulation Journal, 2011, 75, 2316-2317.	1.6	0
232	Weight Change and Cardiovascular Mortality. Circulation Journal, 2013, 77, 602-603.	1.6	0
233	The Association between Adult Height and Stroke Incidence in Japanese Men and Women: A Population-Based Case- Control Study. Journal of Epidemiology, 2021, , .	2.4	0
234	Disparity in metabolic risk factors of non-communicable diseases between Palauans and Filipinos living in Palau. Nagoya Journal of Medical Science, 2017, 79, 157-165.	0.3	0

#	Article	IF	CITATIONS
235	Perceptions and behaviors related to noncommunicable diseases in Palau: a qualitative study. Nagoya Journal of Medical Science, 2021, 83, 287-298.	0.3	O
236	Lack of an association between serum level of transforming growth factor beta -1 and stomach cancer risk in the JACC study. Asian Pacific Journal of Cancer Prevention, 2005, 6, 170-6.	1.2	0
237	Applicability of a web-based 24-hour dietary recall tool for Japanese populations in large-scale epidemiological studies. Journal of Epidemiology, 2022, , .	2.4	O
238	Age and sex differences in factors associated with hypertension among an urban poor population in Bangladesh Nagoya Journal of Medical Science, 2022, 84, 69-79.	0.3	0