

David G Johnson

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/5374105/publications.pdf>

Version: 2024-02-01

61
papers

5,656
citations

101543

36
h-index

128289

60
g-index

61
all docs

61
docs citations

61
times ranked

8140
citing authors

#	ARTICLE	IF	CITATIONS
1	Direct Regulation of DNA Repair by E2F and RB in Mammals and Plants: Core Function or Convergent Evolution?. <i>Cancers</i> , 2021, 13, 934.	3.7	3
2	The E2F1 transcription factor and RB tumor suppressor moonlight as DNA repair factors. <i>Cell Cycle</i> , 2020, 19, 2260-2269.	2.6	17
3	E2F1 acetylation directs p300/CBP-mediated histone acetylation at DNA double-strand breaks to facilitate repair. <i>Nature Communications</i> , 2019, 10, 4951.	12.8	45
4	The <i>p53</i> R72P polymorphism does not affect the physiological response to ionizing radiation in a mouse model. <i>Cell Cycle</i> , 2017, 16, 1153-1163.	2.6	1
5	The Retinoblastoma (RB) Tumor Suppressor: Pushing Back against Genome Instability on Multiple Fronts. <i>International Journal of Molecular Sciences</i> , 2017, 18, 1776.	4.1	72
6	RB localizes to DNA double-strand breaks and promotes DNA end resection and homologous recombination through the recruitment of BRG1. <i>Genes and Development</i> , 2016, 30, 2500-2512.	5.9	83
7	Slug Expression in Mouse Skin and Skin Tumors Is Not Regulated by p53. <i>Journal of Investigative Dermatology</i> , 2014, 134, 566-568.	0.7	0
8	E2F1 Responds to Ultraviolet Radiation by Directly Stimulating DNA Repair and Suppressing Carcinogenesis. <i>Cancer Research</i> , 2014, 74, 3369-3377.	0.9	24
9	Identification of prohibitin and prohibiton as novel factors binding to the p53 induced gene 3 (PIG3) promoter (TGYYC)15 motif. <i>Biochemical and Biophysical Research Communications</i> , 2014, 443, 1239-1244.	2.1	20
10	Modeling gene-environment interactions in oral cavity and esophageal cancers demonstrates a role for the p53 R72P polymorphism in modulating susceptibility. <i>Molecular Carcinogenesis</i> , 2014, 53, 648-658.	2.7	10
11	Chromatin: Receiver and Quarterback for Cellular Signals. <i>Cell</i> , 2013, 152, 685-689.	28.9	62
12	E2F1 and p53 Transcription Factors as Accessory Factors for Nucleotide Excision Repair. <i>International Journal of Molecular Sciences</i> , 2012, 13, 13554-13568.	4.1	13
13	Combined effects of <i>E2F1</i> and <i>E2F2</i> polymorphisms on risk and early onset of squamous cell carcinoma of the head and neck. <i>Molecular Carcinogenesis</i> , 2012, 51, E132-41.	2.7	28
14	Transcriptional and Nontranscriptional Functions of E2F1 in Response to DNA Damage. <i>Cancer Research</i> , 2012, 72, 13-17.	0.9	145
15	Effects of <i>MDM2</i> promoter polymorphisms and <i>p53</i> codon 72 polymorphism on risk and age at onset of squamous cell carcinoma of the head and neck. <i>Molecular Carcinogenesis</i> , 2011, 50, 697-706.	2.7	37
16	E2F1 promotes the recruitment of DNA repair factors to sites of DNA double-strand breaks. <i>Cell Cycle</i> , 2011, 10, 1287-1294.	2.6	66
17	GCN5 and E2F1 stimulate nucleotide excision repair by promoting H3K9 acetylation at sites of damage. <i>Nucleic Acids Research</i> , 2011, 39, 1390-1397.	14.5	135
18	Repression of Androgen Receptor Transcription through the E2F1/DNMT1 Axis. <i>PLoS ONE</i> , 2011, 6, e25187.	2.5	25

#	ARTICLE	IF	CITATIONS
19	A High-Frequency Regulatory Polymorphism in the p53 Pathway Accelerates Tumor Development. <i>Cancer Cell</i> , 2010, 18, 220-230.	16.8	108
20	E2F2 suppresses Myc-induced proliferation and tumorigenesis. <i>Molecular Carcinogenesis</i> , 2010, 49, 152-156.	2.7	37
21	E2F1 Localizes to Sites of UV-induced DNA Damage to Enhance Nucleotide Excision Repair. <i>Journal of Biological Chemistry</i> , 2010, 285, 19308-19315.	3.4	55
22	The RB-E2F1 Pathway Regulates Autophagy. <i>Cancer Research</i> , 2010, 70, 7882-7893.	0.9	107
23	INO80 chromatin remodeling complex promotes the removal of UV lesions by the nucleotide excision repair pathway. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2010, 107, 17274-17279.	7.1	87
24	Mouse Models for the p53 R72P Polymorphism Mimic Human Phenotypes. <i>Cancer Research</i> , 2010, 70, 5851-5859.	0.9	49
25	Enhanced skin carcinogenesis and lack of thymus hyperplasia in transgenic mice expressing human cyclin D1b (<i>CCND1b</i>). <i>Molecular Carcinogenesis</i> , 2009, 48, 508-516.	2.7	8
26	E2F1 and E2F3 activate ATM through distinct mechanisms to promote E1A-induced apoptosis. <i>Cell Cycle</i> , 2008, 7, 391-400.	2.6	17
27	Overexpression of the Low Molecular Weight Cyclin E in Transgenic Mice Induces Metastatic Mammary Carcinomas through the Disruption of the ARF-p53 Pathway. <i>Cancer Research</i> , 2007, 67, 7212-7222.	0.9	64
28	Transgenic E2F1 Expression in the Mouse Brain Induces a Human-Like Bimodal Pattern of Tumors. <i>Cancer Research</i> , 2007, 67, 4005-4009.	0.9	29
29	Distinct and Overlapping Roles for E2F Family Members in Transcription, Proliferation and Apoptosis. <i>Current Molecular Medicine</i> , 2006, 6, 739-748.	1.3	38
30	Putting the Oncogenic and Tumor Suppressive Activities of E2F into Context. <i>Current Molecular Medicine</i> , 2006, 6, 731-738.	1.3	1
31	Oncogenes and the DNA Damage Response: Myc and E2F1 Engage the ATM Signaling Pathway to Activate p53 and Induce Apoptosis. <i>Cell Cycle</i> , 2006, 5, 801-803.	2.6	40
32	E2F3a Stimulates Proliferation, p53-Independent Apoptosis and Carcinogenesis in a Transgenic Mouse Model. <i>Cell Cycle</i> , 2006, 5, 184-190.	2.6	43
33	ATM promotes apoptosis and suppresses tumorigenesis in response to Myc. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2006, 103, 1446-1451.	7.1	142
34	Putting the Oncogenic and Tumor Suppressive Activities of E2F into Context. <i>Current Molecular Medicine</i> , 2006, 6, 731-738.	1.3	151
35	Distinct and Overlapping Roles for E2F Family Members in Transcription, Proliferation and Apoptosis. <i>Current Molecular Medicine</i> , 2006, 6, 739-748.	1.3	423
36	SAGE profiling of UV-induced mouse skin squamous cell carcinomas, comparison with acute UV irradiation effects. <i>Molecular Carcinogenesis</i> , 2005, 42, 40-52.	2.7	40

#	ARTICLE	IF	CITATIONS
37	Expression of Transcription Factor E2F1 and Telomerase in Glioblastomas: Mechanistic Linkage and Prognostic Significance. <i>Journal of the National Cancer Institute</i> , 2005, 97, 1589-1600.	6.3	57
38	Regulation of epidermal apoptosis and DNA repair by E2F1 in response to ultraviolet B radiation. <i>Oncogene</i> , 2005, 24, 2449-2460.	5.9	374
39	Lack of Cyclin-Dependent Kinase 4 Inhibits c- myc Tumorigenic Activities in Epithelial Tissues. <i>Molecular and Cellular Biology</i> , 2004, 24, 7538-7547.	2.3	96
40	E2F1 uses the ATM signaling pathway to induce p53 and Chk2 phosphorylation and apoptosis. <i>Molecular Cancer Research</i> , 2004, 2, 203-14.	3.4	54
41	E2F1 Uses the ATM Signaling Pathway to Induce p53 and Chk2 Phosphorylation and Apoptosis. <i>Molecular Cancer Research</i> , 2004, 2, 203-214.	3.4	125
42	Tumor formation in mice with conditional inactivation of Brca1 in epithelial tissues. <i>Oncogene</i> , 2003, 22, 5415-5426.	5.9	38
43	Inactivating E2f1 reverts apoptosis resistance and cancer sensitivity in Trp53-deficient mice. <i>Nature Cell Biology</i> , 2003, 5, 655-660.	10.3	391
44	ARF Differentially Modulates Apoptosis Induced by E2F1 and Myc. <i>Molecular and Cellular Biology</i> , 2002, 22, 1360-1368.	2.3	75
45	Inactivation of E2f1 enhances tumorigenesis in a Myc transgenic model. <i>Cancer Research</i> , 2002, 62, 3276-81.	0.9	34
46	Molecular Cloning and Characterization of a Novel Mouse Epidermal Differentiation Gene and Its Promoter. <i>Genomics</i> , 2001, 73, 284-290.	2.9	13
47	Deregulated expression of DP1 induces epidermal proliferation and enhances skin carcinogenesis. <i>Molecular Carcinogenesis</i> , 2001, 31, 90-100.	2.7	27
48	Myc lacks E2F1's ability to suppress skin carcinogenesis. <i>Oncogene</i> , 2001, 20, 5341-5349.	5.9	36
49	The paradox of E2F1: Oncogene and tumor suppressor gene. <i>Molecular Carcinogenesis</i> , 2000, 27, 151-157.	2.7	85
50	Regulation of BRCA1 Expression by the Rb-E2F Pathway. <i>Journal of Biological Chemistry</i> , 2000, 275, 4532-4536.	3.4	108
51	E2F4 and E2F1 Have Similar Proliferative Properties but Different Apoptotic and Oncogenic Properties In Vivo. <i>Molecular and Cellular Biology</i> , 2000, 20, 3417-3424.	2.3	87
52	E2F4 and E2F1 Have Similar Proliferative Properties but Different Apoptotic and Oncogenic Properties In Vivo. <i>Molecular and Cellular Biology</i> , 2000, 20, 3417-3424.	2.3	11
53	E2F1 Has Both Oncogenic and Tumor-Suppressive Properties in a Transgenic Model. <i>Molecular and Cellular Biology</i> , 1999, 19, 6408-6414.	2.3	136
54	Deregulated expression of E2F1 induces hyperplasia and cooperates with ras in skin tumor development. <i>Oncogene</i> , 1998, 16, 1267-1276.	5.9	125

#	ARTICLE	IF	CITATIONS
55	Deregulated expression of cell-cycle proteins during premalignant progression in SENCAR mouse skin. <i>Oncogene</i> , 1998, 17, 2251-2258.	5.9	34
56	Differential activities of E2F family members: Unique functions in regulating transcription. <i>Molecular Carcinogenesis</i> , 1998, 22, 190-198.	2.7	35
57	Increased E2F1 activity induces skin tumors in mice heterozygous and nullizygous for p53. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1998, 95, 8858-8863.	7.1	153
58	Role of E2F in cell cycle control and cancer. <i>Frontiers in Bioscience - Landmark</i> , 1998, 3, d447-458.	3.0	166
59	Oncogenic capacity of the E2F1 gene.. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1994, 91, 12823-12827.	7.1	234
60	Expression of transcription factor E2F1 induces quiescent cells to enter S phase. <i>Nature</i> , 1993, 365, 349-352.	27.8	935
61	Variability of Alkaloid Production in Submerged Culture. <i>Quarterly Journal of Crude Drug Research = Vierteljahrliche Zietschrift Fur Drogen-Forschung = Revue Trimestrielle Des Recherches Sur Les Matieres Premieres</i> , 1964, 4, 577-581.	0.2	2