

Antoni Serrano-Blanco

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/537411/publications.pdf>

Version: 2024-02-01

114
papers

4,123
citations

117625

34
h-index

144013

57
g-index

131
all docs

131
docs citations

131
times ranked

5717
citing authors

#	ARTICLE	IF	CITATIONS
1	Assessing depression in primary care with the PHQ-9: Can it be carried out over the telephone?. <i>Journal of General Internal Medicine</i> , 2005, 20, 738-742.	2.6	298
2	Activity monitoring in patients with depression: A systematic review. <i>Journal of Affective Disorders</i> , 2013, 145, 21-28.	4.1	193
3	Prevalence of mental disorders in primary care: results from the diagnosis and treatment of mental disorders in primary care study (DASMAP). <i>Social Psychiatry and Psychiatric Epidemiology</i> , 2010, 45, 201-210.	3.1	137
4	Risk factors for the onset of panic and generalised anxiety disorders in the general adult population: A systematic review of cohort studies. <i>Journal of Affective Disorders</i> , 2014, 168, 337-348.	4.1	130
5	Effectiveness of group acceptance and commitment therapy for fibromyalgia: A 6-month randomized controlled trial (EFFIGACT study). <i>Pain</i> , 2014, 155, 693-702.	4.2	125
6	La carga de enfermedad en España: resultados del Estudio de la Carga Global de las Enfermedades 2016. <i>Medicina Clínica</i> , 2018, 151, 171-190.	0.6	113
7	Prevalence and variability of current depressive disorder in 27 European countries: a population-based study. <i>Lancet Public Health</i> , The, 2021, 6, e729-e738.	10.0	109
8	Perceived injustice in fibromyalgia: Psychometric characteristics of the Injustice Experience Questionnaire and relationship with pain catastrophising and pain acceptance. <i>Journal of Psychosomatic Research</i> , 2012, 73, 86-91.	2.6	105
9	Pilot randomised controlled trial of Help4Mood, an embodied virtual agent-based system to support treatment of depression. <i>Journal of Telemedicine and Telecare</i> , 2016, 22, 348-355.	2.7	94
10	Bifactor analysis and construct validity of the five facet mindfulness questionnaire (FFMQ) in non-clinical Spanish samples. <i>Frontiers in Psychology</i> , 2015, 6, 404.	2.1	90
11	Raloxifene as an Adjunctive Treatment for Postmenopausal Women With Schizophrenia. <i>Journal of Clinical Psychiatry</i> , 2011, 72, 1552-1557.	2.2	88
12	Is major depression adequately diagnosed and treated by general practitioners? Results from an epidemiological study. <i>General Hospital Psychiatry</i> , 2010, 32, 201-209.	2.4	81
13	Gender differences in response to antidepressant treatment prescribed in primary care. Does menopause make a difference?. <i>Journal of Affective Disorders</i> , 2006, 93, 53-60.	4.1	79
14	Psychometric properties of the twelve item World Health Organization Disability Assessment Schedule II (WHO-DAS II) in Spanish primary care patients with a first major depressive episode. <i>Journal of Affective Disorders</i> , 2010, 121, 52-58.	4.1	78
15	Effectiveness of Pharmacist Care in the Improvement of Adherence to Antidepressants: A Systematic Review and Meta-Analysis. <i>Annals of Pharmacotherapy</i> , 2011, 45, 39-48.	1.9	75
16	Brain dysfunction in fibromyalgia and somatization disorder using proton magnetic resonance spectroscopy: a controlled study. <i>Acta Psychiatrica Scandinavica</i> , 2012, 126, 115-125.	4.5	75
17	Adjustment disorders in primary care: prevalence, recognition and use of services. <i>British Journal of Psychiatry</i> , 2012, 201, 137-142.	2.8	73
18	The 12-item World Health Organization Disability Assessment Schedule II (WHO-DAS II): a nonparametric item response analysis. <i>BMC Medical Research Methodology</i> , 2010, 10, 45.	3.1	67

#	ARTICLE	IF	CITATIONS
19	Comorbidity between common mental disorders and chronic somatic diseases in primary care patients. <i>General Hospital Psychiatry</i> , 2010, 32, 240-245.	2.4	67
20	Effects of raloxifene on cognition in postmenopausal women with schizophrenia: A double-blind, randomized, placebo-controlled trial. <i>European Neuropsychopharmacology</i> , 2014, 24, 223-231.	0.7	66
21	Validation of the Spanish version of the Chronic Pain Acceptance Questionnaire (CPAQ) for the assessment of acceptance in fibromyalgia. <i>Health and Quality of Life Outcomes</i> , 2010, 8, 37.	2.4	62
22	Effectiveness of a Psychoeducational Treatment Program Implemented in General Practice for Fibromyalgia Patients. <i>Clinical Journal of Pain</i> , 2011, 27, 383-391.	1.9	60
23	Factor structure, internal consistency and construct validity of the Sheehan Disability Scale in a Spanish primary care sample. <i>Journal of Evaluation in Clinical Practice</i> , 2010, 16, 895-901.	1.8	54
24	Effectiveness of cognitive behaviour therapy for the treatment of catastrophisation in patients with fibromyalgia: a randomised controlled trial. <i>Arthritis Research and Therapy</i> , 2011, 13, R173.	3.5	53
25	Cost-Utility of a Psychoeducational Intervention in Fibromyalgia Patients Compared With Usual Care. <i>Clinical Journal of Pain</i> , 2013, 29, 702-711.	1.9	51
26	Health service use and costs associated with aggressiveness or agitation and containment in adult psychiatric care: a systematic review of the evidence. <i>BMC Psychiatry</i> , 2015, 15, 35.	2.6	48
27	Frequent attendance in primary care: comparison and implications of different definitions. <i>British Journal of General Practice</i> , 2010, 60, e49-e55.	1.4	45
28	Relationship between behavioural coping strategies and acceptance in patients with fibromyalgia syndrome: Elucidating targets of interventions. <i>BMC Musculoskeletal Disorders</i> , 2011, 12, 143.	1.9	42
29	Evaluation of a pharmacist intervention on patients initiating pharmacological treatment for depression: A randomized controlled superiority trial. <i>European Neuropsychopharmacology</i> , 2013, 23, 1057-1066.	0.7	41
30	Cost-utility of cognitive behavioral therapy versus U.S. Food and Drug Administration recommended drugs and usual care in the treatment of patients with fibromyalgia: an economic evaluation alongside a 6-month randomized controlled trial. <i>Arthritis Research and Therapy</i> , 2014, 16, 451.	3.5	41
31	Effectiveness of psychological and/or educational interventions to prevent the onset of episodes of depression: A systematic review of systematic reviews and meta-analyses. <i>Preventive Medicine</i> , 2015, 76, S22-S32.	3.4	41
32	Brain Changes in Long-Term Zen Meditators Using Proton Magnetic Resonance Spectroscopy and Diffusion Tensor Imaging: A Controlled Study. <i>PLoS ONE</i> , 2013, 8, e58476.	2.5	40
33	Initial medication non-adherence: prevalence and predictive factors in a cohort of 1.6 million primary care patients. <i>British Journal of Clinical Pharmacology</i> , 2017, 83, 1328-1340.	2.4	37
34	The burden of disease in Spain: Results from the Global Burden of Disease 2016. <i>Medicina Clínica (English Edition)</i> , 2018, 151, 171-190.	0.2	37
35	Brain dysfunction behind functional symptoms: neuroimaging and somatoform, conversive, and dissociative disorders. <i>Current Opinion in Psychiatry</i> , 2009, 22, 224-231.	6.3	36
36	Cost-Utility of Group Acceptance and Commitment Therapy for Fibromyalgia Versus Recommended Drugs: An Economic Analysis Alongside a 6-Month Randomized Controlled Trial Conducted in Spain (EFFIGACT Study). <i>Journal of Pain</i> , 2017, 18, 868-880.	1.4	35

#	ARTICLE	IF	CITATIONS
37	Cost-utility and biological underpinnings of Mindfulness-Based Stress Reduction (MBSR) versus a psychoeducational programme (FibroQoL) for fibromyalgia: a 12-month randomised controlled trial (EUDAIMON study). <i>BMC Complementary and Alternative Medicine</i> , 2016, 16, 81.	3.7	34
38	Prevalence and age patterns of depression in the United Kingdom. A population-based study. <i>Journal of Affective Disorders</i> , 2021, 279, 164-172.	4.1	34
39	Functional Status, Quality of Life, and Costs Associated With Fibromyalgia Subgroups. <i>Clinical Journal of Pain</i> , 2016, 32, 829-840.	1.9	33
40	Antipsychotic polypharmacy in a regional health service: a population-based study. <i>BMC Psychiatry</i> , 2012, 12, 42.	2.6	32
41	Use of an integrated Atlas of Mental Health Care for evidence informed policy in Catalonia (Spain). <i>Epidemiology and Psychiatric Sciences</i> , 2015, 24, 512-524.	3.9	31
42	Cost-Effectiveness of a Community Pharmacist Intervention in Patients with Depression: A Randomized Controlled Trial (PRODEFAR Study). <i>PLoS ONE</i> , 2013, 8, e70588.	2.5	31
43	Utility of the twelve-item World Health Organization Disability Assessment Schedule II (WHO-DAS II) for discriminating depression and severity in Spanish primary care patients. <i>Quality of Life Research</i> , 2010, 19, 97-101.	3.1	30
44	Recognition of anxiety disorders by the general practitioner: results from the DASMAPP Study. <i>General Hospital Psychiatry</i> , 2012, 34, 227-233.	2.4	30
45	Relationship Between Depression and the Use of Mobile Technologies and Social Media Among Adolescents: Umbrella Review. <i>Journal of Medical Internet Research</i> , 2020, 22, e16388.	4.3	30
46	Dimensionality, Reliability, and Validity of the Revised Fibromyalgia Impact Questionnaire in Two Spanish Samples. <i>Arthritis Care and Research</i> , 2013, 65, 1682-1689.	3.4	29
47	Validation of a Spanish version of the psychological inflexibility in pain scale (PIPS) and an evaluation of its relation with acceptance of pain and mindfulness in sample of persons with fibromyalgia. <i>Health and Quality of Life Outcomes</i> , 2013, 11, 62.	2.4	28
48	Adequacy of Antidepressant Treatment in Spanish Primary Care: A Naturalistic Six-Month Follow-Up Study. <i>Psychiatric Services</i> , 2008, 59, 78-83.	2.0	27
49	Prevalence of mental disorders in non-demented elderly people in primary care. <i>International Psychogeriatrics</i> , 2015, 27, 757-768.	1.0	27
50	Effectiveness of the psychological and pharmacological treatment of catastrophization in patients with fibromyalgia: a randomized controlled trial. <i>Trials</i> , 2009, 10, 24.	1.6	23
51	Intervention to Prevent Major Depression in Primary Care. <i>Annals of Internal Medicine</i> , 2016, 164, 656.	3.9	23
52	Impact of the mass media OBERTAMENT campaign on the levels of stigma among the population of Catalonia, Spain. <i>European Psychiatry</i> , 2016, 31, 44-51.	0.2	23
53	Low intensity vs. self-guided Internet-delivered psychotherapy for major depression: a multicenter, controlled, randomized study. <i>BMC Psychiatry</i> , 2013, 13, 21.	2.6	22
54	Epidemiology of psychiatric morbidity among migrants compared to native born population in Spain: a controlled study. <i>General Hospital Psychiatry</i> , 2013, 35, 93-99.	2.4	22

#	ARTICLE	IF	CITATIONS
55	Once-monthly paliperidone palmitate in recently diagnosed and chronic non-acute patients with schizophrenia. <i>Expert Opinion on Pharmacotherapy</i> , 2016, 17, 1043-1053.	1.8	22
56	Prevalence and characteristics of catatonia on admission to an acute geriatric psychiatry ward. <i>Progress in Neuro-Psychopharmacology and Biological Psychiatry</i> , 2017, 78, 27-33.	4.8	22
57	In-patient costs of agitation and containment in a mental health catchment area. <i>BMC Psychiatry</i> , 2017, 17, 212.	2.6	22
58	Effect of copayment policies on initial medication non-adherence according to income: a population-based study. <i>BMJ Quality and Safety</i> , 2018, 27, 878-891.	3.7	22
59	Cost?utility of selective serotonin reuptake inhibitors for depression in primary care in Catalonia. <i>Acta Psychiatrica Scandinavica</i> , 2006, 114, 39-47.	4.5	21
60	Community pharmacist intervention in depressed primary care patients (PRODEFAR study): randomized controlled trial protocol. <i>BMC Public Health</i> , 2009, 9, 284.	2.9	21
61	Impact of mental disorders and chronic physical conditions in health-related quality of life among primary care patients: results from an epidemiological study. <i>Quality of Life Research</i> , 2009, 18, 1011-1018.	3.1	21
62	Psychotropic prescribing in Catalonia: results from an epidemiological study. <i>Family Practice</i> , 2012, 29, 154-162.	1.9	21
63	A personalized intervention to prevent depression in primary care: cost-effectiveness study nested into a clustered randomized trial. <i>BMC Medicine</i> , 2018, 16, 28.	5.5	21
64	Preventing the onset of major depression based on the level and profile of risk of primary care attendees: protocol of a cluster randomised trial (the predictD-CCRT study). <i>BMC Psychiatry</i> , 2013, 13, 171.	2.6	20
65	Attitudes and intended behaviour to mental disorders and associated factors in catalan population, Spain: cross-sectional population-based survey. <i>BMC Public Health</i> , 2015, 16, 127.	2.9	20
66	Expectations, experiences and attitudes of patients and primary care health professionals regarding online psychotherapeutic interventions for depression: protocol for a qualitative study. <i>BMC Psychiatry</i> , 2013, 13, 64.	2.6	19
67	Effectiveness and cost-effectiveness of antidepressant treatment in primary health care: A six-month randomised study comparing fluoxetine to imipramine. <i>Journal of Affective Disorders</i> , 2006, 91, 153-163.	4.1	18
68	Factors associated with use of community mental health services by schizophrenia patients using multilevel analysis. <i>BMC Health Services Research</i> , 2011, 11, 257.	2.2	18
69	Impact of initial medication non-adherence on use of healthcare services and sick leave: a longitudinal study in a large primary care cohort in Spain. <i>British Journal of General Practice</i> , 2017, 67, e614-e622.	1.4	18
70	Reducing the Mental Healthâ€“Related Stigma of Social Work Students. <i>Research on Social Work Practice</i> , 2018, 28, 164-172.	1.9	18
71	Patientsâ€™ Opinions about Knowing Their Risk for Depression and What to Do about It. The PredictD-Qualitative Study. <i>PLoS ONE</i> , 2014, 9, e92008.	2.5	15
72	Gender differences in the impact of mental disorders and chronic physical conditions on health-related quality of life among non-demented primary care elderly patients. <i>Quality of Life Research</i> , 2016, 25, 1461-1474.	3.1	15

#	ARTICLE	IF	CITATIONS
73	Quality of Life and Adverse Effects of Olanzapine Versus Risperidone Therapy in Patients with Schizophrenia. <i>Psychiatric Quarterly</i> , 2013, 84, 125-135.	2.1	14
74	Is there a case for mental health promotion in the primary care setting? A systematic review. <i>Preventive Medicine</i> , 2015, 76, S5-S11.	3.4	13
75	Complex multiple risk intervention to promote healthy behaviours in people between 45 to 75 years attended in primary health care (EIRA study): study protocol for a hybrid trial. <i>BMC Public Health</i> , 2018, 18, 874.	2.9	13
76	Effectiveness of the "What's Up!" Intervention to Reduce Stigma and Psychometric Properties of the Youth Program Questionnaire (YPQ): Results from a Cluster Non-randomized Controlled Trial Conducted in Catalan High Schools. <i>Frontiers in Psychology</i> , 2017, 8, 1608.	2.1	12
77	Catatonia in elderly psychiatric inpatients is not always associated with intense anxiety: Factor analysis and correlation with psychopathology. <i>International Journal of Geriatric Psychiatry</i> , 2020, 35, 1409-1417.	2.7	12
78	Cost-effectiveness of active monitoring versus antidepressants for major depression in primary health care: a 12-month non-randomized controlled trial (INFAP study). <i>BMC Psychiatry</i> , 2015, 15, 63.	2.6	11
79	Impact of initial medication non-adherence to SSRIs on medical visits and sick leaves. <i>Journal of Affective Disorders</i> , 2018, 226, 282-286.	4.1	11
80	Changes in metabolites in the brain of patients with fibromyalgia after treatment with an NMDA receptor antagonist. <i>Neuroradiology Journal</i> , 2019, 32, 408-419.	1.2	11
81	Comparing watchful waiting with antidepressants for the management of subclinical depression symptoms to mild to moderate depression in primary care: a systematic review. <i>Family Practice</i> , 2017, 34, 639-648.	1.9	10
82	Effectiveness of watchful waiting versus antidepressants for patients diagnosed of mild to moderate depression in primary care: A 12-month pragmatic clinical trial (INFAP study). <i>European Psychiatry</i> , 2018, 53, 66-73.	0.2	10
83	Fluoxetine and imipramine: are there differences in cost-utility for depression in primary care?. <i>Journal of Evaluation in Clinical Practice</i> , 2009, 15, 195-203.	1.8	9
84	Adequacy of Antidepressant Treatment in Spanish Primary Care: A Naturalistic Six-Month Follow-Up Study. <i>Psychiatric Services</i> , 2008, 59, 78-83.	2.0	9
85	Clinical dimensions of fibromyalgia symptoms and development of a combined index of severity: The CODI index. <i>Quality of Life Research</i> , 2013, 22, 153-160.	3.1	8
86	Digging into the construct of fibrofog: Psychometric properties of the Spanish version of the Multidimensional Inventory of Subjective Cognitive Impairment in patients with fibromyalgia. <i>Journal of Applied Biobehavioral Research</i> , 2018, 23, e12134.	2.0	8
87	Investigación traslacional en psiquiatría: el marco Research Domain Criteria (RDoC). <i>Revista De Psiquiatría Y Salud Mental</i> , 2019, 12, 187-195.	1.8	8
88	Accuracy of Self-Reported Items for the Screening of Depression in the General Population. <i>International Journal of Environmental Research and Public Health</i> , 2020, 17, 7955.	2.6	8
89	An integrative cross-design synthesis approach to estimate the cost of illness: An applied case to the cost of depression in Catalonia. <i>Journal of Mental Health</i> , 2013, 22, 135-154.	1.9	7
90	PHQ-8 scores and estimation of depression prevalence – Author's reply. <i>Lancet Public Health</i> , The, 2021, 6, e794.	10.0	7

#	ARTICLE	IF	CITATIONS
91	Qualitative study of the agitation states and their characterization, and the interventions used to attend them. <i>Actas Espanolas De Psiquiatria</i> , 2016, 44, 166-77.	0.1	7
92	Evaluaci3n telef3nica de los sistemas de activaci3n e inhibici3n conductual. <i>Revista De Psiquiatr3a Y Salud Mental</i> , 2009, 2, 66-71.	1.8	6
93	Cost-effectiveness of antidepressants versus active monitoring for mild-to-moderate major depressive disorder: a multisite non-randomized-controlled trial in primary care (INFAP study). <i>European Journal of Health Economics</i> , 2019, 20, 703-713.	2.8	6
94	Identification of the most vulnerable populations in the psychosocial sphere: a cross-sectional study conducted in Catalonia during the strict lockdown imposed against the COVID-19 pandemic. <i>BMJ Open</i> , 2021, 11, e052140.	1.9	6
95	Effectiveness of a psycho-educational program for improving quality of life of fibromyalgia patients. <i>BMC Musculoskeletal Disorders</i> , 2008, 9, .	1.9	5
96	Perfil psicol3gico diferencial entre dolor tor3cico de causa no cardiol3gica y enfermedad coronaria: un estudio controlado. <i>Revista Espanola De Cardiologia</i> , 2010, 63, 357-361.	1.2	5
97	Quality of Life and Hormonal, Biochemical, and Anthropometric Profile Between Olanzapine and Risperidone Users. <i>Psychiatric Quarterly</i> , 2016, 87, 293-304.	2.1	5
98	Diagnostic promiscuity: the use of real-world data to study multimorbidity in mental health. <i>British Journal of Psychiatry</i> , 2021, 218, 237-239.	2.8	5
99	Cost-effectiveness analysis of a multiple health behaviour change intervention in people aged between 45 and 75 years: a cluster randomized controlled trial in primary care (EIRA study). <i>International Journal of Behavioral Nutrition and Physical Activity</i> , 2021, 18, 88.	4.6	5
100	Effectiveness and cost-effectiveness of an intervention to improve Initial Medication Adherence to treatments for cardiovascular diseases and diabetes in primary care: study protocol for a pragmatic cluster randomised controlled trial and economic model (the IMA-cRCT study). , 2022, 23, .		5
101	A model for continuous monitoring of patients with major depression in short and long term periods. <i>Technology and Health Care</i> , 2017, 25, 487-511.	1.2	4
102	Diagnostic accuracy and treatment approach to depression in primary care: predictive factors. <i>Family Practice</i> , 2019, 36, 3-11.	1.9	4
103	Initiation and Single Dispensing in Cardiovascular and Insulin Medications: Prevalence and Explanatory Factors. <i>International Journal of Environmental Research and Public Health</i> , 2020, 17, 3358.	2.6	4
104	Non-initiation of prescribed medication from a Spanish health professionals' perspective: A qualitative exploration based on Grounded Theory. <i>Health and Social Care in the Community</i> , 2022, 30, e213-e221.	1.6	4
105	Intellectual Disability, Autism Spectrum Disorders, Psychiatric Comorbidities and Their Relationship with Challenging Behavior. <i>Journal of Mental Health Research in Intellectual Disabilities</i> , 0, , 1-18.	2.0	4
106	Improving suicide surveillance systems through the use of the Patient Health Questionnaire-9.. <i>Journal of Affective Disorders</i> , 2021, 293, 71-72.	4.1	3
107	Selective serotonin reuptake inhibitors: are there differences in effectiveness in primary care?. <i>Primary Care and Community Psychiatry</i> , 2007, 11, 113-120.	0.4	3
108	MADRIM: A major depression remote intelligent monitor. , 2012, , .		2

#	ARTICLE	IF	CITATIONS
109	Use of Mental Health Services for Patients Diagnosed with Major Depressive Disorders in Primary Care. <i>International Journal of Environmental Research and Public Health</i> , 2021, 18, 885.	2.6	2
110	¿C�mo derivamos a salud mental desde atenci�n primaria?. <i>Atencion Primaria</i> , 2003, 32, 524-530.	1.4	2
111	Evaluation of behavioural activation and inhibition systems by telephone. <i>Revista De Psiquiatr�a Y Salud Mental (English Edition)</i> , 2009, 2, 66-71.	0.3	1
112	Assessment of a Primary Support Program: family physicians and mental health professionals�™ perspective. <i>Family Practice</i> , 2022, 39, 920-931.	1.9	1
113	Impact of the 13-Valent Conjugated Pneumococcal Vaccine on the Direct Costs of Invasive Pneumococcal Disease Requiring Hospital Admission in Children Aged < 5 Years: A Prospective Study. <i>Vaccines</i> , 2020, 8, 387.	4.4	0
114	Cost-effectiveness and cost-utility of a community pharmacist intervention in patients with depression (PRODEFAR study). <i>International Journal of Integrated Care</i> , 2014, 14, .	0.2	0