

Antoni Serrano-Blanco

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/537411/publications.pdf>

Version: 2024-02-01

114
papers

4,123
citations

117625

34
h-index

144013

57
g-index

131
all docs

131
docs citations

131
times ranked

5717
citing authors

#	ARTICLE	IF	CITATIONS
1	Non-initiation of prescribed medication from a Spanish health professionals' perspective: A qualitative exploration based on Grounded Theory. <i>Health and Social Care in the Community</i> , 2022, 30, e213-e221.	1.6	4
2	Assessment of a Primary Support Program: family physicians and mental health professionals' perspective. <i>Family Practice</i> , 2022, 39, 920-931.	1.9	1
3	Effectiveness and cost-effectiveness of an intervention to improve Initial Medication Adherence to treatments for cardiovascular diseases and diabetes in primary care: study protocol for a pragmatic cluster randomised controlled trial and economic model (the IMA-cRCT study). , 2022, 23, .		5
4	Prevalence and age patterns of depression in the United Kingdom. A population-based study. <i>Journal of Affective Disorders</i> , 2021, 279, 164-172.	4.1	34
5	Diagnostic promiscuity: the use of real-world data to study multimorbidity in mental health. <i>British Journal of Psychiatry</i> , 2021, 218, 237-239.	2.8	5
6	Use of Mental Health Services for Patients Diagnosed with Major Depressive Disorders in Primary Care. <i>International Journal of Environmental Research and Public Health</i> , 2021, 18, 885.	2.6	2
7	Cost-effectiveness analysis of a multiple health behaviour change intervention in people aged between 45 and 75 years: a cluster randomized controlled trial in primary care (EIRA study). <i>International Journal of Behavioral Nutrition and Physical Activity</i> , 2021, 18, 88.	4.6	5
8	Prevalence and variability of current depressive disorder in 27 European countries: a population-based study. <i>Lancet Public Health</i> , The, 2021, 6, e729-e738.	10.0	109
9	Improving suicide surveillance systems through the use of the Patient Health Questionnaire-9.. <i>Journal of Affective Disorders</i> , 2021, 293, 71-72.	4.1	3
10	PHQ-8 scores and estimation of depression prevalence – Author's reply. <i>Lancet Public Health</i> , The, 2021, 6, e794.	10.0	7
11	Identification of the most vulnerable populations in the psychosocial sphere: a cross-sectional study conducted in Catalonia during the strict lockdown imposed against the COVID-19 pandemic. <i>BMJ Open</i> , 2021, 11, e052140.	1.9	6
12	Impact of the 13-Valent Conjugated Pneumococcal Vaccine on the Direct Costs of Invasive Pneumococcal Disease Requiring Hospital Admission in Children Aged ≤ 5 Years: A Prospective Study. <i>Vaccines</i> , 2020, 8, 387.	4.4	0
13	Catatonia in elderly psychiatric inpatients is not always associated with intense anxiety: Factor analysis and correlation with psychopathology. <i>International Journal of Geriatric Psychiatry</i> , 2020, 35, 1409-1417.	2.7	12
14	Accuracy of Self-Reported Items for the Screening of Depression in the General Population. <i>International Journal of Environmental Research and Public Health</i> , 2020, 17, 7955.	2.6	8
15	Initiation and Single Dispensing in Cardiovascular and Insulin Medications: Prevalence and Explanatory Factors. <i>International Journal of Environmental Research and Public Health</i> , 2020, 17, 3358.	2.6	4
16	Relationship Between Depression and the Use of Mobile Technologies and Social Media Among Adolescents: Umbrella Review. <i>Journal of Medical Internet Research</i> , 2020, 22, e16388.	4.3	30
17	Investigación traslacional en psiquiatría: el marco Research Domain Criteria (RDoC). <i>Revista De Psiquiatría Y Salud Mental</i> , 2019, 12, 187-195.	1.8	8
18	Changes in metabolites in the brain of patients with fibromyalgia after treatment with an NMDA receptor antagonist. <i>Neuroradiology Journal</i> , 2019, 32, 408-419.	1.2	11

#	ARTICLE	IF	CITATIONS
19	Cost-effectiveness of antidepressants versus active monitoring for mild-to-moderate major depressive disorder: a multisite non-randomized-controlled trial in primary care (INFAP study). <i>European Journal of Health Economics</i> , 2019, 20, 703-713.	2.8	6
20	Diagnostic accuracy and treatment approach to depression in primary care: predictive factors. <i>Family Practice</i> , 2019, 36, 3-11.	1.9	4
21	A personalized intervention to prevent depression in primary care: cost-effectiveness study nested into a clustered randomized trial. <i>BMC Medicine</i> , 2018, 16, 28.	5.5	21
22	Effect of copayment policies on initial medication non-adherence according to income: a population-based study. <i>BMJ Quality and Safety</i> , 2018, 27, 878-891.	3.7	22
23	Reducing the Mental Health-Related Stigma of Social Work Students. <i>Research on Social Work Practice</i> , 2018, 28, 164-172.	1.9	18
24	Impact of initial medication non-adherence to SSRIs on medical visits and sick leaves. <i>Journal of Affective Disorders</i> , 2018, 226, 282-286.	4.1	11
25	Complex multiple risk intervention to promote healthy behaviours in people between 45 to 75 years attended in primary health care (EIRA study): study protocol for a hybrid trial. <i>BMC Public Health</i> , 2018, 18, 874.	2.9	13
26	The burden of disease in Spain: Results from the Global Burden of Disease 2016. <i>Medicina Clínica (English Edition)</i> , 2018, 151, 171-190.	0.2	37
27	Effectiveness of watchful waiting versus antidepressants for patients diagnosed of mild to moderate depression in primary care: A 12-month pragmatic clinical trial (INFAP study). <i>European Psychiatry</i> , 2018, 53, 66-73.	0.2	10
28	Digging into the construct of fibrofog: Psychometric properties of the Spanish version of the Multidimensional Inventory of Subjective Cognitive Impairment in patients with fibromyalgia. <i>Journal of Applied Biobehavioral Research</i> , 2018, 23, e12134.	2.0	8
29	La carga de enfermedad en España: resultados del Estudio de la Carga Global de las Enfermedades 2016. <i>Medicina Clínica</i> , 2018, 151, 171-190.	0.6	113
30	Initial medication non-adherence: prevalence and predictive factors in a cohort of 1.6 million primary care patients. <i>British Journal of Clinical Pharmacology</i> , 2017, 83, 1328-1340.	2.4	37
31	A model for continuous monitoring of patients with major depression in short and long term periods. <i>Technology and Health Care</i> , 2017, 25, 487-511.	1.2	4
32	Comparing watchful waiting with antidepressants for the management of subclinical depression symptoms to mild-to-moderate depression in primary care: a systematic review. <i>Family Practice</i> , 2017, 34, 639-648.	1.9	10
33	Prevalence and characteristics of catatonia on admission to an acute geriatric psychiatry ward. <i>Progress in Neuro-Psychopharmacology and Biological Psychiatry</i> , 2017, 78, 27-33.	4.8	22
34	Cost-Utility of Group Acceptance and Commitment Therapy for Fibromyalgia Versus Recommended Drugs: An Economic Analysis Alongside a 6-Month Randomized Controlled Trial Conducted in Spain (EFFIGACT Study). <i>Journal of Pain</i> , 2017, 18, 868-880.	1.4	35
35	Impact of initial medication non-adherence on use of healthcare services and sick leave: a longitudinal study in a large primary care cohort in Spain. <i>British Journal of General Practice</i> , 2017, 67, e614-e622.	1.4	18
36	In-patient costs of agitation and containment in a mental health catchment area. <i>BMC Psychiatry</i> , 2017, 17, 212.	2.6	22

#	ARTICLE	IF	CITATIONS
37	Effectiveness of the "What's Up!" Intervention to Reduce Stigma and Psychometric Properties of the Youth Program Questionnaire (YPQ): Results from a Cluster Non-randomized Controlled Trial Conducted in Catalan High Schools. <i>Frontiers in Psychology</i> , 2017, 8, 1608.	2.1	12
38	Intervention to Prevent Major Depression in Primary Care. <i>Annals of Internal Medicine</i> , 2016, 164, 656.	3.9	23
39	Once-monthly paliperidone palmitate in recently diagnosed and chronic non-acute patients with schizophrenia. <i>Expert Opinion on Pharmacotherapy</i> , 2016, 17, 1043-1053.	1.8	22
40	Cost-utility and biological underpinnings of Mindfulness-Based Stress Reduction (MBSR) versus a psychoeducational programme (FibroQoL) for fibromyalgia: a 12-month randomised controlled trial (EUDAIMON study). <i>BMC Complementary and Alternative Medicine</i> , 2016, 16, 81.	3.7	34
41	Functional Status, Quality of Life, and Costs Associated With Fibromyalgia Subgroups. <i>Clinical Journal of Pain</i> , 2016, 32, 829-840.	1.9	33
42	Impact of the mass media OBERTAMENT campaign on the levels of stigma among the population of Catalonia, Spain. <i>European Psychiatry</i> , 2016, 31, 44-51.	0.2	23
43	Gender differences in the impact of mental disorders and chronic physical conditions on health-related quality of life among non-demented primary care elderly patients. <i>Quality of Life Research</i> , 2016, 25, 1461-1474.	3.1	15
44	Pilot randomised controlled trial of Help4Mood, an embodied virtual agent-based system to support treatment of depression. <i>Journal of Telemedicine and Telecare</i> , 2016, 22, 348-355.	2.7	94
45	Quality of Life and Hormonal, Biochemical, and Anthropometric Profile Between Olanzapine and Risperidone Users. <i>Psychiatric Quarterly</i> , 2016, 87, 293-304.	2.1	5
46	Qualitative study of the agitation states and their characterization, and the interventions used to attend them. <i>Actas Espanolas De Psiquiatria</i> , 2016, 44, 166-77.	0.1	7
47	Use of an integrated Atlas of Mental Health Care for evidence informed policy in Catalonia (Spain). <i>Epidemiology and Psychiatric Sciences</i> , 2015, 24, 512-524.	3.9	31
48	Attitudes and intended behaviour to mental disorders and associated factors in catalan population, Spain: cross-sectional population-based survey. <i>BMC Public Health</i> , 2015, 16, 127.	2.9	20
49	Bifactor analysis and construct validity of the five facet mindfulness questionnaire (FFMQ) in non-clinical Spanish samples. <i>Frontiers in Psychology</i> , 2015, 6, 404.	2.1	90
50	Is there a case for mental health promotion in the primary care setting? A systematic review. <i>Preventive Medicine</i> , 2015, 76, S5-S11.	3.4	13
51	Effectiveness of psychological and/or educational interventions to prevent the onset of episodes of depression: A systematic review of systematic reviews and meta-analyses. <i>Preventive Medicine</i> , 2015, 76, S22-S32.	3.4	41
52	Health service use and costs associated with aggressiveness or agitation and containment in adult psychiatric care: a systematic review of the evidence. <i>BMC Psychiatry</i> , 2015, 15, 35.	2.6	48
53	Cost-effectiveness of active monitoring versus antidepressants for major depression in primary health care: a 12-month non-randomized controlled trial (INFAP study). <i>BMC Psychiatry</i> , 2015, 15, 63.	2.6	11
54	Prevalence of mental disorders in non-demented elderly people in primary care. <i>International Psychogeriatrics</i> , 2015, 27, 757-768.	1.0	27

#	ARTICLE	IF	CITATIONS
55	Patients' Opinions about Knowing Their Risk for Depression and What to Do about It. The PredictD-Qualitative Study. PLoS ONE, 2014, 9, e92008.	2.5	15
56	Cost-utility of cognitive behavioral therapy versus U.S. Food and Drug Administration recommended drugs and usual care in the treatment of patients with fibromyalgia: an economic evaluation alongside a 6-month randomized controlled trial. Arthritis Research and Therapy, 2014, 16, 451.	3.5	41
57	Effectiveness of group acceptance and commitment therapy for fibromyalgia: A 6-month randomized controlled trial (EFFIGACT study). Pain, 2014, 155, 693-702.	4.2	125
58	Risk factors for the onset of panic and generalised anxiety disorders in the general adult population: A systematic review of cohort studies. Journal of Affective Disorders, 2014, 168, 337-348.	4.1	130
59	Effects of raloxifene on cognition in postmenopausal women with schizophrenia: A double-blind, randomized, placebo-controlled trial. European Neuropsychopharmacology, 2014, 24, 223-231.	0.7	66
60	Cost-effectiveness and cost-utility of a community pharmacist intervention in patients with depression (PRODEFAR study). International Journal of Integrated Care, 2014, 14, .	0.2	0
61	Validation of a Spanish version of the psychological inflexibility in pain scale (PIPS) and an evaluation of its relation with acceptance of pain and mindfulness in sample of persons with fibromyalgia. Health and Quality of Life Outcomes, 2013, 11, 62.	2.4	28
62	Preventing the onset of major depression based on the level and profile of risk of primary care attendees: protocol of a cluster randomised trial (the predictD-CCRT study). BMC Psychiatry, 2013, 13, 171.	2.6	20
63	Expectations, experiences and attitudes of patients and primary care health professionals regarding online psychotherapeutic interventions for depression: protocol for a qualitative study. BMC Psychiatry, 2013, 13, 64.	2.6	19
64	Low intensity vs. self-guided Internet-delivered psychotherapy for major depression: a multicenter, controlled, randomized study. BMC Psychiatry, 2013, 13, 21.	2.6	22
65	Quality of Life and Adverse Effects of Olanzapine Versus Risperidone Therapy in Patients with Schizophrenia. Psychiatric Quarterly, 2013, 84, 125-135.	2.1	14
66	Clinical dimensions of fibromyalgia symptoms and development of a combined index of severity: The CODI index. Quality of Life Research, 2013, 22, 153-160.	3.1	8
67	Evaluation of a pharmacist intervention on patients initiating pharmacological treatment for depression: A randomized controlled superiority trial. European Neuropsychopharmacology, 2013, 23, 1057-1066.	0.7	41
68	Epidemiology of psychiatric morbidity among migrants compared to native born population in Spain: a controlled study. General Hospital Psychiatry, 2013, 35, 93-99.	2.4	22
69	Activity monitoring in patients with depression: A systematic review. Journal of Affective Disorders, 2013, 145, 21-28.	4.1	193
70	Cost-Utility of a Psychoeducational Intervention in Fibromyalgia Patients Compared With Usual Care. Clinical Journal of Pain, 2013, 29, 702-711.	1.9	51
71	Dimensionality, Reliability, and Validity of the Revised Fibromyalgia Impact Questionnaire in Two Spanish Samples. Arthritis Care and Research, 2013, 65, 1682-1689.	3.4	29
72	An integrative cross-design synthesis approach to estimate the cost of illness: An applied case to the cost of depression in Catalonia. Journal of Mental Health, 2013, 22, 135-154.	1.9	7

#	ARTICLE	IF	CITATIONS
73	Brain Changes in Long-Term Zen Meditators Using Proton Magnetic Resonance Spectroscopy and Diffusion Tensor Imaging: A Controlled Study. PLoS ONE, 2013, 8, e58476.	2.5	40
74	Cost-Effectiveness of a Community Pharmacist Intervention in Patients with Depression: A Randomized Controlled Trial (PRODEFAR Study). PLoS ONE, 2013, 8, e70588.	2.5	31
75	Psychotropic prescribing in Catalonia: results from an epidemiological study. Family Practice, 2012, 29, 154-162.	1.9	21
76	Adjustment disorders in primary care: prevalence, recognition and use of services. British Journal of Psychiatry, 2012, 201, 137-142.	2.8	73
77	MADRID: A major depression remote intelligent monitor. , 2012, , .		2
78	Perceived injustice in fibromyalgia: Psychometric characteristics of the Injustice Experience Questionnaire and relationship with pain catastrophising and pain acceptance. Journal of Psychosomatic Research, 2012, 73, 86-91.	2.6	105
79	Antipsychotic polypharmacy in a regional health service: a population-based study. BMC Psychiatry, 2012, 12, 42.	2.6	32
80	Recognition of anxiety disorders by the general practitioner: results from the DASMAP Study. General Hospital Psychiatry, 2012, 34, 227-233.	2.4	30
81	Brain dysfunction in fibromyalgia and somatization disorder using proton magnetic resonance spectroscopy: a controlled study. Acta Psychiatrica Scandinavica, 2012, 126, 115-125.	4.5	75
82	Effectiveness of Pharmacist Care in the Improvement of Adherence to Antidepressants: A Systematic Review and Meta-Analysis. Annals of Pharmacotherapy, 2011, 45, 39-48.	1.9	75
83	Effectiveness of cognitive behaviour therapy for the treatment of catastrophisation in patients with fibromyalgia: a randomised controlled trial. Arthritis Research and Therapy, 2011, 13, R173.	3.5	53
84	Effectiveness of a Psychoeducational Treatment Program Implemented in General Practice for Fibromyalgia Patients. Clinical Journal of Pain, 2011, 27, 383-391.	1.9	60
85	Relationship between behavioural coping strategies and acceptance in patients with fibromyalgia syndrome: Elucidating targets of interventions. BMC Musculoskeletal Disorders, 2011, 12, 143.	1.9	42
86	Factors associated with use of community mental health services by schizophrenia patients using multilevel analysis. BMC Health Services Research, 2011, 11, 257.	2.2	18
87	Raloxifene as an Adjunctive Treatment for Postmenopausal Women With Schizophrenia. Journal of Clinical Psychiatry, 2011, 72, 1552-1557.	2.2	88
88	Prevalence of mental disorders in primary care: results from the diagnosis and treatment of mental disorders in primary care study (DASMAP). Social Psychiatry and Psychiatric Epidemiology, 2010, 45, 201-210.	3.1	137
89	Utility of the twelve-item World Health Organization Disability Assessment Schedule II (WHO-DAS II) for discriminating depression "caseness" and severity in Spanish primary care patients. Quality of Life Research, 2010, 19, 97-101.	3.1	30
90	The 12-item World Health Organization Disability Assessment Schedule II (WHO-DAS II): a nonparametric item response analysis. BMC Medical Research Methodology, 2010, 10, 45.	3.1	67

#	ARTICLE	IF	CITATIONS
91	Validation of the Spanish version of the Chronic Pain Acceptance Questionnaire (CPAQ) for the assessment of acceptance in fibromyalgia. <i>Health and Quality of Life Outcomes</i> , 2010, 8, 37.	2.4	62
92	Is major depression adequately diagnosed and treated by general practitioners? Results from an epidemiological study. <i>General Hospital Psychiatry</i> , 2010, 32, 201-209.	2.4	81
93	Comorbidity between common mental disorders and chronic somatic diseases in primary care patients. <i>General Hospital Psychiatry</i> , 2010, 32, 240-245.	2.4	67
94	Psychometric properties of the twelve item World Health Organization Disability Assessment Schedule II (WHO-DAS II) in Spanish primary care patients with a first major depressive episode. <i>Journal of Affective Disorders</i> , 2010, 121, 52-58.	4.1	78
95	Factor structure, internal consistency and construct validity of the Sheehan Disability Scale in a Spanish primary care sample. <i>Journal of Evaluation in Clinical Practice</i> , 2010, 16, 895-901.	1.8	54
96	Frequent attendance in primary care: comparison and implications of different definitions. <i>British Journal of General Practice</i> , 2010, 60, e49-e55.	1.4	45
97	Perfil psicológico diferencial entre dolor torácico de causa no cardíaca y enfermedad coronaria: un estudio controlado. <i>Revista Española De Cardiología</i> , 2010, 63, 357-361.	1.2	5
98	Community pharmacist intervention in depressed primary care patients (PRODEFAR study): randomized controlled trial protocol. <i>BMC Public Health</i> , 2009, 9, 284.	2.9	21
99	Impact of mental disorders and chronic physical conditions in health-related quality of life among primary care patients: results from an epidemiological study. <i>Quality of Life Research</i> , 2009, 18, 1011-1018.	3.1	21
100	Effectiveness of the psychological and pharmacological treatment of catastrophization in patients with fibromyalgia: a randomized controlled trial. <i>Trials</i> , 2009, 10, 24.	1.6	23
101	Fluoxetine and imipramine: are there differences in cost-utility for depression in primary care?. <i>Journal of Evaluation in Clinical Practice</i> , 2009, 15, 195-203.	1.8	9
102	Evaluación telefónica de los sistemas de activación e inhibición conductual. <i>Revista De Psiquiatría Y Salud Mental</i> , 2009, 2, 66-71.	1.8	6
103	Evaluation of behavioural activation and inhibition systems by telephone. <i>Revista De Psiquiatría Y Salud Mental (English Edition)</i> , 2009, 2, 66-71.	0.3	1
104	Brain dysfunction behind functional symptoms: neuroimaging and somatoform, conversive, and dissociative disorders. <i>Current Opinion in Psychiatry</i> , 2009, 22, 224-231.	6.3	36
105	Effectiveness of a psycho-educational program for improving quality of life of fibromyalgia patients. <i>BMC Musculoskeletal Disorders</i> , 2008, 9, .	1.9	5
106	Adequacy of Antidepressant Treatment in Spanish Primary Care: A Naturalistic Six-Month Follow-Up Study. <i>Psychiatric Services</i> , 2008, 59, 78-83.	2.0	27
107	Adequacy of Antidepressant Treatment in Spanish Primary Care: A Naturalistic Six-Month Follow-Up Study. <i>Psychiatric Services</i> , 2008, 59, 78-83.	2.0	9
108	Selective serotonin reuptake inhibitors: are there differences in effectiveness in primary care?. <i>Primary Care and Community Psychiatry</i> , 2007, 11, 113-120.	0.4	3

#	ARTICLE	IF	CITATIONS
109	Cost?utility of selective serotonin reuptake inhibitors for depression in primary care in Catalonia. Acta Psychiatrica Scandinavica, 2006, 114, 39-47.	4.5	21
110	Effectiveness and cost-effectiveness of antidepressant treatment in primary health care: A six-month randomised study comparing fluoxetine to imipramine. Journal of Affective Disorders, 2006, 91, 153-163.	4.1	18
111	Gender differences in response to antidepressant treatment prescribed in primary care. Does menopause make a difference?. Journal of Affective Disorders, 2006, 93, 53-60.	4.1	79
112	Assessing depression in primary care with the PHQ-9: Can it be carried out over the telephone?. Journal of General Internal Medicine, 2005, 20, 738-742.	2.6	298
113	¿CÃ³mo derivamos a salud mental desde atenciÃ³n primaria?. Atencion Primaria, 2003, 32, 524-530.	1.4	2
114	Intellectual Disability, Autism Spectrum Disorders, Psychiatric Comorbidities and Their Relationship with Challenging Behavior. Journal of Mental Health Research in Intellectual Disabilities, 0, , 1-18.	2.0	4