

Cinzia Rinaldo

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/538217/publications.pdf>

Version: 2024-02-01

38
papers

2,051
citations

236925

25
h-index

315739

38
g-index

41
all docs

41
docs citations

41
times ranked

2883
citing authors

#	ARTICLE	IF	CITATIONS
1	HIPK2 Is Required for Midbody Remnant Removal Through Autophagy-Mediated Degradation. <i>Frontiers in Cell and Developmental Biology</i> , 2020, 8, 572094.	3.7	7
2	An Alternative Splice Variant of HIPK2 with Intron Retention Contributes to Cytokinesis. <i>Cells</i> , 2020, 9, 484.	4.1	13
3	Spastin recovery in hereditary spastic paraplegia by preventing neddylation-dependent degradation. <i>Life Science Alliance</i> , 2020, 3, e202000799.	2.8	19
4	HIPK2 Phosphorylates the Microtubule-Severing Enzyme Spastin at S268 for Abscission. <i>Cells</i> , 2019, 8, 684.	4.1	31
5	p53 mitotic centrosome localization preserves centrosome integrity and works as sensor for the mitotic surveillance pathway. <i>Cell Death and Disease</i> , 2019, 10, 850.	6.3	26
6	Extrachromosomal Histone H2B Contributes to the Formation of the Abscission Site for Cell Division. <i>Cells</i> , 2019, 8, 1391.	4.1	4
7	ESCRT-III is necessary for the integrity of the nuclear envelope in micronuclei but is aberrant at ruptured micronuclear envelopes generating damage. <i>Oncogenesis</i> , 2019, 8, 29.	4.9	57
8	HIPK2 and extrachromosomal histone H2B are separately recruited by Aurora-B for cytokinesis. <i>Oncogene</i> , 2018, 37, 3562-3574.	5.9	15
9	Effects of Y361 auto-phosphorylation on structural plasticity of the HIPK2 kinase domain. <i>Protein Science</i> , 2018, 27, 725-737.	7.6	4
10	CDKL5 localizes at the centrosome and midbody and is required for faithful cell division. <i>Scientific Reports</i> , 2017, 7, 6228.	3.3	27
11	<i>Hmga1</i> null mouse embryonic fibroblasts display downregulation of spindle assembly checkpoint gene expression associated to nuclear and karyotypic abnormalities. <i>Cell Cycle</i> , 2016, 15, 812-818.	2.6	9
12	HIPK2 deficiency causes chromosomal instability by cytokinesis failure and increases tumorigenicity. <i>Oncotarget</i> , 2015, 6, 10320-10334.	1.8	30
13	Deregulation of HMGA1 expression induces chromosome instability through regulation of spindle assembly checkpoint genes. <i>Oncotarget</i> , 2015, 6, 17342-17353.	1.8	27
14	Co-targeting of Bcl-2 and mTOR pathway triggers synergistic apoptosis in BH3 mimetics resistant acute lymphoblastic leukemia. <i>Oncotarget</i> , 2015, 6, 32089-32103.	1.8	36
15	HIPK2 sustains apoptotic response by phosphorylating Che-1/AATF and promoting its degradation. <i>Cell Death and Disease</i> , 2014, 5, e1414-e1414.	6.3	11
16	Therapeutic targeting of Chk1 in NSCLC stem cells during chemotherapy. <i>Cell Death and Differentiation</i> , 2012, 19, 768-778.	11.2	157
17	HIPK2 Controls Cytokinesis and Prevents Tetraploidization by Phosphorylating Histone H2B at the Midbody. <i>Molecular Cell</i> , 2012, 47, 87-98.	9.7	58
18	Updates on HIPK2: a resourceful oncosuppressor for clearing cancer. <i>Journal of Experimental and Clinical Cancer Research</i> , 2012, 31, 63.	8.6	81

#	ARTICLE	IF	CITATIONS
19	HIPK2 phosphorylates p53 and promotes its degradation in response to DNA damage. <i>Oncogene</i> , 2011, 30, 4802-4813.	5.9	57
20	Homeodomain-interacting Protein Kinase-2 Stabilizes p27kip1 by Its Phosphorylation at Serine 10 and Contributes to Cell Motility. <i>Journal of Biological Chemistry</i> , 2011, 286, 29005-29013.	3.4	9
21	MYCN Sensitizes Human Neuroblastoma to Apoptosis by HIPK2 Activation through a DNA Damage Response. <i>Molecular Cancer Research</i> , 2011, 9, 67-77.	3.4	30
22	The Loss of the p53 Activator HIPK2 Is Responsible for Galectin-3 Overexpression in Well Differentiated Thyroid Carcinomas. <i>PLoS ONE</i> , 2011, 6, e20665.	2.5	54
23	Pre Clinical mTOR-Inhibition of Acute Lymphoblastic Leukemia Cells Synergizes with Pro-Apoptotic Target Therapy Through Mcl-1 Down-Regulation. <i>Blood</i> , 2011, 118, 3581-3581.	1.4	0
24	HIPK2 Regulation by MDM2 Determines Tumor Cell Response to the p53-Reactivating Drugs Nutlin-3 and RITA. <i>Cancer Research</i> , 2009, 69, 6241-6248.	0.9	49
25	Sgk1 activates MDM2-dependent p53 degradation and affects cell proliferation, survival, and differentiation. <i>Journal of Molecular Medicine</i> , 2009, 87, 1221-1239.	3.9	88
26	p53 is stimulated by gain-of-function p53 mutations and modulates chemoresistance in anaplastic thyroid carcinomas. <i>Journal of Pathology</i> , 2009, 218, 66-75.	4.5	33
27	HIPK2 is involved in cell proliferation and its suppression promotes growth arrest independently of DNA damage. <i>Cell Proliferation</i> , 2009, 42, 373-384.	5.3	33
28	MDM4 (MDMX) localizes at the mitochondria and facilitates the p53-mediated intrinsic-apoptotic pathway. <i>EMBO Journal</i> , 2009, 28, 1926-1939.	7.8	75
29	HIPKs: Jack of all trades in basic nuclear activities. <i>Biochimica Et Biophysica Acta - Molecular Cell Research</i> , 2008, 1783, 2124-2129.	4.1	58
30	MDM2-Regulated Degradation of HIPK2 Prevents p53Ser46 Phosphorylation and DNA Damage-Induced Apoptosis. <i>Molecular Cell</i> , 2007, 25, 739-750.	9.7	161
31	HIPK2: a multitasking partner for transcription factors in DNA damage response and development This paper is one of a selection of papers published in this Special Issue, entitled 28th International West Coast Chromatin and Chromosome Conference, and has undergone the Journal's usual peer review process. <i>Biochemistry and Cell Biology</i> , 2007, 85, 411-418.	2.0	115
32	High-mobility group A1 inhibits p53 by cytoplasmic relocalization of its proapoptotic activator HIPK2. <i>Journal of Clinical Investigation</i> , 2007, 117, 693-702.	8.2	88
33	High Mobility Group A1 (HMGA1) proteins interact with p53 and inhibit its apoptotic activity. <i>Cell Death and Differentiation</i> , 2006, 13, 1554-1563.	11.2	65
34	Mutant p53 gain of function: reduction of tumor malignancy of human cancer cell lines through abrogation of mutant p53 expression. <i>Oncogene</i> , 2006, 25, 304-309.	5.9	188
35	Repression of the Antiapoptotic Molecule Galectin-3 by Homeodomain-Interacting Protein Kinase 2-Activated p53 Is Required for p53-Induced Apoptosis. <i>Molecular and Cellular Biology</i> , 2006, 26, 4746-4757.	2.3	93
36	Homeodomain-interacting protein kinase-2 activity and p53 phosphorylation are critical events for cisplatin-mediated apoptosis. <i>Experimental Cell Research</i> , 2004, 293, 311-320.	2.6	99

#	ARTICLE	IF	CITATIONS
37	Roles for <i>Caenorhabditis elegans rad-51</i> in Meiosis and in Resistance to Ionizing Radiation During Development. <i>Genetics</i> , 2002, 160, 471-479.	2.9	119
38	The <i>Caenorhabditis elegans</i> RAD51 homolog is transcribed into two alternative mRNAs potentially encoding proteins of different sizes. <i>Molecular Genetics and Genomics</i> , 1998, 260, 289-294.	2.4	24