

# San-Gang Wu

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/538474/publications.pdf>

Version: 2024-02-01

143  
papers

2,335  
citations

304743

22  
h-index

330143

37  
g-index

149  
all docs

149  
docs citations

149  
times ranked

3681  
citing authors

#	ARTICLE	IF	CITATIONS
1	Distribution, Chemotherapy Use, and Outcome of the 21-Gene Recurrence Score Between Chinese and White breast Cancer in the United States. <i>Clinical Breast Cancer</i> , 2022, 22, 279-287.	2.4	3
2	OUP accepted manuscript. <i>BJS Open</i> , 2022, 6, .	1.7	0
3	Olanzapine 5â€‰mg for Nausea and Vomiting in Patients with Nasopharyngeal Carcinoma Receiving Cisplatin-Based Concurrent Chemoradiotherapy. <i>Journal of Oncology</i> , 2022, 2022, 1-7.	1.3	1
4	Prognostic significance of the skeletal muscle index and systemic inflammatory index in patients with lymph node-positive breast cancer after radical mastectomy. <i>BMC Cancer</i> , 2022, 22, 234.	2.6	10
5	The prognostic effect of residual tumor for advanced epithelial ovarian cancer treated with neoadjuvant chemotherapy or primary debulking surgery. <i>Cancer Medicine</i> , 2022, , .	2.8	5
6	Local treatment improves survival in patients with stage IVB cervical cancer. <i>Gynecologic Oncology</i> , 2022, 165, 538-545.	1.4	5
7	The Predictive Effect of the 8th AJCC Pathological Prognostic Staging on the Benefit of Postmastectomy Radiotherapy in N2/N3 Breast Cancer. <i>Breast Cancer: Targets and Therapy</i> , 2022, Volume 14, 133-144.	1.8	1
8	The prognostic and predictive value of the 8th American Joint Committee on Cancer (AJCC) staging system among early breast cancer patients aged <50 years. <i>Gland Surgery</i> , 2021, 10, 233-241.	1.1	2
9	Additional radiotherapy to breast-conserving surgery is an optional treatment for de novo stage IV breast cancer: A population-based analysis. <i>Cancer Medicine</i> , 2021, 10, 1634-1643.	2.8	6
10	Evaluation of Subjective Sleep Disturbances in Cancer Patients: A Cross-Sectional Study in a Radiotherapy Department. <i>Frontiers in Psychiatry</i> , 2021, 12, 648896.	2.6	11
11	Identification of MEG8/miR-378d/SOBP axis as a novel regulatory network and associated with immune infiltrates in ovarian carcinoma by integrated bioinformatics analysis. <i>Cancer Medicine</i> , 2021, 10, 2924-2939.	2.8	9
12	circ-PTK2 (hsa_circ_0008305) regulates the pathogenic processes of ovarian cancer via miR-639 and FOXC1 regulatory cascade. <i>Cancer Cell International</i> , 2021, 21, 277.	4.1	9
13	Survival Outcomes and Treatment Decision by Human Papillomavirus Status Among Patients With Stage IVC Head and Neck Squamous Cell Carcinoma. <i>Frontiers in Oncology</i> , 2021, 11, 668066.	2.8	7
14	Adjuvant chemotherapy and survival outcome in node-negative breast cancer with a 21-gene recurrence score of 26â€“30. <i>Future Oncology</i> , 2021, 17, 2183-2192.	2.4	1
15	Metronomic capecitabine as adjuvant therapy in locoregionally advanced nasopharyngeal carcinoma: a multicentre, open-label, parallel-group, randomised, controlled, phase 3 trial. <i>Lancet</i> , The, 2021, 398, 303-313.	13.7	98
16	Biopsy of cervical lymph node does not impact the survival of nasopharyngeal carcinoma. <i>Cancer Medicine</i> , 2021, 10, 6687-6696.	2.8	2
17	Increased risk of cerebrovascular mortality in head and neck cancer survivors aged 65 years treated with definitive radiotherapy: a population-based cohort study. <i>Radiation Oncology</i> , 2021, 16, 185.	2.7	7
18	The patterns of distant metastasis and prognostic factors in patients with primary metastatic Ewing sarcoma of the bone. <i>Journal of Bone Oncology</i> , 2021, 30, 100385.	2.4	7

#	ARTICLE	IF	CITATIONS
19	Triple-negative breast cancer outcomes: Does AJCC 8th staging improve chemotherapy decision-making. <i>Breast</i> , 2021, 59, 117-123.	2.2	4
20	Long-Term Survival Among Histological Subtypes in Advanced Epithelial Ovarian Cancer: Population-Based Study Using the Surveillance, Epidemiology, and End Results Database. <i>JMIR Public Health and Surveillance</i> , 2021, 7, e25976.	2.6	8
21	Rare Metastasis to the Submandibular Gland in Oral Squamous Cell Carcinoma. <i>Frontiers in Oncology</i> , 2021, 11, 728230.	2.8	2
22	Should women with early breast cancer under 40 years of age have a routine 21-gene recurrence score testing: A SEER database study. <i>Breast</i> , 2020, 49, 233-241.	2.2	8
23	Integration the biologic factors into the staging of breast cancer patients with ipsilateral supraclavicular lymph node metastasis. <i>Journal of Cancer</i> , 2020, 11, 6834-6840.	2.5	0
24	The effect of postmastectomy radiotherapy in node-positive triple-negative breast cancer. <i>BMC Cancer</i> , 2020, 20, 1146.	2.6	7
25	Cognitive dysfunction in patients with nasopharyngeal carcinoma after induction chemotherapy. <i>Oral Oncology</i> , 2020, 111, 104921.	1.5	6
26	Incorporation of biologic factors for the staging of de novo stage IV breast cancer. <i>Npj Breast Cancer</i> , 2020, 6, 43.	5.2	7
27	Prognostic and Predictive Value of the American Joint Committee on Cancer Pathological Prognostic Staging System in Nodal Micrometastatic Breast Cancer. <i>Frontiers in Oncology</i> , 2020, 10, 570175.	2.8	1
28	Aggressive Local Treatment Improves Survival in Stage IV Breast Cancer With Synchronous Metastasis. <i>Frontiers in Oncology</i> , 2020, 10, 522580.	2.8	7
29	Real-world impact of postmastectomy radiotherapy in T1â€“2 breast cancer with one to three positive lymph nodes. <i>Annals of Translational Medicine</i> , 2020, 8, 489-489.	1.7	6
30	Evaluation of the 8th edition of the American joint committee on cancerâ€™s pathological staging system in prognosis assessment and treatment decision making for stage T1-2N1 breast cancer after mastectomy. <i>Breast</i> , 2020, 51, 2-10.	2.2	13
31	Prognostic validation and therapeutic decisionâ€™making of the AJCC eighth pathological prognostic staging for T3N0 breast cancer after mastectomy. <i>Clinical and Translational Medicine</i> , 2020, 10, 125-136.	4.0	11
32	Staging for Breast Cancer With Internal Mammary Lymph Nodes Metastasis: Utility of Incorporating Biologic Factors. <i>Frontiers in Oncology</i> , 2020, 10, 584009.	2.8	3
33	Chemotherapy and 21-gene recurrence score testing for older breast cancer patients: A competing-risks analysis. <i>Breast</i> , 2020, 54, 319-327.	2.2	8
34	Thyroid-like low-grade nasopharyngeal papillary adenocarcinoma: a case report and literature review. <i>Translational Cancer Research</i> , 2020, 9, 4457-4463.	1.0	3
35	The longitudinal risk of mortality between invasive ductal carcinoma and metaplastic breast carcinoma. <i>Scientific Reports</i> , 2020, 10, 22070.	3.3	8
36	Histological Tumor Type is Associated with One-Year Cause-Specific Survival in Women with Stage IIIâ€“IV Epithelial Ovarian Cancer: A Surveillance, Epidemiology, and End Results (SEER) Database Population Study, 2004â€“2014. <i>Medical Science Monitor</i> , 2020, 26, e920531.	1.1	6

#	ARTICLE	IF	CITATIONS
37	Validation of the 8th edition of the American Joint Committee on Cancer Pathological Prognostic Staging for young breast cancer patients. <i>Aging</i> , 2020, 12, 7549-7560.	3.1	4
38	Prognostic validation and treatment decision making of the 8th edition of the American Joint Committee on Cancer pathological staging system for elderly women with early-stage breast cancer. <i>Aging</i> , 2020, 12, 15077-15090.	3.1	2
39	Analysis of the Trends in Publications on Clinical Cancer Research in Mainland China from the Surveillance, Epidemiology, and End Results (SEER) Database: Bibliometric Study. <i>JMIR Medical Informatics</i> , 2020, 8, e21931.	2.6	4
40	The 1-year mortality after radiotherapy for nasopharyngeal carcinoma: a population-based analysis. <i>Future Oncology</i> , 2019, 15, 3357-3365.	2.4	2
41	Lung Large Cell Neuroendocrine Carcinoma: An Analysis of Patients from the Surveillance, Epidemiology, and End-Results (SEER) Database. <i>Medical Science Monitor</i> , 2019, 25, 3636-3646.	1.1	21
42	Progesterone receptor status and tumor grade predict the 21-gene recurrence score of invasive lobular breast cancer. <i>Biomarkers in Medicine</i> , 2019, 13, 1005-1012.	1.4	8
43	Omission of adjuvant radiotherapy following breast-conserving surgery for elderly women with early-stage pure mucinous breast carcinoma. <i>Radiation Oncology</i> , 2019, 14, 190.	2.7	9
44	Real-World Impact of Survival by Period of Diagnosis in Epithelial Ovarian Cancer Between 1990 and 2014. <i>Frontiers in Oncology</i> , 2019, 9, 639.	2.8	31
45	The Effect of Post-mastectomy Radiotherapy in Patients With Metaplastic Breast Cancer: An Analysis of SEER Database. <i>Frontiers in Oncology</i> , 2019, 9, 747.	2.8	11
46	Lymph Node Status and Outcomes for Nasopharyngeal Carcinoma According to Histological Subtypes: A SEER Population-Based Retrospective Analysis. <i>Advances in Therapy</i> , 2019, 36, 3123-3133.	2.9	18
47	The Role of Axillary Lymph Node Dissection in Tubular Carcinoma of the Breast: A Population Database Study. <i>Medical Science Monitor</i> , 2019, 25, 880-887.	1.1	5
48	21-Gene Recurrence Score Assay Could Not Predict Benefit of Post-mastectomy Radiotherapy in T1-2 N1mic ER-Positive HER2-Negative Breast Cancer. <i>Frontiers in Oncology</i> , 2019, 9, 270.	2.8	8
49	Noninferior Outcome After Breast-Conserving Treatment Compared to Mastectomy in Breast Cancer Patients With Four or More Positive Lymph Nodes. <i>Frontiers in Oncology</i> , 2019, 9, 143.	2.8	9
50	The 21-gene recurrence score and effects of adjuvant radiotherapy after breast conserving surgery in early-stage breast cancer. <i>Future Oncology</i> , 2019, 15, 1629-1639.	2.4	8
51	<p>&lt;p>Effect of 21-gene recurrence score in decision-making for surgery in early stage breast cancer&lt;/p>. <i>OncoTargets and Therapy</i> , 2019, Volume 12, 2071-2078.	2.0	2
52	21-Gene Recurrence Score Assay and Outcomes of Adjuvant Radiotherapy in Elderly Women With Early-Stage Breast Cancer After Breast-Conserving Surgery. <i>Frontiers in Oncology</i> , 2019, 9, 1.	2.8	139
53	The effect of histological subtypes on survival outcome in nasopharyngeal carcinoma after extensive follow up. <i>Annals of Translational Medicine</i> , 2019, 7, 768-768.	1.7	22
54	<p>Development and validation of a novel diagnostic model for assessing lung cancer metastasis in a Chinese population based on multicenter real-world data</p>. <i>Cancer Management and Research</i> , 2019, Volume 11, 9213-9223.	1.9	2

#	ARTICLE	IF	CITATIONS
55	Inflammatory breast cancer outcomes by breast cancer subtype: a population-based study. <i>Future Oncology</i> , 2019, 15, 507-516.	2.4	16
56	21-gene recurrence score and adjuvant chemotherapy decisions in patients with invasive lobular breast cancer. <i>Biomarkers in Medicine</i> , 2019, 13, 83-93.	1.4	19
57	Impact of the 21-gene recurrence score assay on chemotherapy decision making and outcomes for breast cancer patients with four or more positive lymph nodes. <i>Annals of Translational Medicine</i> , 2019, 7, 446-446.	1.7	5
58	KIF11 Functions as an Oncogene and Is Associated with Poor Outcomes from Breast Cancer. <i>Cancer Research and Treatment</i> , 2019, 51, 1207-1221.	3.0	47
59	Impact of 21-Gene Recurrence Score on Chemotherapy Decision in Invasive Ductal Carcinoma of Breast with Nodal Micrometastases. <i>Cancer Research and Treatment</i> , 2019, 51, 1437-1448.	3.0	4
60	Bioinformatics-Based Discovery of CKLF-Like MARVEL Transmembrane Member 5 as a Novel Biomarker for Breast Cancer. <i>Frontiers in Cell and Developmental Biology</i> , 2019, 7, 361.	3.7	4
61	Development and validation of a novel diagnostic nomogram model based on tumor markers for assessing cancer risk of pulmonary lesions: A multicenter study in Chinese population. <i>Cancer Letters</i> , 2018, 420, 236-241.	7.2	16
62	The effect of lymphadenectomy in advanced ovarian cancer according to residual tumor status: A population-based study. <i>International Journal of Surgery</i> , 2018, 52, 11-15.	2.7	19
63	Comparison of the effects of local treatment strategies in non-metastatic Ewing sarcoma of bone. <i>Expert Review of Anticancer Therapy</i> , 2018, 18, 501-506.	2.4	4
64	Survival in signet ring cell carcinoma varies based on primary tumor location: a Surveillance, Epidemiology, and End Results database analysis. <i>Expert Review of Gastroenterology and Hepatology</i> , 2018, 12, 209-214.	3.0	50
65	The effects of postoperative radiotherapy on survival outcomes in patients under 65 with estrogen receptor positive tubular breast carcinoma. <i>Radiation Oncology</i> , 2018, 13, 226.	2.7	7
66	The Distribution and Outcomes of the 21-Gene Recurrence Score in T1-T2N0 Estrogen Receptor-Positive Breast Cancer With Different Histologic Subtypes. <i>Frontiers in Genetics</i> , 2018, 9, 638.	2.3	23
67	The Effect of Histological Subtypes on Outcomes of Stage IV Epithelial Ovarian Cancer. <i>Frontiers in Oncology</i> , 2018, 8, 577.	2.8	25
68	Clinicopathologic characteristics and clinical outcomes of pure type and mixed type of tubular carcinoma of the breast: a single-institution cohort study. <i>Cancer Management and Research</i> , 2018, Volume 10, 4509-4515.	1.9	6
69	The Clinicopathological Features and Survival Outcomes of Different Histological Subtypes in Triple-negative Breast Cancer. <i>Journal of Cancer</i> , 2018, 9, 296-303.	2.5	60
70	Prognostic value of ductal carcinoma in situ component in invasive ductal carcinoma of the breast: a Surveillance, Epidemiology, and End Results database analysis. <i>Cancer Management and Research</i> , 2018, Volume 10, 527-534.	1.9	5
71	Trends and Outcomes of Sentinel Lymph Node Biopsy in Early-stage Vulvar Squamous Cell Carcinoma: A Population-based Study. <i>Journal of Cancer</i> , 2018, 9, 1951-1957.	2.5	7
72	The prognosis and effects of local treatment strategies for orbital embryonal rhabdomyosarcoma: a population-based study. <i>Cancer Management and Research</i> , 2018, Volume 10, 1727-1734.	1.9	3

#	ARTICLE	IF	CITATIONS
73	The Effect of Marital Status on Nasopharyngeal Carcinoma Survival: A Surveillance, Epidemiology and End Results Study. <i>Journal of Cancer</i> , 2018, 9, 1870-1876.	2.5	23
74	Long-term survival effect of the interval between mastectomy and radiotherapy in locally advanced breast cancer. <i>Cancer Management and Research</i> , 2018, Volume 10, 2047-2054.	1.9	11
75	Comparison of survival outcomes of locally advanced breast cancer patients receiving post-mastectomy radiotherapy with and without immediate breast reconstruction: a population-based analysis. <i>Cancer Management and Research</i> , 2018, Volume 10, 1993-2002.	1.9	9
76	Downregulation of hsa_circ_0011946 suppresses the migration and invasion of the breast cancer cell line MCF-7 by targeting RFC3. <i>Cancer Management and Research</i> , 2018, Volume 10, 535-544.	1.9	75
77	Prognostic Value of the Number of Removed Lymph Nodes in Vulvar Squamous Cell Carcinoma Patients With Node-Positive Disease: A Population-Based Study. <i>Frontiers in Oncology</i> , 2018, 8, 184.	2.8	1
78	Omission of Postoperative Radiotherapy in Women Aged 65 Years or Older With Tubular Carcinoma of the Breast After Breast-Conserving Surgery. <i>Frontiers in Oncology</i> , 2018, 8, 190.	2.8	9
79	Tubular carcinomas of the breast: an epidemiologic study. <i>Future Oncology</i> , 2018, 14, 3037-3047.	2.4	4
80	Widowed status increases the risk of death in vulvar cancer. <i>Future Oncology</i> , 2018, 14, 2589-2598.	2.4	8
81	Patterns of Distant Metastasis Between Histological Types in Esophageal Cancer. <i>Frontiers in Oncology</i> , 2018, 8, 302.	2.8	52
82	Survival benefits with the addition of adjuvant hysterectomy to radiochemotherapy for treatment of stage I adenocarcinoma of the uterine cervix. <i>Journal of Surgical Oncology</i> , 2018, 118, 574-580.	1.7	5
83	Lymph node ratio has prognostic value related to the number of positive lymph nodes in patients with vulvar cancer. <i>Future Oncology</i> , 2018, 14, 2343-2351.	2.4	3
84	Clinical Features of Brain Metastases in Small Cell Lung Cancer: an Implication for Hippocampal Sparing Whole Brain Radiation Therapy. <i>Translational Oncology</i> , 2017, 10, 54-58.	3.7	14
85	Up-Regulation of RFC3 Promotes Triple Negative Breast Cancer Metastasis and is Associated With Poor Prognosis Via EMT. <i>Translational Oncology</i> , 2017, 10, 1-9.	3.7	46
86	The prognostic value of histologic subtype in node-positive early-stage cervical cancer after hysterectomy and adjuvant radiotherapy. <i>International Journal of Surgery</i> , 2017, 44, 1-6.	2.7	13
87	Men and women show similar survival outcome in stage IV breast cancer. <i>Breast</i> , 2017, 34, 115-121.	2.2	6
88	The effect of local treatment modalities in patients with early-stage adenocarcinoma of the uterine cervix: A population-based analysis. <i>International Journal of Surgery</i> , 2017, 41, 16-22.	2.7	14
89	Adjuvant radiation therapy and survival for adenoid cystic carcinoma of the breast. <i>Breast</i> , 2017, 31, 214-218.	2.2	21
90	Multimodal treatment including hysterectomy improves survival in patients with locally advanced cervical cancer: A population-based, propensity score-matched analysis. <i>International Journal of Surgery</i> , 2017, 48, 122-127.	2.7	7

#	ARTICLE	IF	CITATIONS
91	Demographic and clinicopathological characteristics of nasopharyngeal carcinoma and survival outcomes according to age at diagnosis: A population-based analysis. <i>Oral Oncology</i> , 2017, 73, 83-87.	1.5	40
92	Preoperative radiotherapy improves survival in rectal signet-ring cell carcinoma-a population-based study. <i>Radiation Oncology</i> , 2017, 12, 141.	2.7	12
93	The effect of distant metastases sites on survival in de novo stage-IV breast cancer: A SEER database analysis. <i>Tumor Biology</i> , 2017, 39, 101042831770508.	1.8	56
94	Therapeutic role of axillary lymph node dissection in patients with stage IV breast cancer: a population-based analysis. <i>Journal of Cancer Research and Clinical Oncology</i> , 2017, 143, 467-474.	2.5	7
95	Comparison of clinical outcomes of squamous cell carcinoma, adenocarcinoma, and adenosquamous carcinoma of the uterine cervix after definitive radiotherapy: a population-based analysis. <i>Journal of Cancer Research and Clinical Oncology</i> , 2017, 143, 115-122.	2.5	59
96	Early-stage node negative cervical adenocarcinoma and squamous cell carcinoma show similar survival outcomes after hysterectomy: a population-based study. <i>Journal of Gynecologic Oncology</i> , 2017, 28, e81.	2.2	10
97	Sites of metastasis and overall survival in esophageal cancer: a population-based study. <i>Cancer Management and Research</i> , 2017, Volume 9, 781-788.	1.9	68
98	Postoperative radiotherapy for invasive micropapillary carcinoma of the breast: an analysis of Surveillance, Epidemiology, and End Results database. <i>Cancer Management and Research</i> , 2017, Volume 9, 453-459.	1.9	14
99	The impact of examined lymph node count on survival in squamous cell carcinoma and adenocarcinoma of the uterine cervix. <i>Cancer Management and Research</i> , 2017, Volume 9, 315-322.	1.9	19
100	Comparison of survival outcomes between radical hysterectomy and definitive radiochemotherapy in stage IB1 and IIA1 cervical cancer. <i>Cancer Management and Research</i> , 2017, Volume 9, 813-819.	1.9	13
101	Comparable Survival between Additional Radiotherapy and Local Surgery in Occult Breast Cancer after Axillary Lymph Node Dissection: A Population-based Analysis. <i>Journal of Cancer</i> , 2017, 8, 3849-3855.	2.5	13
102	The survival benefits of local surgery in stage IV breast cancer are not affected by breast cancer subtypes: a population-based analysis. <i>Oncotarget</i> , 2017, 8, 67851-67860.	1.8	10
103	Incorporation of the number of positive lymph nodes leads to better prognostic discrimination of node-positive early stage cervical cancer. <i>Oncotarget</i> , 2017, 8, 26057-26065.	1.8	12
104	Clinicopathological features of small cell carcinoma of the uterine cervix in the surveillance, epidemiology, and end results database. <i>Oncotarget</i> , 2017, 8, 40425-40433.	1.8	17
105	Progesterone receptor loss identifies hormone receptor-positive and HER2-negative breast cancer subgroups at higher risk of relapse: a retrospective cohort study. <i>OncoTargets and Therapy</i> , 2016, 9, 1707.	2.0	9
106	Use of CEA and CA15-3 to Predict Axillary Lymph Node Metastasis in Patients with Breast Cancer. <i>Journal of Cancer</i> , 2016, 7, 37-41.	2.5	23
107	Patterns of distant metastasis in Chinese women according to breast cancer subtypes. <i>Oncotarget</i> , 2016, 7, 47975-47984.	1.8	23
108	Clinical features of brain metastases in breast cancer: an implication for hippocampal-sparing whole-brain radiation therapy. <i>Therapeutics and Clinical Risk Management</i> , 2016, Volume 12, 1849-1853.	2.0	11

#	ARTICLE	IF	CITATIONS
109	Differences in esophageal cancer characteristics and survival between Chinese and Caucasian patients in the SEER database. <i>OncoTargets and Therapy</i> , 2016, Volume 9, 6435-6444.	2.0	12
110	Impact of the number of resected lymph nodes on survival after preoperative radiotherapy for esophageal cancer. <i>Oncotarget</i> , 2016, 7, 22497-22507.	1.8	14
111	Prognostic value of lymph node ratio in stage IIIC epithelial ovarian cancer with node-positive in a SEER population-based study. <i>Oncotarget</i> , 2016, 7, 7952-7959.	1.8	22
112	Surgery Combined with Radiotherapy Improved Survival in Metastatic Esophageal Cancer in a Surveillance Epidemiology and End Results Population-based Study. <i>Scientific Reports</i> , 2016, 6, 28280.	3.3	31
113	Lymph node dissection improved survival in patients with metastatic thoracic esophageal cancer: An analysis of 220 patients from the SEER database. <i>International Journal of Surgery</i> , 2016, 35, 13-18.	2.7	6
114	Patterns of Regional Lymph Node Recurrence After Radical Surgery for Thoracic Esophageal Squamous Cell Carcinoma. <i>Annals of Thoracic Surgery</i> , 2016, 101, 551-557.	1.3	22
115	Risk factors for lymph node metastasis in ovarian cancer: Implications for systematic lymphadenectomy. <i>International Journal of Surgery</i> , 2016, 29, 123-127.	2.7	20
116	The local treatment modalities in FIGO stage I-III small-cell carcinoma of the cervix are determined by disease stage and lymph node status. <i>Cancer Medicine</i> , 2016, 5, 1108-1115.	2.8	17
117	The Activation of ERK1/2 and JNK MAPK Signaling by Insulin/IGF-1 Is Responsible for the Development of Colon Cancer with Type 2 Diabetes Mellitus. <i>PLoS ONE</i> , 2016, 11, e0149822.	2.5	38
118	Effect of postoperative radiotherapy for squamous cell cancer of the breast in a surveillance epidemiology and end results population-based study. <i>Oncotarget</i> , 2016, 7, 10684-10693.	1.8	4
119	Lymph node ratio may predict the benefit of postoperative radiotherapy in node-positive cervical cancer. <i>Oncotarget</i> , 2016, 7, 29420-29428.	1.8	16
120	Number of Negative Lymph Nodes Can Predict Survival after Postmastectomy Radiotherapy According to Different Breast Cancer Subtypes. <i>Journal of Cancer</i> , 2015, 6, 261-269.	2.5	3
121	Tailoring Pelvic Lymphadenectomy for Patients with Stage IA2, IB1, and IIA1 Uterine Cervical Cancer. <i>Journal of Cancer</i> , 2015, 6, 377-381.	2.5	18
122	Prognostic Impact of ABO Blood Group on the Survival in Patients with Ovarian Cancer. <i>Journal of Cancer</i> , 2015, 6, 970-975.	2.5	17
123	Effect of blood type on survival of Chinese patients with esophageal squamous cell carcinoma. <i>OncoTargets and Therapy</i> , 2015, 8, 947.	2.0	7
124	Prognostic value of lymph node ratio in patients with small-cell carcinoma of the cervix based on data from a large national registry. <i>OncoTargets and Therapy</i> , 2015, 9, 67.	2.0	2
125	Influence of different treatment modalities on survival of patients with low-grade endometrial stromal sarcoma: A retrospective cohort study. <i>International Journal of Surgery</i> , 2015, 23, 147-151.	2.7	24
126	Prognostic Value of Different Lymph Node Staging Methods in Esophageal Squamous Cell Carcinoma After Esophagectomy. <i>Annals of Thoracic Surgery</i> , 2015, 99, 284-290.	1.3	19

#	ARTICLE	IF	CITATIONS
127	Number of negative lymph nodes is associated with disease-free survival in patients with breast cancer. <i>BMC Cancer</i> , 2015, 15, 43.	2.6	10
128	Postmastectomy Radiotherapy Improves Disease-Free Survival of High Risk of Locoregional Recurrence Breast Cancer Patients with T1-2 and 1 to 3 Positive Nodes. <i>PLoS ONE</i> , 2015, 10, e0119105.	2.5	22
129	Prognosis of patients with esophageal squamous cell carcinoma after esophagectomy using the log odds of positive lymph nodes. <i>Oncotarget</i> , 2015, 6, 36911-36922.	1.8	26
130	Distribution of metastatic disease in the brain in relation to the hippocampus: a retrospective single-center analysis of 6064 metastases in 632 patients. <i>Oncotarget</i> , 2015, 6, 44030-44036.	1.8	25
131	Using the Lymph Node Ratio to Evaluate the Prognosis of Stage II/III Breast Cancer Patients Who Received Neoadjuvant Chemotherapy and Mastectomy. <i>Cancer Research and Treatment</i> , 2015, 47, 757-764.	3.0	20
132	Number of negative lymph nodes should be considered for incorporation into staging for breast cancer. <i>American Journal of Cancer Research</i> , 2015, 5, 844-53.	1.4	20
133	Transdermal fentanyl for pain due to chemoradiotherapy-induced oral mucositis in nasopharyngeal cancer patients: evaluating efficacy, safety, and improvement in quality of life. <i>Drug Design, Development and Therapy</i> , 2014, 8, 497.	4.3	16
134	Prognostic Value of Ki-67 in Breast Cancer Patients with Positive Axillary Lymph Nodes: A Retrospective Cohort Study. <i>PLoS ONE</i> , 2014, 9, e87264.	2.5	33
135	The value of radiotherapy in breast cancer patients with isolated ipsilateral supraclavicular lymph node metastasis without distant metastases at diagnosis: a retrospective analysis of Chinese patients. <i>OncoTargets and Therapy</i> , 2014, 7, 281.	2.0	6
136	Number of negative lymph nodes can predict survival of breast cancer patients with four or more positive lymph nodes after postmastectomy radiotherapy. <i>Radiation Oncology</i> , 2014, 9, 284.	2.7	12
137	Dosimetric analysis of the brachial plexus among patients with breast cancer treated with post-mastectomy radiotherapy to the ipsilateral supraclavicular area: report of 3 cases of radiation-induced brachial plexus neuropathy. <i>Radiation Oncology</i> , 2014, 9, 292.	2.7	16
138	Serum levels of CEA and CA15-3 in different molecular subtypes and prognostic value in Chinese breast cancer. <i>Breast</i> , 2014, 23, 88-93.	2.2	90
139	Using the lymph nodal ratio to predict the risk of locoregional recurrence in lymph node-positive breast cancer patients treated with mastectomy without radiation therapy. <i>Radiation Oncology</i> , 2013, 8, 119.	2.7	14
140	Locoregional recurrence of pT3N0M0 breast cancer after mastectomy is not higher than that of pT1-2N0M0: An analysis for radiotherapy. <i>Cancer Science</i> , 2013, 104, 599-603.	3.9	2
141	Prognostic Value of Metastatic Axillary Lymph Node Ratio for Chinese Breast Cancer Patients. <i>PLoS ONE</i> , 2013, 8, e61410.	2.5	34
142	Predictive value of breast cancer molecular subtypes in Chinese patients with four or more positive nodes after postmastectomy radiotherapy. <i>Breast</i> , 2012, 21, 657-661.	2.2	33
143	Ovarian Ablation Using Goserelin Improves Survival of Premenopausal Patients with Stage II/III Hormone Receptor-Positive Breast Cancer without Chemotherapy-Induced Amenorrhea. <i>Cancer Research and Treatment</i> , 1970, 47, 55-63.	3.0	8