

Riobaldo M R Cintra

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/547388/publications.pdf>

Version: 2024-02-01

15
papers

199
citations

1163117

8
h-index

1058476

14
g-index

17
all docs

17
docs citations

17
times ranked

365
citing authors

#	ARTICLE	IF	CITATIONS
1	Rebound inflammatory response during the acute phase of myocardial infarction after simvastatin withdrawal. <i>Atherosclerosis</i> , 2009, 207, 191-194.	0.8	61
2	Dapagliflozin effect on endothelial dysfunction in diabetic patients with atherosclerotic disease: a randomized active-controlled trial. <i>Cardiovascular Diabetology</i> , 2021, 20, 74.	6.8	44
3	Glycosylated hemoglobin is associated with decreased endothelial function, high inflammatory response, and adverse clinical outcome in non-diabetic STEMI patients. <i>Atherosclerosis</i> , 2015, 243, 124-130.	0.8	17
4	Validation of surrogate indexes of insulin sensitivity in acute phase of myocardial infarction based on euglycemic-hyperinsulinemic clamp. <i>American Journal of Physiology - Endocrinology and Metabolism</i> , 2014, 306, E399-E403.	3.5	15
5	Inhibition of the sodium-glucose co-transporter 2 in the elderly: clinical and mechanistic insights into safety and efficacy. <i>Revista Da Associação Médica Brasileira</i> , 2019, 65, 70-86.	0.7	15
6	Central role of obesity in endothelial cell dysfunction and cardiovascular risk. <i>Revista Da Associação Médica Brasileira</i> , 2019, 65, 87-97.	0.7	10
7	High plasma HDL-C attenuates stress hyperglycemia during acute phase of myocardial infarction. <i>Atherosclerosis</i> , 2012, 220, 231-236.	0.8	9
8	Assessment of dapagliflozin effect on diabetic endothelial dysfunction of brachial artery (ADDENDA-BHS2 trial): rationale, design, and baseline characteristics of a randomized controlled trial. <i>Diabetology and Metabolic Syndrome</i> , 2019, 11, 62.	2.7	9
9	Dapagliflozin increases retinal thickness in type 2 diabetic patients as compared with glibenclamide: A randomized controlled trial. <i>Diabetes and Metabolism</i> , 2021, 47, 101280.	2.9	6
10	High-density lipoprotein levels are strongly associated with the recovery rate of insulin sensitivity during the acute phase of myocardial infarction: A study by euglycemic hyperinsulinemic clamp. <i>Journal of Clinical Lipidology</i> , 2013, 7, 24-28.	1.5	4
11	Adiponectin concentration data improve the estimation of atherosclerotic risk in normal and in overweight subjects. <i>Clinical Endocrinology</i> , 2018, 88, 388-396.	2.4	4
12	TCF7L2 polymorphism is associated with low nitric oxide release, endothelial dysfunction and enhanced inflammatory response after myocardial infarction. <i>BBA Clinical</i> , 2016, 5, 159-165.	4.1	2
13	Statin Short-term Inhibition of Insulin Sensitivity and Secretion During Acute Phase of ST-Elevation Myocardial Infarction. <i>Scientific Reports</i> , 2019, 9, 16401.	3.3	2
14	Adverse outcome has a U-shaped relation with acute phase change in insulin sensitivity after ST-Elevation Myocardial Infarction. <i>International Journal of Cardiology</i> , 2018, 254, 16-22.	1.7	1
15	POSCH trial 25-year follow-up results: latest news from an old kid on the block. <i>Clinical Lipidology</i> , 2010, 5, 651-653.	0.4	0