
SÃ;vio Sandes

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/554143/publications.pdf Version: 2024-02-01



SÃ:NO SANDES

#	Article	IF	CITATIONS
1	Co-infection by Salmonella enterica subsp. Enterica serovar typhimurium and Entamoeba dispar pathogenic strains enhances colitis and the expression of amoebic virulence factors. Microbial Pathogenesis, 2021, 158, 105010.	2.9	1
2	Weissella paramesenteroides WpK4 plays an immunobiotic role in gut-brain axis, reducing gut permeability, anxiety-like and depressive-like behaviors in murine models of colitis and chronic stress. Food Research International, 2020, 137, 109741.	6.2	24
3	Differential Immune Response of Lactobacillus plantarum 286 Against Salmonella Typhimurium Infection in Conventional and Germ-Free Mice. Advances in Experimental Medicine and Biology, 2020, 1323, 1-17.	1.6	5
4	Viability of Staphylococcus aureus and expression of its toxins (SEC and TSST-1) in cheeses using Lactobacillus rhamnosus D1 or Weissella paramesenteroides GIR16L4 or both as starter cultures. Journal of Dairy Science, 2020, 103, 4100-4108.	3.4	9
5	Milk Fermented by Lactobacillus paracasei NCC 2461 (ST11) Modulates the Immune Response and Microbiota to Exert its Protective Effects Against Salmonella typhimurium Infection in Mice. Probiotics and Antimicrobial Proteins, 2020, 12, 1398-1408.	3.9	11
6	Virulence factors and antimicrobial resistance of Staphylococcus aureus isolated from the production process of Minas artisanal cheese from the region of Campo das Vertentes, Brazil. Journal of Dairy Science, 2020, 103, 2098-2110.	3.4	18
7	"Physicochemical, immunomodulatory and safety aspects of milks fermented with Lactobacillus paracasei isolated from kefirâ€: Food Research International, 2019, 123, 48-55.	6.2	27
8	Microbial shifts in Minas artisanal cheeses from the Serra do Salitre region of Minas Gerais, Brazil throughout ripening time. Food Microbiology, 2019, 82, 349-362.	4.2	32
9	Probiotic <i>Propionibacterium freudenreichii</i> requires SlpB protein to mitigate mucositis induced by chemotherapy. Oncotarget, 2019, 10, 7198-7219.	1.8	34
10	Selection of starter cultures for the production of sour cassava starch in a pilot-scale fermentation process. Brazilian Journal of Microbiology, 2018, 49, 823-831.	2.0	22
11	<i>Coagulase-Negative Staphylococci</i> Isolated from Human Bloodstream Infections Showed Multidrug Resistance Profile. Microbial Drug Resistance, 2018, 24, 635-647.	2.0	28
12	Protective Effect of Lactobacillus diolivorans 1Z, Isolated From Brazilian Kefir, Against Salmonella enterica Serovar Typhimurium in Experimental Murine Models. Frontiers in Microbiology, 2018, 9, 2856.	3.5	16
13	Microencapsulation of Lactic Acid Bacteria Improves the Gastrointestinal Delivery and in situ Expression of Recombinant Fluorescent Protein. Frontiers in Microbiology, 2018, 9, 2398.	3.5	20
14	In vitro and in vivo evaluation of two potential probiotic lactobacilli isolated from cocoa fermentation (Theobroma cacao L.). Journal of Functional Foods, 2018, 47, 184-191.	3.4	16
15	Selection of Lactic Acid Bacteria with Probiotic Potential Isolated from the Fermentation Process of "Cupua§u―(Theobroma grandiflorum). Advances in Experimental Medicine and Biology, 2017, 973, 1-16.	1.6	6
16	Protective effects of milk fermented by Lactobacillus plantarum B7 from Brazilian artisanal cheese on a Salmonella enterica serovar Typhimurium infection in BALB/c mice. Journal of Functional Foods, 2017, 33, 436-445.	3.4	24
17	Selection of new lactic acid bacteria strains bearing probiotic features from mucosal microbiota of healthy calves: Looking for immunobiotics through in vitro and in vivo approaches for immunobiotics through Research, 2017, 200, 1-13.	5.3	43
18	Milk fermented by <i>Lactobacillus</i> species from Brazilian artisanal cheese protect germ-free-mice against <i>Salmonella</i> Typhimurium infection. Beneficial Microbes, 2017, 8, 579-588.	2.4	21

SÃivio Sandes

#	Article	IF	CITATIONS
19	Evaluation of colonisation resistance in stool of human donors using ex vivo, in vitro and in vivo assays. Beneficial Microbes, 2017, 8, 217-230.	2.4	2
20	Selection of a candidate probiotic strain of <i>Pediococcus pentosaceus</i> from the faecal microbiota of horses by <i>inÂvitro</i> testing and health claims in a mouse model of <i>Salmonella</i> infection. Journal of Applied Microbiology, 2017, 122, 225-238.	3.1	25
21	Culture and molecular identification of microorganisms from Digital Dermatitis lesions in dairy cattle: Leptospira, an unexpected finding. Arquivo Brasileiro De Medicina Veterinaria E Zootecnia, 2017, 69, 559-569.	0.4	0
22	Isolation and identification of lactic acid bacteria from Brazilian Minas artisanal cheese. CYTA - Journal of Food, 2016, , 1-4.	1.9	11
23	Biofilm and toxin profile: A phenotypic and genotypic characterization of coagulase-negative staphylococci isolated from human bloodstream infections. Microbial Pathogenesis, 2016, 100, 312-318.	2.9	20
24	Weissella paramesenteroides WpK4 reduces gene expression of intestinal cytokines, and hepatic and splenic injuries in a murine model of typhoid fever. Beneficial Microbes, 2016, 7, 61-73.	2.4	17
25	Lactic acid microbiota identification in water, raw milk, endogenous starter culture, and fresh Minas artisanal cheese from the Campo das Vertentes region of Brazil during the dry and rainy seasons. Journal of Dairy Science, 2016, 99, 6086-6096.	3.4	39
26	Selection of lactic acid bacteria from Brazilian kefir grains for potential use as starter or probiotic cultures. Anaerobe, 2015, 32, 70-76.	2.1	107
27	Detecção de genes toxigênicos, susceptibilidade antimicrobiana e antagonismo in vitro de Staphylococcus spp. isolados de queijos artesanais. Vigilância Sanitária Em Debate: Sociedade, Ciência & Tecnologia, 2015, .	0.1	Ο
28	Isolation, enumeration, molecular identification and probiotic potential evaluation of lactic acid bacteria isolated from sheep milk. Arquivo Brasileiro De Medicina Veterinaria E Zootecnia, 2014, 66, 940-948.	0.4	14
29	Lactobacillus species identification by amplified ribosomal 16S-23S rRNA restriction fragment length polymorphism analysis. Beneficial Microbes, 2014, 5, 471-481.	2.4	18
30	In vitro assessment of functional properties of lactic acid bacteria isolated from faecal microbiota of healthy dogs for potential use as probiotics. Beneficial Microbes, 2013, 4, 267-275.	2.4	29
31	Genetic diversity and population genetic structure in giant earthworm Rhinodrilus alatus (Annelida:) Tj ETQq1 1	0.784314	rgBT /Overloc