

Ping Ye

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/5565719/publications.pdf>

Version: 2024-02-01

85
papers

1,141
citations

516710

16
h-index

477307

29
g-index

96
all docs

96
docs citations

96
times ranked

2105
citing authors

#	ARTICLE	IF	CITATIONS
1	Prevalence of dyslipidaemia in patients treated with lipid-lowering agents in China: Results of the DYSLipidemia International Study (DYSIS). <i>Atherosclerosis</i> , 2014, 235, 463-469.	0.8	76
2	Mesenchymal stem cells promote type 2 macrophage polarization to ameliorate the myocardial injury caused by diabetic cardiomyopathy. <i>Journal of Translational Medicine</i> , 2019, 17, 251.	4.4	71
3	Controlling Nutritional Status (CONUT) score as a predictor of all-cause mortality in elderly hypertensive patients: a prospective follow-up study. <i>BMJ Open</i> , 2017, 7, e015649.	1.9	63
4	Association between serum homocysteine and arterial stiffness in elderly: a community-based study. <i>Journal of Geriatric Cardiology</i> , 2014, 11, 32-8.	0.2	62
5	Effects of Xuezhikang in patients with dyslipidemia: A multicenter, randomized, placebo-controlled study. <i>Journal of Clinical Lipidology</i> , 2014, 8, 568-575.	1.5	49
6	Brain Natriuretic Peptide and Its Biochemical, Analytical, and Clinical Issues in Heart Failure: A Narrative Review. <i>Frontiers in Physiology</i> , 2018, 9, 692.	2.8	49
7	Effect of Xuezhikang on Cardiovascular Events and Mortality in Elderly Patients with a History of Myocardial Infarction: A Subgroup Analysis of Elderly Subjects from the China Coronary Secondary Prevention Study. <i>Journal of the American Geriatrics Society</i> , 2007, 55, 1015-1022.	2.6	44
8	Mesenchymal stem cells ameliorate myocardial fibrosis in diabetic cardiomyopathy via the secretion of prostaglandin E2. <i>Stem Cell Research and Therapy</i> , 2020, 11, 122.	5.5	43
9	Effect of Aging on the Expression of Peroxisome Proliferator-Activated Receptor β and the Possible Relation to Insulin Resistance. <i>Gerontology</i> , 2006, 52, 69-75.	2.8	37
10	Triglycerides are a predictive factor for arterial stiffness: a community-based 4.8-year prospective study. <i>Lipids in Health and Disease</i> , 2016, 15, 97.	3.0	36
11	Lipid-lowering therapy and lipid goal attainment in patients with metabolic syndrome in China: Subgroup analysis of the Dyslipidemia International Study-China (DYSIS-China). <i>Atherosclerosis</i> , 2014, 237, 99-105.	0.8	25
12	Epidemiological associations between hyperuricemia and cardiometabolic risk factors: a comprehensive study from Chinese community. <i>BMC Cardiovascular Disorders</i> , 2015, 15, 129.	1.7	23
13	Plasma Homocysteine Is Associated with Aortic Arterial Stiffness but not Wave Reflection in Chinese Hypertensive Subjects. <i>PLoS ONE</i> , 2014, 9, e85938.	2.5	23
14	Oncological miR-182-3p, a Novel Smooth Muscle Cell Phenotype Modulator, Evidences From Model Rats and Patients. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2016, 36, 1386-1397.	2.4	22
15	The neutrophil-to-lymphocyte ratio on admission is a good predictor for all-cause mortality in hypertensive patients over 80 years of age. <i>BMC Cardiovascular Disorders</i> , 2017, 17, 167.	1.7	21
16	The prognostic value of the plasma N-terminal pro-brain natriuretic peptide level on all-cause death and major cardiovascular events in a community-based population. <i>Clinical Interventions in Aging</i> , 2016, 11, 245.	2.9	19
17	Biomarkers in Cardiorenal Syndromes. <i>BioMed Research International</i> , 2018, 2018, 1-8.	1.9	19
18	Predictors of Percutaneous Catheter Drainage (PCD) after Abdominal Paracentesis Drainage (APD) in Patients with Moderately Severe or Severe Acute Pancreatitis along with Fluid Collections. <i>PLoS ONE</i> , 2015, 10, e0115348.	2.5	16

#	ARTICLE	IF	CITATIONS
19	Atorvastatin attenuating down-regulation of peroxisome proliferator-activated receptor gamma in preventing cardiac hypertrophy of rats in vitro and in vivo. <i>Journal of Pharmacy and Pharmaceutical Sciences</i> , 2006, 9, 365-75.	2.1	16
20	High-density lipoprotein 3 cholesterol is a predictive factor for arterial stiffness: a community-based 4.8-year prospective study. <i>Lipids in Health and Disease</i> , 2018, 17, 5.	3.0	15
21	The influence of apolipoprotein B and E gene polymorphisms on the response to simvastatin therapy in patients with hyperlipidemia. <i>Chinese Medical Sciences Journal</i> , 2003, 18, 9-13.	0.4	15
22	Plasma Homocysteine is a Predictive Factor for Arterial Stiffness: A Community-Based 4.8-Year Prospective Study. <i>Journal of Clinical Hypertension</i> , 2015, 17, 594-600.	2.0	14
23	Atorvastatin improves pathological changes in the aged kidney by upregulating peroxisome proliferator-activated receptor expression and reducing matrix metalloproteinase-9 and transforming growth factor- β 1 levels. <i>Experimental Gerontology</i> , 2016, 74, 37-42.	2.8	14
24	The abilities of new anthropometric indices in identifying cardiometabolic abnormalities, and influence of residence area and lifestyle on these anthropometric indices in a Chinese community-dwelling population. <i>Clinical Interventions in Aging</i> , 2014, 9, 179.	2.9	13
25	Multimarker Analysis for New Biomarkers in Relation to Central Arterial Stiffness and Hemodynamics in a Chinese Community-Dwelling Population. <i>Angiology</i> , 2015, 66, 950-956.	1.8	13
26	Association of high-sensitivity cardiac troponin T with mortality and cardiovascular events in a community-based prospective study in Beijing. <i>BMJ Open</i> , 2017, 7, e013431.	1.9	13
27	The ability of NT-proBNP to detect chronic heart failure and predict all-cause mortality is higher in elderly Chinese coronary artery disease patients with chronic kidney disease. <i>Clinical Interventions in Aging</i> , 2013, 8, 409.	2.9	12
28	The Association of Homocysteine with Metabolic Syndrome in a Community-Dwelling Population: Homocysteine Might Be Concomitant with Metabolic Syndrome. <i>PLoS ONE</i> , 2014, 9, e113148.	2.5	12
29	Transverse False Tendons in the Left Ventricular Cavity Are Associated with Early Repolarization. <i>PLoS ONE</i> , 2015, 10, e0125173.	2.5	12
30	Effect of physician characteristics and knowledge on the quality of dyslipidemia management and LDL-C target goal achievement in China: Subgroup analysis of the Dyslipidemia International Study. <i>Journal of Global Health</i> , 2017, 7, 020702.	2.7	12
31	Effect of peroxisome proliferator-activated receptor activators on tumor necrosis factor-alpha expression in neonatal rat cardiac myocytes. <i>Chinese Medical Sciences Journal</i> , 2004, 19, 243-7.	0.4	12
32	Age-related decrease in expression of peroxisome proliferator-activated receptor alpha and its effects on development of dyslipidemia. <i>Chinese Medical Journal</i> , 2005, 118, 1093-8.	2.3	12
33	A Multiregional, Randomized Evaluation of the Lipid-Modifying Efficacy and Tolerability of Anacetrapib Added to Ongoing Statin Therapy in Patients With Hypercholesterolemia or Low High-Density Lipoprotein Cholesterol. <i>American Journal of Cardiology</i> , 2017, 120, 569-576.	1.6	11
34	Rapid improvement in carotid adventitial angiogenesis and plaque neovascularization after rosuvastatin therapy in statin treatment-naïve subjects. <i>Journal of Clinical Lipidology</i> , 2019, 13, 847-853.	1.5	11
35	Deep analyses of the associations of a series of biomarkers with insulin resistance, metabolic syndrome, and diabetes risk in nondiabetic middle-aged and elderly individuals: results from a Chinese community-based study. <i>Clinical Interventions in Aging</i> , 2016, Volume 11, 1531-1538.	2.9	10
36	Changes in carotid plaque tissue composition in subjects who continued and discontinued statin therapy. <i>Journal of Clinical Lipidology</i> , 2016, 10, 587-593.	1.5	10

#	ARTICLE	IF	CITATIONS
37	Potential Protective Effect of Long-Term Therapy with Xuezhikang on Left Ventricular Diastolic Function in Patients with Essential Hypertension. <i>Journal of Alternative and Complementary Medicine</i> , 2009, 15, 719-725.	2.1	9
38	Association between high-sensitivity cardiac troponin T and predicted cardiovascular risks in a community-based population. <i>International Journal of Cardiology</i> , 2011, 149, 253-256.	1.7	9
39	Xuezhikang Therapy Increases miR-33 Expression in Patients with Low HDL-C Levels. <i>Disease Markers</i> , 2014, 2014, 1-5.	1.3	9
40	Improving heart function by modulating myocardiocyte autophagy: a possible novel mechanism for cardiovascular protection of high-density lipoprotein. <i>Lipids in Health and Disease</i> , 2014, 13, 163.	3.0	9
41	Noninvasive central pulse pressure is an independent determinant of renal function. <i>Journal of Clinical Hypertension</i> , 2020, 22, 234-242.	2.0	9
42	The predictive capacity and additional prognostic power of N-terminal pro-B-type natriuretic peptide in Chinese elderly with chronic heart failure. <i>Clinical Interventions in Aging</i> , 2015, 10, 359.	2.9	8
43	Persistent lipid abnormalities in statin-treated coronary artery disease patients with and without diabetes in China. <i>International Journal of Cardiology</i> , 2015, 182, 469-475.	1.7	8
44	The relationship of serum alanine aminotransferase normal-range levels to arterial stiffness and metabolic syndrome in non-drinkers and drinkers: a Chinese community-based analysis. <i>BMC Gastroenterology</i> , 2017, 17, 49.	2.0	8
45	The alteration of plasminogen activator inhibitor-1 expression by linoleic acid and fenofibrate in HepG2 cells. <i>Blood Coagulation and Fibrinolysis</i> , 2007, 18, 15-19.	1.0	7
46	Roles of fasting and postprandial blood glucose in the effect of type 2 diabetes on central arterial stiffness: a 5-year prospective community-based analysis. <i>Diabetology and Metabolic Syndrome</i> , 2017, 9, 33.	2.7	7
47	The increase in plasminogen activator inhibitor type-1 expression by stimulation of activators for peroxisome proliferator-activated receptors in human endothelial cells. <i>Chinese Medical Sciences Journal</i> , 2002, 17, 112-6.	0.4	7
48	Urine Albumin Excretion Is Associated with Cardiac Troponin T Detected with a Highly Sensitive Assay in a Community-Based Population. <i>PLoS ONE</i> , 2015, 10, e0135747.	2.5	6
49	The predictive value of arterial stiffness on major adverse cardiovascular events in individuals with mildly impaired renal function. <i>Clinical Interventions in Aging</i> , 2016, Volume 11, 1175-1181.	2.9	6
50	Peripheral arterial stiffness is associated with higher baseline plasma uric acid: A prospective cohort study. <i>Saudi Journal of Biological Sciences</i> , 2017, 24, 574-581.	3.8	6
51	Relationship between Central Arterial Stiffness and Insulin Resistance in Chinese Community-Dwelling Population without Diabetes Mellitus. <i>International Journal of Endocrinology</i> , 2017, 2017, 1-4.	1.5	6
52	Effect of Atorvastatin on Expression of Peroxisome Proliferator-activated Receptor Beta/delta in Angiotensin II-induced Hypertrophic Myocardial Cells In Vitro. <i>Chinese Medical Sciences Journal</i> , 2015, 30, 245-251.	0.4	5
53	Baseline type 2 diabetes had a significant association with elevated high sensitivity cardiac troponin T levels in Chinese community-dwelling population: a 5-year prospective analysis. <i>Nutrition and Metabolism</i> , 2017, 14, 73.	3.0	5
54	Single-marker and multi-marker approaches to appraise the relationships between biomarkers and microalbuminuria in Chinese middle-aged and elderly from communities: a cross-sectional analysis. <i>BMC Nephrology</i> , 2018, 19, 93.	1.8	5

#	ARTICLE	IF	CITATIONS
55	The association between Hcpidin and arterial stiffness in a community-dwelling population. <i>Lipids in Health and Disease</i> , 2018, 17, 244.	3.0	5
56	Body mass index is an independent predictive factor for kidney function evaluated by glomerular filtration rate in a community-dwelling population. <i>Eating and Weight Disorders</i> , 2019, 24, 731-738.	2.5	5
57	Plasma Homocysteine Is a Predictive Factor for Accelerated Renal Function Decline and Chronic Kidney Disease in a Community-Dwelling Population. <i>Kidney and Blood Pressure Research</i> , 2021, 46, 541-549.	2.0	5
58	Correlation between small and dense low-density lipoprotein cholesterol and cardiovascular events in Beijing community population. <i>Journal of Clinical Hypertension</i> , 2021, 23, 345-351.	2.0	5
59	Different types of atrial fibrillation, renal function, and mortality in elderly Chinese patients with coronary artery disease. <i>Clinical Interventions in Aging</i> , 2014, 9, 301.	2.9	4
60	D-4F decreases the expression of A β 2 protein through up-regulating long non coding RNA sirt1-as in SAMP8 mice. <i>Saudi Pharmaceutical Journal</i> , 2017, 25, 517-522.	2.7	4
61	Relationships of drinking and smoking with peripheral arterial stiffness in Chinese community-dwelling population without symptomatic peripheral arterial disease. <i>Tobacco Induced Diseases</i> , 2017, 15, 39.	0.6	4
62	Relationship between age, osteoporosis and coronary artery calcification detected by high-definition computerized tomography in Chinese elderly men. <i>Archives of Gerontology and Geriatrics</i> , 2018, 79, 8-12.	3.0	4
63	Association of arterial stiffness and central hemodynamics with moderately reduced glomerular filtration rate in Chinese middle-aged and elderly community residents: a cross-sectional analysis. <i>BMC Nephrology</i> , 2018, 19, 103.	1.8	4
64	Right ventricle may be involved in regional diastolic dysfunction earliest in primary hypertension patients. <i>Journal of Cellular Biochemistry</i> , 2019, 120, 18088-18093.	2.6	4
65	Valor Prognóstico dos Níveis Plasmáticos de NT-proBNP em Pacientes Hospitalizados com Mais de 80 Anos de Idade em um Hospital em Pequim, China. <i>Arquivos Brasileiros De Cardiologia</i> , 2021, 116, 1027-1036.	0.8	4
66	Homozygous familial hypercholesterolemia in China: Genetic and clinical characteristics from a real-world, multi-center, cohort study. <i>Journal of Clinical Lipidology</i> , 2022, 16, 306-314.	1.5	4
67	Association between resting heart rate and N-terminal pro-brain natriuretic peptide in a community-based population study in Beijing. <i>Clinical Interventions in Aging</i> , 2014, 10, 55.	2.9	3
68	Characteristics of clinical drugs for elderly chronic heart failure complicated with different degrees of renal insufficiency. <i>Pakistan Journal of Medical Sciences</i> , 2018, 34, 135-138.	0.6	3
69	Effect of Intravenous Iron Supplementation on Acute Mountain Sickness: A Preliminary Randomized Controlled Study. <i>Medical Science Monitor</i> , 2015, 21, 2050-2057.	1.1	3
70	Relationship of Arterial Compliance and Blood Pressure with Microalbuminuria and Mildly Decreased Glomerular Filtration Rate: A Chinese Community-Based Analysis. <i>PLoS ONE</i> , 2014, 9, e101013.	2.5	3
71	Plasma homocysteine levels are independently associated with alterations of large artery stiffness in men but not in women. <i>Journal of Geriatric Cardiology</i> , 2015, 12, 251-6.	0.2	3
72	Renal function had an independent relationship with coronary artery calcification in Chinese elderly men. <i>BMC Geriatrics</i> , 2017, 17, 80.	2.7	2

#	ARTICLE	IF	CITATIONS
73	Predictive abilities of cardiovascular biomarkers to rapid decline of renal function in Chinese community-dwelling population: a 5-year prospective analysis. <i>BMC Nephrology</i> , 2017, 18, 331.	1.8	2
74	Relationships of pancreatic beta-cell function with microalbuminuria and glomerular filtration rate in middle-aged and elderly population without type 2 diabetes mellitus: a Chinese community-based analysis. <i>Clinical Interventions in Aging</i> , 2017, Volume 12, 753-757.	2.9	2
75	PCSK9 positively correlates with plasma sdLDL in community-dwelling population but not in diabetic participants after confounder adjustment. <i>Medicine (United States)</i> , 2019, 98, e15062.	1.0	2
76	High-Sensitivity Cardiac Troponin T Is a Risk Factor for Major Adverse Cardiovascular Events and All-Cause Mortality: A 9.5-Year Follow-Up Study. <i>Cardiology Research and Practice</i> , 2021, 2021, 1-8.	1.1	2
77	Activation of peroxisome proliferator-activated receptor alpha in human endothelial cells increases plasminogen activator inhibitor type-1 expression. <i>Chinese Medical Journal</i> , 2003, 116, 29-33.	2.3	2
78	Association between arterial stiffness and risk of coronary artery disease in a community-based population. <i>Chinese Medical Journal</i> , 2014, 127, 3944-7.	2.3	2
79	Homocysteine is associated with plasma high-sensitivity cardiac troponin T levels in a community-dwelling population. <i>Clinical Interventions in Aging</i> , 2013, 9, 79.	2.9	1
80	Relationship between drug application and mortality rate in Chinese older coronary artery disease/chronic heart failure patients with and without low glomerular filtration rate. <i>BMC Pharmacology & Toxicology</i> , 2019, 20, 44.	2.4	1
81	Functions of Monocytes and Macrophages and the Associated Effective Molecules and Mechanisms at the Early Stage of Atherosclerosis. <i>Acta Cardiologica Sinica</i> , 2021, 37, 522-533.	0.2	1
82	Association between high-sensitivity cardiac troponin T and N-terminal pro-brain natriuretic peptide in a community based population. <i>Chinese Medical Journal</i> , 2014, 127, 638-44.	2.3	1
83	Baseline Ratio of Soluble Fas/FasL Predicts Onset of Pulmonary Hypertension in Elder Patients Undergoing Maintenance Hemodialysis: A Prospective Cohort Study. <i>Frontiers in Physiology</i> , 2022, 13, 847172.	2.8	1
84	Effect of Xuezhikang Therapy on Expression of Pulmonary Hypertension Related miR-638 in Patients With Low HDL-C Levels. <i>Frontiers in Pharmacology</i> , 2021, 12, 764046.	3.5	1
85	Serum Alanine Aminotransferase Levels within Normal Range Have Different Associations with Augmentation Index and Other Cardiometabolic Risk Factors in Nondrinkers and Drinkers: A Chinese Community-Based Analysis. <i>Gastroenterology Research and Practice</i> , 2017, 2017, 1-5.	1.5	0