

Claude B Sirlin

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/5573331/publications.pdf>

Version: 2024-02-01

323
papers

35,782
citations

2565

99
h-index

4511

177
g-index

325
all docs

325
docs citations

325
times ranked

26241
citing authors

#	ARTICLE	IF	CITATIONS
1	Diagnosis, Staging, and Management of Hepatocellular Carcinoma: 2018 Practice Guidance by the American Association for the Study of Liver Diseases. <i>Hepatology</i> , 2018, 68, 723-750.	3.6	3,096
2	AASLD guidelines for the treatment of hepatocellular carcinoma. <i>Hepatology</i> , 2018, 67, 358-380.	3.6	2,932
3	Liver Imaging Reporting and Data System (LI-RADS) Version 2018: Imaging of Hepatocellular Carcinoma in At-Risk Patients. <i>Radiology</i> , 2018, 289, 816-830.	3.6	634
4	Quantitative assessment of liver fat with magnetic resonance imaging and spectroscopy. <i>Journal of Magnetic Resonance Imaging</i> , 2011, 34, 729-749.	1.9	613
5	Decoding global gene expression programs in liver cancer by noninvasive imaging. <i>Nature Biotechnology</i> , 2007, 25, 675-680.	9.4	510
6	In vivo characterization of the liver fat ¹ H MR spectrum. <i>NMR in Biomedicine</i> , 2011, 24, 784-790.	1.6	452
7	Utility of magnetic resonance imaging versus histology for quantifying changes in liver fat in nonalcoholic fatty liver disease trials. <i>Hepatology</i> , 2013, 58, 1930-1940.	3.6	434
8	CT and MR Imaging Diagnosis and Staging of Hepatocellular Carcinoma: Part II. Extracellular Agents, Hepatobiliary Agents, and Ancillary Imaging Features. <i>Radiology</i> , 2014, 273, 30-50.	3.6	430
9	Heritability of Nonalcoholic Fatty Liver Disease. <i>Gastroenterology</i> , 2009, 136, 1585-1592.	0.6	419
10	LI-RADS (Liver Imaging Reporting and Data System): Summary, discussion, and consensus of the LI-RADS Management Working Group and future directions. <i>Hepatology</i> , 2015, 61, 1056-1065.	3.6	412
11	Nonalcoholic Fatty Liver Disease: MR Imaging of Liver Proton Density Fat Fraction to Assess Hepatic Steatosis. <i>Radiology</i> , 2013, 267, 422-431.	3.6	410
12	CT and MR Imaging Diagnosis and Staging of Hepatocellular Carcinoma: Part I. Development, Growth, and Spread: Key Pathologic and Imaging Aspects. <i>Radiology</i> , 2014, 272, 635-654.	3.6	401
13	Magnetic resonance elastography predicts advanced fibrosis in patients with nonalcoholic fatty liver disease: A prospective study. <i>Hepatology</i> , 2014, 60, 1920-1928.	3.6	388
14	Proton density fat fraction: A standardized mrâ€¢based biomarker of tissue fat concentration. <i>Journal of Magnetic Resonance Imaging</i> , 2012, 36, 1011-1014.	1.9	385
15	SAFETY Study: Alanine Aminotransferase Cutoff Values Are Set Too High for Reliable Detection of Pediatric Chronic Liver Disease. <i>Gastroenterology</i> , 2010, 138, 1357-1364.e2.	0.6	377
16	Fatty Liver: Imaging Patterns and Pitfalls. <i>Radiographics</i> , 2006, 26, 1637-1653.	1.4	362
17	Insulin resistance drives hepatic de novo lipogenesis in nonalcoholic fatty liver disease. <i>Journal of Clinical Investigation</i> , 2020, 130, 1453-1460.	3.9	362
18	CT and MR Imaging of Extrahepatic Fatty Masses of the Abdomen and Pelvis: Techniques, Diagnosis, Differential Diagnosis, and Pitfalls. <i>Radiographics</i> , 2005, 25, 69-85.	1.4	356

#	ARTICLE	IF	CITATIONS
19	Relaxation effects in the quantification of fat using gradient echo imaging. <i>Magnetic Resonance Imaging</i> , 2008, 26, 347-359.	1.0	356
20	Quantification of Hepatic Steatosis with T1-independent, T2*-corrected MR Imaging with Spectral Modeling of Fat: Blinded Comparison with MR Spectroscopy. <i>Radiology</i> , 2011, 258, 767-775.	3.6	345
21	Epidemiology of hepatocellular carcinoma: target population for surveillance and diagnosis. <i>Abdominal Radiology</i> , 2018, 43, 13-25.	1.0	338
22	Portal chronic inflammation in nonalcoholic fatty liver disease (NAFLD): A histologic marker of advanced NAFLD-Clinicopathologic correlations from the nonalcoholic steatohepatitis clinical research network. <i>Hepatology</i> , 2009, 49, 809-820.	3.6	335
23	Imaging for the diagnosis of hepatocellular carcinoma: A systematic review and meta-analysis. <i>Hepatology</i> , 2018, 67, 401-421.	3.6	329
24	Gadoxetate Disodium-Enhanced MRI of the Liver: Part 1, Protocol Optimization and Lesion Appearance in the Noncirrhotic Liver. <i>American Journal of Roentgenology</i> , 2010, 195, 13-28.	1.0	313
25	Noninvasive, Quantitative Assessment of Liver Fat by MRI-PDF as an Endpoint in NASH Trials. <i>Hepatology</i> , 2018, 68, 763-772.	3.6	299
26	Ezetimibe for the treatment of nonalcoholic steatohepatitis: Assessment by novel magnetic resonance imaging and magnetic resonance elastography in a randomized trial (MOZART trial). <i>Hepatology</i> , 2015, 61, 1239-1250.	3.6	296
27	Nonalcoholic Fatty Liver Disease: Diagnostic and Fat-Grading Accuracy of Low-Flip-Angle Multiecho Gradient-Recalled-Echo MR Imaging at 1.5 T. <i>Radiology</i> , 2009, 251, 67-76.	3.6	287
28	Correlation between liver histology and novel magnetic resonance imaging in adult patients with nonalcoholic fatty liver disease - MRI accurately quantifies hepatic steatosis in NAFLD. <i>Alimentary Pharmacology and Therapeutics</i> , 2012, 36, 22-29.	1.9	285
29	A computed tomography radiogenomic biomarker predicts microvascular invasion and clinical outcomes in hepatocellular carcinoma. <i>Hepatology</i> , 2015, 62, 792-800.	3.6	276
30	MRI and MRE for non-invasive quantitative assessment of hepatic steatosis and fibrosis in NAFLD and NASH: Clinical trials to clinical practice. <i>Journal of Hepatology</i> , 2016, 65, 1006-1016.	1.8	275
31	Sitagliptin vs. placebo for non-alcoholic fatty liver disease: A randomized controlled trial. <i>Journal of Hepatology</i> , 2016, 65, 369-376.	1.8	264
32	Quantification of Liver Fat with Magnetic Resonance Imaging. <i>Magnetic Resonance Imaging Clinics of North America</i> , 2010, 18, 337-357.	0.6	260
33	Estimation of Hepatic Proton-Density Fat Fraction by Using MR Imaging at 3.0 T. <i>Radiology</i> , 2011, 258, 749-759.	3.6	259
34	Optimal threshold of controlled attenuation parameter with MRI-PDF as the gold standard for the detection of hepatic steatosis. <i>Hepatology</i> , 2018, 67, 1348-1359.	3.6	250
35	Fatty Liver Disease: MR Imaging Techniques for the Detection and Quantification of Liver Steatosis. <i>Radiographics</i> , 2009, 29, 231-260.	1.4	246
36	CHCCCA: Consensus terminology for primary liver carcinomas with both hepatocytic and cholangiocytic differentiation. <i>Hepatology</i> , 2018, 68, 113-126.	3.6	244

#	ARTICLE	IF	CITATIONS
37	Accuracy of MR Imagingâ€“estimated Proton Density Fat Fraction for Classification of Dichotomized Histologic Steatosis Grades in Nonalcoholic Fatty Liver Disease. <i>Radiology</i> , 2015, 274, 416-425.	3.6	239
38	Advances in pediatric nonalcoholic fatty liver disease. <i>Hepatology</i> , 2009, 50, 1282-1293.	3.6	235
39	Abdominal Wall Hernias: Imaging Features, Complications, and Diagnostic Pitfalls at Multiâ€“Detector Row CT. <i>Radiographics</i> , 2005, 25, 1501-1520.	1.4	230
40	Evidence Supporting LI-RADS Major Features for CT- and MR Imagingâ€“based Diagnosis of Hepatocellular Carcinoma: A Systematic Review. <i>Radiology</i> , 2018, 286, 29-48.	3.6	230
41	Linearity, Bias, and Precision of Hepatic Proton Density Fat Fraction Measurements by Using MR Imaging: A Meta-Analysis. <i>Radiology</i> , 2018, 286, 486-498.	3.6	225
42	How to perform Contrast-Enhanced Ultrasound (CEUS). <i>Ultrasound International Open</i> , 2018, 04, E2-E15.	0.3	222
43	MR Imaging of Liver Fibrosis: Current State of the Art. <i>Radiographics</i> , 2009, 29, 1615-1635.	1.4	220
44	Effect of colesevelam on liver fat quantified by magnetic resonance in nonalcoholic steatohepatitis: A randomized controlled trial. <i>Hepatology</i> , 2012, 56, 922-932.	3.6	218
45	Quantitative Elastography Methods in Liver Disease: Current Evidence and Future Directions. <i>Radiology</i> , 2018, 286, 738-763.	3.6	215
46	CT Evaluation of Appendicitis and Its Complications: Imaging Techniques and Key Diagnostic Findings. <i>American Journal of Roentgenology</i> , 2005, 185, 406-417.	1.0	214
47	Pediatric Nonalcoholic Fatty Liver Disease. <i>Journal of Pediatric Gastroenterology and Nutrition</i> , 2006, 43, 413-427.	0.9	214
48	Agreement Between Magnetic Resonance Imaging Proton Density Fat Fraction Measurements and Pathologist-Assigned Steatosis Grades of Liver Biopsies From Adults With Nonalcoholic Steatohepatitis. <i>Gastroenterology</i> , 2017, 153, 753-761.	0.6	209
49	Quantification of liver iron with MRI: State of the art and remaining challenges. <i>Journal of Magnetic Resonance Imaging</i> , 2014, 40, 1003-1021.	1.9	208
50	Effect of PRESS and STEAM sequences on magnetic resonance spectroscopic liver fat quantification. <i>Journal of Magnetic Resonance Imaging</i> , 2009, 30, 145-152.	1.9	201
51	Noninvasive Diagnosis of Nonalcoholic Fatty Liver Disease and Quantification of Liver Fat Using a New Quantitative Ultrasound Technique. <i>Clinical Gastroenterology and Hepatology</i> , 2015, 13, 1337-1345.e6.	2.4	200
52	Gadoxetate Disodiumâ€“Enhanced MRI of the Liver: Part 2, Protocol Optimization and Lesion Appearance in the Cirrhotic Liver. <i>American Journal of Roentgenology</i> , 2010, 195, 29-41.	1.0	198
53	Magnetic resonance elastography for staging liver fibrosis in non-alcoholic fatty liver disease: a diagnostic accuracy systematic review and individual participant data pooled analysis. <i>European Radiology</i> , 2016, 26, 1431-1440.	2.3	195
54	T_1 independent, T_2^* corrected chemical shift based fatâ€“water separation with multiâ€“peak fat spectral modeling is an accurate and precise measure of hepatic steatosis. <i>Journal of Magnetic Resonance Imaging</i> , 2011, 33, 873-881.	1.9	183

#	ARTICLE	IF	CITATIONS
55	Disproportionate Fat Stranding: A Helpful CT Sign in Patients with Acute Abdominal Pain. <i>Radiographics</i> , 2004, 24, 703-715.	1.4	180
56	Cirrhosis-associated Hepatocellular Nodules: Correlation of Histopathologic and MR Imaging Features. <i>Radiographics</i> , 2008, 28, 747-769.	1.4	176
57	Liver Fibrosis: Noninvasive Diagnosis with Double Contrast Material-enhanced MR Imaging. <i>Radiology</i> , 2006, 239, 425-437.	3.6	172
58	Magnetic Resonance Imaging Quantification of Liver Iron. <i>Magnetic Resonance Imaging Clinics of North America</i> , 2010, 18, 359-381.	0.6	170
59	Noninvasive screening of diabetics in primary care for NAFLD and advanced fibrosis by MRI and MRE. <i>Alimentary Pharmacology and Therapeutics</i> , 2016, 43, 83-95.	1.9	168
60	Magnetic resonance elastography is superior to acoustic radiation force impulse for the Diagnosis of fibrosis in patients with biopsy-proven nonalcoholic fatty liver disease: A prospective study. <i>Hepatology</i> , 2016, 63, 453-461.	3.6	168
61	Combination of complex-based and magnitude-based multiecho water-fat separation for accurate quantification of fat fraction. <i>Magnetic Resonance in Medicine</i> , 2011, 66, 199-206.	1.9	166
62	Current status of imaging in nonalcoholic fatty liver disease. <i>World Journal of Hepatology</i> , 2018, 10, 530-542.	0.8	166
63	Ultrasound Elastography and MR Elastography for Assessing Liver Fibrosis: Part 2, Diagnostic Performance, Confounders, and Future Directions. <i>American Journal of Roentgenology</i> , 2015, 205, 33-40.	1.0	164
64	Liver fat imaging—a clinical overview of ultrasound, CT, and MR imaging. <i>British Journal of Radiology</i> , 2018, 91, 20170959.	1.0	164
65	Comparative 13-year meta-analysis of the sensitivity and positive predictive value of ultrasound, CT, and MRI for detecting hepatocellular carcinoma. <i>Abdominal Radiology</i> , 2016, 41, 71-90.	1.0	163
66	Effect of a Low Free Sugar Diet vs Usual Diet on Nonalcoholic Fatty Liver Disease in Adolescent Boys. <i>JAMA - Journal of the American Medical Association</i> , 2019, 321, 256.	3.8	163
67	Novel 3D Magnetic Resonance Elastography for the Noninvasive Diagnosis of Advanced Fibrosis in NAFLD: A Prospective Study. <i>American Journal of Gastroenterology</i> , 2016, 111, 986-994.	0.2	160
68	Ultrasound Elastography and MR Elastography for Assessing Liver Fibrosis: Part 1, Principles and Techniques. <i>American Journal of Roentgenology</i> , 2015, 205, 22-32.	1.0	159
69	Longitudinal correlations between MRE, MRI-PDFF, and liver histology in patients with non-alcoholic steatohepatitis: Analysis of data from a phase II trial of selonsertib. <i>Journal of Hepatology</i> , 2019, 70, 133-141.	1.8	149
70	CEUS LI-RADS: algorithm, implementation, and key differences from CT/MRI. <i>Abdominal Radiology</i> , 2018, 43, 127-142.	1.0	147
71	Whole-Body CT Screening: Spectrum of Findings and Recommendations in 1192 Patients. <i>Radiology</i> , 2005, 237, 385-394.	3.6	146
72	MR Contrast Agents for Liver Imaging: What, When, How. <i>Radiographics</i> , 2006, 26, 1621-1636.	1.4	144

#	ARTICLE	IF	CITATIONS
73	Radiogenomic Analysis to Identify Imaging Phenotypes Associated with Drug Response Gene Expression Programs in Hepatocellular Carcinoma. <i>Journal of Vascular and Interventional Radiology</i> , 2007, 18, 821-830.	0.2	144
74	Blunt Abdominal Trauma: Screening US in 2,693 Patients. <i>Radiology</i> , 2001, 218, 352-358.	3.6	142
75	Link between gut microbiome derived metabolite and shared gene effects with hepatic steatosis and fibrosis in NAFLD. <i>Hepatology</i> , 2018, 68, 918-932.	3.6	141
76	Association of Coronary Artery and Aortic Calcium With Lumbar Bone Density: The MESA Abdominal Aortic Calcium Study. <i>American Journal of Epidemiology</i> , 2008, 169, 186-194.	1.6	140
77	Magnetic resonance imaging and liver histology as biomarkers of hepatic steatosis in children with nonalcoholic fatty liver disease. <i>Hepatology</i> , 2015, 61, 1887-1895.	3.6	138
78	Nonalcoholic fatty liver disease with cirrhosis increases familial risk for advanced fibrosis. <i>Journal of Clinical Investigation</i> , 2017, 127, 2697-2704.	3.9	137
79	Chronic thromboembolism: diagnosis with helical CT and MR imaging with angiographic and surgical correlation.. <i>Radiology</i> , 1997, 204, 695-702.	3.6	136
80	Protection from liver fibrosis by a peroxisome proliferator-activated receptor γ agonist. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2012, 109, E1369-76.	3.3	136
81	Quality of life in adults with nonalcoholic fatty liver disease: Baseline data from the nonalcoholic steatohepatitis clinical research network. <i>Hepatology</i> , 2009, 49, 1904-1912.	3.6	133
82	Reproducibility of MRI-determined proton density fat fraction across two different MR scanner platforms. <i>Journal of Magnetic Resonance Imaging</i> , 2011, 34, 928-934.	1.9	130
83	Prevalence of Nonalcoholic Fatty Liver Disease in Children with Obesity. <i>Journal of Pediatrics</i> , 2019, 207, 64-70.	0.9	130
84	Noninvasive Assessment of Hepatic Steatosis. <i>Clinical Gastroenterology and Hepatology</i> , 2009, 7, 135-140.	2.4	129
85	Effect of Weight Loss on Magnetic Resonance Imaging Estimation of Liver Fat and Volume in Patients With Nonalcoholic Steatohepatitis. <i>Clinical Gastroenterology and Hepatology</i> , 2015, 13, 561-568.e1.	2.4	128
86	Comparative diagnostic accuracy of magnetic resonance elastography vs. eight clinical prediction rules for noninvasive diagnosis of advanced fibrosis in biopsy-proven nonalcoholic fatty liver disease: a prospective study. <i>Alimentary Pharmacology and Therapeutics</i> , 2015, 41, 1271-1280.	1.9	125
87	Review article: epidemiology, pathogenesis and potential treatments of paediatric nonalcoholic fatty liver disease. <i>Alimentary Pharmacology and Therapeutics</i> , 2008, 28, 13-24.	1.9	124
88	Association of noninvasive quantitative decline in liver fat content on MRI with histologic response in nonalcoholic steatohepatitis. <i>Therapeutic Advances in Gastroenterology</i> , 2016, 9, 692-701.	1.4	123
89	Diagnostic Challenges and Pitfalls in MR Imaging with Hepatocyte-specific Contrast Agents. <i>Radiographics</i> , 2011, 31, 1547-1568.	1.4	116
90	A Pilot Comparative Study of Quantitative Ultrasound, Conventional Ultrasound, and MRI for Predicting Histology-Determined Steatosis Grade in Adult Nonalcoholic Fatty Liver Disease. <i>American Journal of Roentgenology</i> , 2017, 208, W168-W177.	1.0	113

#	ARTICLE	IF	CITATIONS
91	Hepatocellular carcinoma detection: diagnostic performance of a simulated abbreviated MRI protocol combining diffusion-weighted and T1-weighted imaging at the delayed phase post gadoxetic acid. <i>Abdominal Radiology</i> , 2017, 42, 179-190.	1.0	113
92	Diagnostic accuracy of magnetic resonance imaging hepatic proton density fat fraction in pediatric nonalcoholic fatty liver disease. <i>Hepatology</i> , 2018, 67, 858-872.	3.6	112
93	Carotid Arteries: Contrast-enhanced US Angiographyâ€”Preliminary Clinical Experience. <i>Radiology</i> , 2004, 230, 561-568.	3.6	107
94	Evidence and Recommendations for Imaging Liver Fat in Children, Based on Systematic Review. <i>Clinical Gastroenterology and Hepatology</i> , 2014, 12, 765-773.	2.4	106
95	Shell Osteochondral Allografts of the Knee: Comparison of MR Imaging Findings and Immunologic Responses. <i>Radiology</i> , 2001, 219, 35-43.	3.6	105
96	Cross-sectional Imaging of Extranodal Involvement in Abdominopelvic Lymphoproliferative Malignancies. <i>Radiographics</i> , 2007, 27, 1613-1634.	1.4	105
97	Diagnostic Per-Patient Accuracy of an Abbreviated Hepatobiliary Phase Gadoxetic Acidâ€”Enhanced MRI for Hepatocellular Carcinoma Surveillance. <i>American Journal of Roentgenology</i> , 2015, 204, 527-535.	1.0	105
98	Hepatobiliary agents and their role in LI-RADS. <i>Abdominal Imaging</i> , 2015, 40, 613-625.	2.0	105
99	Association between novel <scp>MRI</scp>â€”estimated pancreatic fat and liver histologyâ€”determined steatosis and fibrosis in nonâ€”alcoholic fatty liver disease. <i>Alimentary Pharmacology and Therapeutics</i> , 2013, 37, 630-639.	1.9	104
100	Magnetic resonance elastography measured shear stiffness as a biomarker of fibrosis in pediatric nonalcoholic fatty liver disease. <i>Hepatology</i> , 2017, 66, 1474-1485.	3.6	103
101	Multisite, multivendor validation of the accuracy and reproducibility of proton-density fat-fraction quantification at 1.5T and 3T using a fat-water phantom. <i>Magnetic Resonance in Medicine</i> , 2017, 77, 1516-1524.	1.9	99
102	Imaging Features of Perivascular Fatty Infiltration of the Liver: Initial Observations. <i>Radiology</i> , 2005, 237, 159-169.	3.6	98
103	Hepatic Fat Quantification. <i>Investigative Radiology</i> , 2012, 47, 368-375.	3.5	98
104	Repeatability of MR Elastography of Liver: A Meta-Analysis. <i>Radiology</i> , 2017, 285, 92-100.	3.6	96
105	Contrast-enhanced ultrasound (CEUS) liver imaging reporting and data system (LI-RADS) 2017 â€” a review of important differences compared to the CT/MRI system. <i>Clinical and Molecular Hepatology</i> , 2017, 23, 280-289.	4.5	96
106	Hypotensive Patients with Blunt Abdominal Trauma: Performance of Screening US. <i>Radiology</i> , 2005, 235, 436-443.	3.6	93
107	<p>LI-RADS: a conceptual and historical review from its beginning to its recent integration into AASLD clinical practice guidance<p>. <i>Journal of Hepatocellular Carcinoma</i> , 2019, Volume 6, 49-69.	1.8	93
108	Blunt Abdominal Trauma: Clinical Value of Negative Screening US Scans. <i>Radiology</i> , 2004, 230, 661-668.	3.6	89

#	ARTICLE	IF	CITATIONS
109	Consensus report from the 7th International Forum for Liver Magnetic Resonance Imaging. <i>European Radiology</i> , 2016, 26, 674-682.	2.3	86
110	Spatial distribution of MRI-determined hepatic proton density fat fraction in adults with nonalcoholic fatty liver disease. <i>Journal of Magnetic Resonance Imaging</i> , 2014, 39, 1525-1532.	1.9	85
111	Association Between Quantity of Liver Fat and Cardiovascular Risk in Patients With Nonalcoholic Fatty Liver Disease Independent of Nonalcoholic Steatohepatitis. <i>Clinical Gastroenterology and Hepatology</i> , 2015, 13, 1513-1520.e1.	2.4	85
112	Imaging Outcomes of Liver Imaging Reporting and Data System Version 2014 Category 2, 3, and 4 Observations Detected at CT and MR Imaging. <i>Radiology</i> , 2016, 281, 129-139.	3.6	85
113	Interreader Reliability of LI-RADS Version 2014 Algorithm and Imaging Features for Diagnosis of Hepatocellular Carcinoma: A Large International Multireader Study. <i>Radiology</i> , 2018, 286, 173-185.	3.6	84
114	LI-RADS Version 2018 Ancillary Features at MRI. <i>Radiographics</i> , 2018, 38, 1973-2001.	1.4	83
115	CT appearance of the normal appendix in adults. <i>European Radiology</i> , 2005, 15, 2096-2103.	2.3	81
116	Optimal phased-array combination for spectroscopy. <i>Magnetic Resonance Imaging</i> , 2008, 26, 847-850.	1.0	81
117	Sebelipase alfa over 52 weeks reduces serum transaminases, liver volume and improves serum lipids in patients with lysosomal acid lipase deficiency. <i>Journal of Hepatology</i> , 2014, 61, 1135-1142.	1.8	81
118	Noninvasive Diagnosis of Nonalcoholic Fatty Liver Disease and Quantification of Liver Fat with Radiofrequency Ultrasound Data Using One-dimensional Convolutional Neural Networks. <i>Radiology</i> , 2020, 295, 342-350.	3.6	79
119	Cisterna Chyli at Routine Abdominal MR Imaging: A Normal Anatomic Structure in the Retrocrural Space. <i>Radiographics</i> , 2004, 24, 809-817.	1.4	76
120	Focal hepatic lesions in Gd-EOB-DTPA enhanced MRI: the atlas. <i>Insights Into Imaging</i> , 2012, 3, 451-474.	1.6	69
121	Predictors of Patient Response to Pulmonary Thromboendarterectomy. <i>American Journal of Roentgenology</i> , 2000, 174, 509-515.	1.0	68
122	Reproducibility of hepatic fat fraction measurement by magnetic resonance imaging. <i>Journal of Magnetic Resonance Imaging</i> , 2013, 37, 1359-1370.	1.9	68
123	Reproducibility of MR-based liver fat quantification across field strength: Same-day comparison between 1.5T and 3T in obese subjects. <i>Journal of Magnetic Resonance Imaging</i> , 2015, 42, 811-817.	1.9	67
124	Shared genetic effects between hepatic steatosis and fibrosis: A prospective twin study. <i>Hepatology</i> , 2016, 64, 1547-1558.	3.6	64
125	Diagnostic per-lesion performance of a simulated gadoxetate disodium-enhanced abbreviated MRI protocol for hepatocellular carcinoma screening. <i>Clinical Radiology</i> , 2018, 73, 485-493.	0.5	63
126	Imaging-Based Diagnostic Systems for Hepatocellular Carcinoma. <i>American Journal of Roentgenology</i> , 2013, 201, 41-55.	1.0	61

#	ARTICLE	IF	CITATIONS
127	Adaptive local window for level set segmentation of CT and MRI liver lesions. <i>Medical Image Analysis</i> , 2017, 37, 46-55.	7.0	59
128	Double-Contrast MRI for Accurate Staging of Hepatocellular Carcinoma in Patients with Cirrhosis. <i>American Journal of Roentgenology</i> , 2008, 190, 47-57.	1.0	58
129	Gadolinium-DTPA-dextran: A macromolecular MR blood pool contrast agent1. <i>Academic Radiology</i> , 2004, 11, 1361-1369.	1.3	57
130	Indeterminate Observations (Liver Imaging Reporting and Data System Category 3) on MRI in the Cirrhotic Liver: Fate and Clinical Implications. <i>American Journal of Roentgenology</i> , 2013, 201, 993-1001.	1.0	57
131	Intra- and inter-examination repeatability of magnetic resonance spectroscopy, magnitude-based MRI, and complex-based MRI for estimation of hepatic proton density fat fraction in overweight and obese children and adults. <i>Abdominal Imaging</i> , 2015, 40, 3070-3077.	2.0	57
132	Associations between histologic features of nonalcoholic fatty liver disease (NAFLD) and quantitative diffusion-weighted MRI measurements in adults. <i>Journal of Magnetic Resonance Imaging</i> , 2015, 41, 1629-1638.	1.9	57
133	Clinical Utility of an Increase in Magnetic Resonance Elastography in Predicting Fibrosis Progression in Nonalcoholic Fatty Liver Disease. <i>Hepatology</i> , 2020, 71, 849-860.	3.6	57
134	Assessment of Hepatic Steatosis in Nonalcoholic Fatty Liver Disease by Using Quantitative US. <i>Radiology</i> , 2020, 295, 106-113.	3.6	57
135	Screening Sonography in Pregnant Patients With Blunt Abdominal Trauma. <i>Journal of Ultrasound in Medicine</i> , 2005, 24, 175-181.	0.8	56
136	Ovarian imaging by magnetic resonance in obese adolescent girls with polycystic ovary syndrome: a pilot study. <i>Fertility and Sterility</i> , 2005, 84, 985-995.	0.5	56
137	White paper of the Society of Abdominal Radiology hepatocellular carcinoma diagnosis disease-focused panel on LI-RADS v2018 for CT and MRI. <i>Abdominal Radiology</i> , 2018, 43, 2625-2642.	1.0	56
138	Liver Imaging Reporting and Data System Category 5: MRI Predictors of Microvascular Invasion and Recurrence After Hepatectomy for Hepatocellular Carcinoma. <i>American Journal of Roentgenology</i> , 2019, 213, 821-830.	1.0	56
139	Consensus report from the 8th International Forum for Liver Magnetic Resonance Imaging. <i>European Radiology</i> , 2020, 30, 370-382.	2.3	55
140	Liver fibrosis imaging: A clinical review of ultrasound and magnetic resonance elastography. <i>Journal of Magnetic Resonance Imaging</i> , 2020, 51, 25-42.	1.9	53
141	Accuracy and the effect of possible subject-based confounders of magnitude-based MRI for estimating hepatic proton density fat fraction in adults, using MR spectroscopy as reference. <i>Journal of Magnetic Resonance Imaging</i> , 2016, 43, 398-406.	1.9	52
142	Noninvasive classification of hepatic fibrosis based on texture parameters from double contrast-enhanced magnetic resonance images. <i>Journal of Magnetic Resonance Imaging</i> , 2012, 36, 1154-1161.	1.9	51
143	Does the Functional Liver Imaging Score Derived from Gadoteric Acid-enhanced MRI Predict Outcomes in Chronic Liver Disease?. <i>Radiology</i> , 2020, 294, 98-107.	3.6	51
144	Acetaminophen Pharmacokinetics in Children With Nonalcoholic Fatty Liver Disease. <i>Journal of Pediatric Gastroenterology and Nutrition</i> , 2011, 52, 198-202.	0.9	50

#	ARTICLE	IF	CITATIONS
145	Bone mineral density and atherosclerosis: The Multi-Ethnic Study of Atherosclerosis, Abdominal Aortic Calcium Study. <i>Atherosclerosis</i> , 2010, 209, 283-289.	0.4	49
146	Serum metabolites detect the presence of advanced fibrosis in derivation and validation cohorts of patients with non-alcoholic fatty liver disease. <i>Gut</i> , 2019, 68, 1884-1892.	6.1	48
147	Inter-examination precision of magnitude-based MRI for estimation of segmental hepatic proton density fat fraction in obese subjects. <i>Journal of Magnetic Resonance Imaging</i> , 2014, 39, 1265-1271.	1.9	47
148	Hepatocellular carcinoma imaging systems: why they exist, how they have evolved, and how they differ. <i>Abdominal Radiology</i> , 2018, 43, 3-12.	1.0	47
149	US of Blunt Abdominal Trauma: Importance of Free Pelvic Fluid in Women of Reproductive Age. <i>Radiology</i> , 2001, 219, 229-235.	3.6	46
150	Application of Modified Spin-Echo-based Sequences for Hepatic MR Elastography: Evaluation, Comparison with the Conventional Gradient-Echo Sequence, and Preliminary Clinical Experience. <i>Radiology</i> , 2017, 282, 390-398.	3.6	46
151	Intravenous Gadoxetate Disodium Administration Reduces Breath-holding Capacity in the Hepatic Arterial Phase: A Multi-Center Randomized Placebo-controlled Trial. <i>Radiology</i> , 2017, 282, 361-368.	3.6	46
152	Liver Imaging Reporting and Data System: an expert consensus statement. <i>Journal of Hepatocellular Carcinoma</i> , 2017, Volume 4, 29-39.	1.8	46
153	How bariatric surgery affects liver volume and fat density in NAFLD patients. <i>Surgical Endoscopy and Other Interventional Techniques</i> , 2018, 32, 1675-1682.	1.3	46
154	Role of US LI-RADS in the LI-RADS Algorithm. <i>Radiographics</i> , 2019, 39, 690-708.	1.4	45
155	Hepatocellular carcinoma detection in liver cirrhosis: diagnostic performance of contrast-enhanced CT vs. MRI with extracellular contrast vs. gadoxetic acid. <i>European Radiology</i> , 2020, 30, 1020-1030.	2.3	45
156	Lower serum hepcidin and greater parenchymal iron in nonalcoholic fatty liver disease patients with C282Y/HFE mutations. <i>Hepatology</i> , 2012, 56, 1730-1740.	3.6	44
157	In vivo triglyceride composition of abdominal adipose tissue measured by ¹ H MRS at 3T. <i>Journal of Magnetic Resonance Imaging</i> , 2017, 45, 1455-1463.	1.9	44
158	Contrast-enhanced B-mode US angiography in the assessment of experimental in vivo and in vitro atherosclerotic disease. <i>Academic Radiology</i> , 2001, 8, 162-172.	1.3	43
159	Importance of evaluating organ parenchyma during screening abdominal ultrasonography after blunt trauma.. <i>Journal of Ultrasound in Medicine</i> , 2001, 20, 577-583.	0.8	43
160	Chest Radiography with a Flat-Panel Detector: Image Quality with Dose Reduction after Copper Filtration. <i>Radiology</i> , 2005, 237, 691-700.	3.6	43
161	Assessment of liver fat quantification in the presence of iron. <i>Magnetic Resonance Imaging</i> , 2010, 28, 767-776.	1.0	43
162	Effect of flip angle on the accuracy and repeatability of hepatic proton density fat fraction estimation by complex data-based, T1-independent, T2*-corrected, spectrum-modeled MRI. <i>Journal of Magnetic Resonance Imaging</i> , 2014, 39, 440-447.	1.9	43

#	ARTICLE	IF	CITATIONS
163	LI-RADS Categorization of Benign and Likely Benign Findings in Patients at Risk of Hepatocellular Carcinoma: A Pictorial Atlas. <i>American Journal of Roentgenology</i> , 2014, 203, W48-W69.	1.0	43
164	Repeatability and Reproducibility of the Ultrasonic Attenuation Coefficient and Backscatter Coefficient Measured in the Right Lobe of the Liver in Adults With Known or Suspected Nonalcoholic Fatty Liver Disease. <i>Journal of Ultrasound in Medicine</i> , 2018, 37, 1913-1927.	0.8	43
165	Abbreviated MRI for Hepatocellular Carcinoma Screening and Surveillance. <i>Radiographics</i> , 2020, 40, 1916-1931.	1.4	43
166	Gadoxetate-enhanced abbreviated MRI is highly accurate for hepatocellular carcinoma screening. <i>European Radiology</i> , 2020, 30, 6003-6013.	2.3	43
167	MRI of the Female Pelvis Using Vaginal Gel. <i>American Journal of Roentgenology</i> , 2005, 185, 1221-1227.	1.0	42
168	Insulin Resistance Increases MRI-Estimated Pancreatic Fat in Nonalcoholic Fatty Liver Disease and Normal Controls. <i>Gastroenterology Research and Practice</i> , 2013, 2013, 1-8.	0.7	42
169	Pediatric Hepatoblastoma, Hepatocellular Carcinoma, and Other Hepatic Neoplasms: Consensus Imaging Recommendations from American College of Radiology Pediatric Liver Reporting and Data System (LI-RADS) Working Group. <i>Radiology</i> , 2020, 296, 493-497.	3.6	42
170	Introduction to the Liver Imaging Reporting and Data System for Hepatocellular Carcinoma. <i>Clinical Gastroenterology and Hepatology</i> , 2019, 17, 1228-1238.	2.4	41
171	Pediatric Nonalcoholic Fatty Liver Disease: A Comprehensive Review. <i>Advances in Pediatrics</i> , 2010, 57, 85-140.	0.5	40
172	Consensus report from the 6th International forum for liver MRI using gadoxetic acid. <i>Journal of Magnetic Resonance Imaging</i> , 2014, 40, 516-529.	1.9	40
173	Toward a standardized system for hepatocellular carcinoma diagnosis using computed tomography and MRI. <i>Expert Review of Gastroenterology and Hepatology</i> , 2013, 7, 269-279.	1.4	39
174	Understanding LI-RADS. <i>Magnetic Resonance Imaging Clinics of North America</i> , 2014, 22, 337-352.	0.6	39
175	LI-RADS® algorithm: CT and MRI. <i>Abdominal Radiology</i> , 2018, 43, 111-126.	1.0	39
176	Linearity and Bias of Proton Density Fat Fraction as a Quantitative Imaging Biomarker: A Multicenter, Multiplatform, Multivendor Phantom Study. <i>Radiology</i> , 2021, 298, 640-651.	3.6	39
177	Screening US for Blunt Abdominal Trauma: Objective Predictors of False-Negative Findings and Missed Injuries. <i>Radiology</i> , 2003, 229, 766-774.	3.6	38
178	Consensus Report of the Fifth International Forum for Liver MRI. <i>American Journal of Roentgenology</i> , 2013, 201, 97-107.	1.0	38
179	Quantifying Abdominal Adipose Tissue and Thigh Muscle Volume and Hepatic Proton Density Fat Fraction: Repeatability and Accuracy of an MR Imaging-based, Semiautomated Analysis Method. <i>Radiology</i> , 2017, 283, 438-449.	3.6	38
180	Deep learning assisted differentiation of hepatocellular carcinoma from focal liver lesions: choice of four-phase and three-phase CT imaging protocol. <i>Abdominal Radiology</i> , 2020, 45, 2688-2697.	1.0	37

#	ARTICLE	IF	CITATIONS
181	Acute appendicitis: diagnostic value of nonenhanced CT with selective use of contrast in routine clinical settings. <i>European Radiology</i> , 2007, 17, 2055-2061.	2.3	35
182	Cross-sectional investigation of correlation between hepatic steatosis and IVIM perfusion on MR imaging. <i>Magnetic Resonance Imaging</i> , 2012, 30, 572-578.	1.0	35
183	Cardiac motion in diffusion-weighted MRI of the liver: artifact and a method of correction. <i>Journal of Magnetic Resonance Imaging</i> , 2012, 35, 318-327.	1.9	35
184	Estimation of Fish and ω -3 Fatty Acid Intake in Pediatric Nonalcoholic Fatty Liver Disease. <i>Journal of Pediatric Gastroenterology and Nutrition</i> , 2013, 57, 627-633.	0.9	35
185	How reader perception of capsule affects interpretation of washout in hypervascular liver nodules in patients at risk for hepatocellular carcinoma. <i>Journal of Magnetic Resonance Imaging</i> , 2016, 43, 1337-1345.	1.9	35
186	Randomised clinical trial: a leucine-metformin-sildenafil combination (NS0200) vs placebo in patients with non-alcoholic fatty liver disease. <i>Alimentary Pharmacology and Therapeutics</i> , 2018, 47, 1639-1651.	1.9	35
187	Constraining the initial phase in water-fat separation. <i>Magnetic Resonance Imaging</i> , 2011, 29, 216-221.	1.0	34
188	Iron Deficiency in Patients With Nonalcoholic Fatty Liver Disease Is Associated With Obesity, Female Gender, and Low Serum Hepcidin. <i>Clinical Gastroenterology and Hepatology</i> , 2014, 12, 1170-1178.	2.4	34
189	Repeatability and reproducibility of 2D and 3D hepatic MR elastography with rigid and flexible drivers at end-expiration and end-inspiration in healthy volunteers. <i>Abdominal Radiology</i> , 2017, 42, 2843-2854.	1.0	34
190	LI-RADS 2017: An update. <i>Journal of Magnetic Resonance Imaging</i> , 2018, 47, 1459-1474.	1.9	34
191	Screening Ultrasound in Blunt Abdominal Trauma. <i>Journal of Intensive Care Medicine</i> , 2003, 18, 253-260.	1.3	33
192	Cirrhotic liver: What's that nodule? The LI-RADS approach. <i>Journal of Magnetic Resonance Imaging</i> , 2016, 43, 281-294.	1.9	33
193	LI-RADS categories: concepts, definitions, and criteria. <i>Abdominal Radiology</i> , 2018, 43, 101-110.	1.0	33
194	MRI Assessment of Treatment Response in HIV-associated NAFLD: A Randomized Trial of a Stearoyl-Coenzyme A Desaturase Inhibitor (ARRIVE Trial). <i>Hepatology</i> , 2019, 70, 1531-1545.	3.6	33
195	Effect of acquisition rate on liver and portal vein enhancement with microbubble contrast. <i>Ultrasound in Medicine and Biology</i> , 1999, 25, 331-338.	0.7	32
196	In vivo breath-hold ¹ H MRS simultaneous estimation of liver proton density fat fraction, and <i>T</i> ₁ and <i>T</i> ₂ of water and fat, with a multi-TR, multi-TE sequence. <i>Journal of Magnetic Resonance Imaging</i> , 2015, 42, 1538-1543.	1.9	32
197	CT/MRI and CEUS LI-RADS Major Features Association with Hepatocellular Carcinoma: Individual Patient Data Meta-Analysis. <i>Radiology</i> , 2022, 302, 326-335.	3.6	32
198	Magnetic resonance elastography identifies fibrosis in adults with alpha ₁ -antitrypsin deficiency liver disease: a prospective study. <i>Alimentary Pharmacology and Therapeutics</i> , 2016, 44, 287-299.	1.9	31

#	ARTICLE	IF	CITATIONS
199	Increased severity of liver fat content and liver fibrosis in non-alcoholic fatty liver disease correlate with epicardial fat volume in type 2 diabetes: A prospective study. <i>European Radiology</i> , 2018, 28, 1345-1355.	2.3	31
200	Unusual cause of ureteral obstruction in transplant kidney. <i>Abdominal Imaging</i> , 2006, 31, 379-382.	2.0	30
201	Nephrogenic systemic fibrosis in liver disease: A systematic review. <i>Journal of Magnetic Resonance Imaging</i> , 2009, 30, 1313-1322.	1.9	29
202	Elastography to assess the stage of liver fibrosis in children: Concepts, opportunities, and challenges. <i>Clinical Liver Disease</i> , 2017, 9, 5-10.	1.0	29
203	Weight Loss Decreases Magnetic Resonance Elastography Estimated Liver Stiffness in Nonalcoholic Fatty Liver Disease. <i>Clinical Gastroenterology and Hepatology</i> , 2017, 15, 463-464.	2.4	29
204	Hepatocarcinogenesis and LI-RADS. <i>Abdominal Radiology</i> , 2018, 43, 158-168.	1.0	29
205	Inter-sonographer reproducibility of quantitative ultrasound outcomes and shear wave speed measured in the right lobe of the liver in adults with known or suspected non-alcoholic fatty liver disease. <i>European Radiology</i> , 2018, 28, 4992-5000.	2.3	29
206	Evaluation of Liver Fibrosis Using Texture Analysis on Combined-Contrast-Enhanced Magnetic Resonance Images at 3.0T. <i>BioMed Research International</i> , 2015, 2015, 1-12.	0.9	28
207	Screening and Surveillance of Hepatocellular Carcinoma. <i>Radiologic Clinics of North America</i> , 2017, 55, 1197-1209.	0.9	28
208	CT/MR LI-RADS 2018: clinical implications and management recommendations. <i>Abdominal Radiology</i> , 2019, 44, 1306-1322.	1.0	28
209	Hepatic R2* is more strongly associated with proton density fat fraction than histologic liver iron scores in patients with nonalcoholic fatty liver disease. <i>Journal of Magnetic Resonance Imaging</i> , 2019, 49, 1456-1466.	1.9	28
210	Deep convolutional neural network applied to the liver imaging reporting and data system (LI-RADS) version 2014 category classification: a pilot study. <i>Abdominal Radiology</i> , 2020, 45, 24-35.	1.0	28
211	Nonstandardized Terminology to Describe Focal Liver Lesions in Patients at Risk for Hepatocellular Carcinoma: Implications Regarding Clinical Communication. <i>American Journal of Roentgenology</i> , 2018, 210, 85-90.	1.0	27
212	MRI proton density fat fraction is robust across the biologically plausible range of triglyceride spectra in adults with nonalcoholic steatohepatitis. <i>Journal of Magnetic Resonance Imaging</i> , 2018, 47, 995-1002.	1.9	27
213	Recommendation for terminology: Nodules without arterial phase hyperenhancement and with hepatobiliary phase hypointensity in chronic liver disease. <i>Journal of Magnetic Resonance Imaging</i> , 2018, 48, 1169-1171.	1.9	27
214	Quantification of fluid on screening ultrasonography for blunt abdominal trauma: a simple scoring system to predict severity of injury. <i>Journal of Ultrasound in Medicine</i> , 2001, 20, 359-364.	0.8	26
215	Inter-platform reproducibility of ultrasonic attenuation and backscatter coefficients in assessing NAFLD. <i>European Radiology</i> , 2019, 29, 4699-4708.	2.3	26
216	Accuracy of multiecho magnitude-based MRI (M ₀ -MRI) for estimation of hepatic proton density fat fraction (PDFF) in children. <i>Journal of Magnetic Resonance Imaging</i> , 2015, 42, 1223-1232.	1.9	25

#	ARTICLE	IF	CITATIONS
217	Liver histology and diffusion-weighted MRI in children with nonalcoholic fatty liver disease: A MAGNET study. <i>Journal of Magnetic Resonance Imaging</i> , 2017, 46, 1149-1158.	1.9	25
218	LI-RADS and transplantation for hepatocellular carcinoma. <i>Abdominal Radiology</i> , 2018, 43, 193-202.	1.0	24
219	Relationship between resolution of nonalcoholic steatohepatitis and changes in lipoprotein subfractions: a posthoc analysis of the PIVENS trial. <i>Alimentary Pharmacology and Therapeutics</i> , 2019, 49, 1205-1213.	1.9	24
220	Patterns of fluid accumulation on screening ultrasonography for blunt abdominal trauma: comparison with site of injury. <i>Journal of Ultrasound in Medicine</i> , 2001, 20, 351-357.	0.8	23
221	Pitfalls and problems to be solved in the diagnostic CT/MRI Liver Imaging Reporting and Data System (LI-RADS). <i>European Radiology</i> , 2019, 29, 1124-1132.	2.3	23
222	Use of gadoxetate disodium in patients with chronic liver disease and its implications for liver imaging reporting and data system (LI-RADS). <i>Journal of Magnetic Resonance Imaging</i> , 2019, 49, 1236-1252.	1.9	23
223	Normal range for MR elastography measured liver stiffness in children without liver disease. <i>Journal of Magnetic Resonance Imaging</i> , 2020, 51, 919-927.	1.9	23
224	Effects of intravenous gadolinium administration and flip angle on the assessment of liver fat signal fraction with opposed-phase and in-phase imaging. <i>Journal of Magnetic Resonance Imaging</i> , 2008, 28, 246-251.	1.9	22
225	Update on the Liver Imaging Reporting and Data System. <i>Advances in Anatomic Pathology</i> , 2015, 22, 314-322.	2.4	22
226	Cross-sectional and longitudinal evaluation of liver volume and total liver fat burden in adults with nonalcoholic steatohepatitis. <i>Abdominal Imaging</i> , 2015, 40, 26-37.	2.0	22
227	The Role of Preoperative Dynamic Contrast-enhanced 3.0-T MR Imaging in Predicting Early Recurrence in Patients With Early-Stage Hepatocellular Carcinomas After Curative Resection. <i>Frontiers in Oncology</i> , 2019, 9, 1336.	1.3	22
228	Monitoring Fatty Liver Disease with MRI Following Bariatric Surgery: A Prospective, Dual-Center Study. <i>Radiology</i> , 2019, 290, 682-690.	3.6	22
229	Superparamagnetic iron oxides and low molecular weight gadolinium chelates are synergistic for direct visualization of advanced liver fibrosis. <i>Journal of Magnetic Resonance Imaging</i> , 2007, 26, 728-737.	1.9	21
230	Assessment of treatment response in nonalcoholic steatohepatitis using advanced magnetic resonance imaging. <i>Alimentary Pharmacology and Therapeutics</i> , 2017, 45, 844-854.	1.9	21
231	Collagen Formation Assessed by N-Terminal Propeptide of Type 3 Procollagen Is a Heritable Trait and Is Associated With Liver Fibrosis Assessed by Magnetic Resonance Elastography. <i>Hepatology</i> , 2019, 70, 127-141.	3.6	21
232	Feasibility of and agreement between MR imaging and spectroscopic estimation of hepatic proton density fat fraction in children with known or suspected nonalcoholic fatty liver disease. <i>Abdominal Imaging</i> , 2015, 40, 3084-3090.	2.0	20
233	Optimization of region-of-interest sampling strategies for hepatic MRI proton density fat fraction quantification. <i>Journal of Magnetic Resonance Imaging</i> , 2018, 47, 988-994.	1.9	20
234	Longitudinal evolution of CT and MRI LI-RADS v2014 category 1, 2, 3, and 4 observations. <i>European Radiology</i> , 2019, 29, 5073-5081.	2.3	20

#	ARTICLE	IF	CITATIONS
235	Hepatocellular adenomas: Understanding the pathomolecular lexicon, MRI features, terminology, and pitfalls to inform a standardized approach. <i>Journal of Magnetic Resonance Imaging</i> , 2020, 51, 1630-1640.	1.9	20
236	Evaluation of MRI fat fraction in the liver and spine pre and post SPIO infusion. <i>Magnetic Resonance Imaging</i> , 2013, 31, 1012-1016.	1.0	19
237	Risk of nephrogenic systemic fibrosis is low in patients with chronic liver disease exposed to gadolinium-based contrast agents. <i>Journal of Magnetic Resonance Imaging</i> , 2015, 41, 1259-1267.	1.9	19
238	Accuracy of PDFF estimation by magnitude-based and complex-based MRI in children with MR spectroscopy as a reference. <i>Journal of Magnetic Resonance Imaging</i> , 2017, 46, 1641-1647.	1.9	19
239	Cross-sectional correlation between hepatic R2* and proton density fat fraction (PDFF) in children with hepatic steatosis. <i>Journal of Magnetic Resonance Imaging</i> , 2018, 47, 418-424.	1.9	19
240	Spectrum of Pitfalls, Pseudolesions, and Potential Misdiagnoses in Cirrhosis. <i>American Journal of Roentgenology</i> , 2018, 211, 87-96.	1.0	19
241	Inter- and intra-reader agreement for gadoxetic acid-enhanced MRI parameter readings in patients with chronic liver diseases. <i>European Radiology</i> , 2019, 29, 6600-6610.	2.3	19
242	Comparative diagnostic performance of ultrasound shear wave elastography and magnetic resonance elastography for classifying fibrosis stage in adults with biopsy-proven nonalcoholic fatty liver disease. <i>European Radiology</i> , 2022, 32, 2457-2469.	2.3	19
243	Polydioxanone Biodegradable Pins in the Knee. <i>American Journal of Roentgenology</i> , 2001, 176, 83-90.	1.0	18
244	Science to Practice: Can T1-weighted Imaging Be Used to Diagnose and Assess the Severity of Hepatic Fibrosis?. <i>Radiology</i> , 2011, 259, 619-620.	3.6	18
245	Robustness of fat quantification using chemical shift imaging. <i>Magnetic Resonance Imaging</i> , 2012, 30, 151-157.	1.0	18
246	Effect of echo-sampling strategy on the accuracy of out-of-phase and in-phase multiecho gradient-echo MRI hepatic fat fraction estimation. <i>Journal of Magnetic Resonance Imaging</i> , 2014, 39, 567-575.	1.9	18
247	User and system pitfalls in liver imaging with LI-RADS. <i>Journal of Magnetic Resonance Imaging</i> , 2019, 50, 1673-1686.	1.9	18
248	Technology Insight: advances in liver imaging. <i>Nature Reviews Gastroenterology & Hepatology</i> , 2007, 4, 215-228.	1.7	17
249	Automated registration of sequential breath-hold dynamic contrast-enhanced MR images: a comparison of three techniques. <i>Magnetic Resonance Imaging</i> , 2011, 29, 668-682.	1.0	17
250	Accurate diagnosis of nonalcoholic fatty liver disease in human participants via quantitative ultrasound. , 2014, , .		16
251	Reader agreement and accuracy of ultrasound features for hepatic steatosis. <i>Abdominal Radiology</i> , 2019, 44, 54-64.	1.0	16
252	Pilot study on longitudinal change in pancreatic proton density fat fraction during a weight-loss surgery program in adults with obesity. <i>Journal of Magnetic Resonance Imaging</i> , 2019, 50, 1092-1102.	1.9	16

#	ARTICLE	IF	CITATIONS
253	A Pilot Genome-Wide Analysis Study Identifies Loci Associated With Response to Obeticholic Acid in Patients With NASH. <i>Hepatology Communications</i> , 2019, 3, 1571-1584.	2.0	16
254	Accuracy of common proton density fat fraction thresholds for magnitude- and complex-based chemical shift-encoded MRI for assessing hepatic steatosis in patients with obesity. <i>Abdominal Radiology</i> , 2020, 45, 661-671.	1.0	16
255	Up-to-Date Role of CT/MRI LI-RADS in Hepatocellular Carcinoma. <i>Journal of Hepatocellular Carcinoma</i> , 2021, Volume 8, 513-527.	1.8	16
256	Capsular retraction: an uncommon imaging finding in hepatic inflammatory pseudotumour. <i>British Journal of Radiology</i> , 2009, 82, e256-e260.	1.0	15
257	Management implications and outcomes of LI-RADS-2, -3, -4, and -M category observations. <i>Abdominal Radiology</i> , 2018, 43, 143-148.	1.0	15
258	Diagnostic Accuracy of Preoperative Gadoteric Acid-enhanced 3-T MR Imaging for Malignant Liver Lesions by Using Ex Vivo MR Imaging-matched Pathologic Findings as the Reference Standard. <i>Radiology</i> , 2015, 276, 775-786.	3.6	14
259	Bone marrow fat content is correlated with hepatic fat content in paediatric non-alcoholic fatty liver disease. <i>Clinical Radiology</i> , 2017, 72, 425.e9-425.e14.	0.5	14
260	Effect of threshold growth as a major feature on LI-RADS categorization. <i>Abdominal Radiology</i> , 2017, 42, 2089-2100.	1.0	14
261	Sources of systematic error in proton density fat fraction (PDFF) quantification in the liver evaluated from magnitude images with different numbers of echoes. <i>NMR in Biomedicine</i> , 2018, 31, e3843.	1.6	14
262	Is It Time to Expand the Definition of Washout Appearance in LI-RADS?. <i>Radiology</i> , 2019, 291, 658-659.	3.6	14
263	A Quantitative Approach to Sequence and Image Weighting. <i>Journal of Computer Assisted Tomography</i> , 2010, 34, 317-331.	0.5	13
264	Amphicrine carcinoma of the liver. <i>Annals of Diagnostic Pathology</i> , 2011, 15, 355-357.	0.6	13
265	Error model for reduction of cardiac and respiratory motion effects in quantitative liver DW-MRI. <i>Magnetic Resonance in Medicine</i> , 2013, 70, 1460-1469.	1.9	13
266	LI-RADS version 2018: What is new and what does this mean to my radiology reports?. <i>Abdominal Radiology</i> , 2019, 44, 41-42.	1.0	13
267	How to Use LI-RADS to Report Liver CT and MRI Observations. <i>Radiographics</i> , 2021, 41, 1352-1367.	1.4	13
268	LI-RADS ancillary features on contrast-enhanced ultrasonography. <i>Ultrasonography</i> , 2020, 39, 221-228.	1.0	13
269	Liver imaging: it is time to adopt standardized terminology. <i>European Radiology</i> , 2022, 32, 6291-6301.	2.3	13
270	B-Mode Enhancement of the Liver with Microbubble Contrast Agent. <i>Academic Radiology</i> , 2001, 8, 734-740.	1.3	12

#	ARTICLE	IF	CITATIONS
271	Cardiovascular risk assessment in the treatment of nonalcoholic steatohepatitis: a secondary analysis of the MOZART trial. <i>Therapeutic Advances in Gastroenterology</i> , 2016, 9, 152-161.	1.4	12
272	LI-RADS: a glimpse into the future. <i>Abdominal Radiology</i> , 2018, 43, 231-236.	1.0	12
273	Cirrhosis and LI-RADS. <i>Abdominal Radiology</i> , 2018, 43, 26-40.	1.0	12
274	Editorial on the Current Role of Ultrasound. <i>Applied Sciences (Switzerland)</i> , 2019, 9, 3512.	1.3	12
275	Time to Clarify Common Misconceptions about the Liver Imaging Reporting and Data System for Contrast-enhanced US. <i>Radiology</i> , 2020, 295, 245-247.	3.6	12
276	CT and MRI of diffuse lobar involvement pattern in liver pathology. <i>Diagnostic and Interventional Radiology</i> , 2010, 17, 334-42.	0.7	12
277	MR elastography in nonalcoholic fatty liver disease: inter-center and inter-analysis-method measurement reproducibility and accuracy at 3T. <i>European Radiology</i> , 2022, 32, 2937-2948.	2.3	12
278	Direct Comparison of Quantitative US versus Controlled Attenuation Parameter for Liver Fat Assessment Using MRI Proton Density Fat Fraction as the Reference Standard in Patients Suspected of Having NAFLD. <i>Radiology</i> , 2022, , 211131.	3.6	12
279	Invited Commentary. <i>Radiographics</i> , 2009, 29, 1277-1280.	1.4	11
280	Apparent Diffusion Coefficient of Fibrosis and Regenerative Nodules in the Cirrhotic Liver at MRI. <i>American Journal of Roentgenology</i> , 2010, 194, 1515-1522.	1.0	11
281	Histogram Analysis of Hepatobiliary Phase MR Imaging as a Quantitative Value for Liver Cirrhosis: Preliminary Observations. <i>Yonsei Medical Journal</i> , 2014, 55, 651.	0.9	11
282	Liver Imaging Reporting and Data System: Review of Ancillary Imaging Features. <i>Seminars in Roentgenology</i> , 2016, 51, 301-307.	0.2	11
283	Hepatic steatosis and reduction in steatosis following bariatric weight loss surgery differs between segments and lobes. <i>European Radiology</i> , 2019, 29, 2474-2480.	2.3	11
284	Effect of shot number on the calculated apparent diffusion coefficient in phantoms and in human liver in diffusion-weighted echo-planar imaging. <i>Journal of Magnetic Resonance Imaging</i> , 2009, 30, 547-553.	1.9	10
285	The LI-RADS adventure—a personal statement. <i>Abdominal Radiology</i> , 2018, 43, 1-2.	1.0	10
286	Diagnostic performance of LI-RADS version 2018 in differentiating hepatocellular carcinoma from other hepatic malignancies in patients with hepatitis B virus infection. <i>Bosnian Journal of Basic Medical Sciences</i> , 2020, 20, 401-410.	0.6	10
287	Staging of fibrosis in experimental non-alcoholic steatohepatitis by quantitative molecular imaging in rat models. <i>Nuclear Medicine and Biology</i> , 2016, 43, 179-187.	0.3	9
288	Epidemiology of Hepatic Steatosis at a Tertiary Care Center. <i>Academic Radiology</i> , 2018, 25, 317-327.	1.3	9

#	ARTICLE	IF	CITATIONS
289	Surveillance of hepatocellular carcinoma by medical imaging. <i>Quantitative Imaging in Medicine and Surgery</i> , 2019, 9, 1904-1910.	1.1	9
290	Assessment of a high-SNR chemical-shift-encoded MRI with complex reconstruction for proton density fat fraction (PDFF) estimation overall and in the low-fat range. <i>Journal of Magnetic Resonance Imaging</i> , 2019, 49, 229-238.	1.9	9
291	The relationship between liver triglyceride composition and proton density fat fraction as assessed by ¹ H MRS. <i>NMR in Biomedicine</i> , 2020, 33, e4286.	1.6	9
292	LI-RADS ancillary feature prediction of longitudinal category changes in LR-3 observations: an exploratory study. <i>Abdominal Radiology</i> , 2020, 45, 3092-3102.	1.0	9
293	Female Pelvis. <i>Magnetic Resonance Imaging Clinics of North America</i> , 2005, 13, 381-395.	0.6	8
294	Body MRI artefacts: from image degradation to diagnostic utility. <i>Radiologia Medica</i> , 2009, 114, 18-31.	4.7	8
295	Inter-reader agreement of magnetic resonance imaging proton density fat fraction and its longitudinal change in a clinical trial of adults with nonalcoholic steatohepatitis. <i>Abdominal Radiology</i> , 2019, 44, 482-492.	1.0	8
296	The Dose-Response Effects of Consuming High Fructose Corn Syrup-Sweetened Beverages on Hepatic Lipid Content and Insulin Sensitivity in Young Adults. <i>Nutrients</i> , 2022, 14, 1648.	1.7	8
297	Noninvasive imaging biomarkers for steatosis assessment. <i>Liver Transplantation</i> , 2009, 15, 1389-1391.	1.3	7
298	On confirmation bias in imaging research. <i>Journal of Magnetic Resonance Imaging</i> , 2015, 41, 1163-1164.	1.9	7
299	Measurement of spleen fat on MRI-proton density fat fraction arises from reconstruction of noise. <i>Abdominal Radiology</i> , 2019, 44, 3295-3303.	1.0	7
300	Imaging Features at the Periphery: Hemodynamics, Pathophysiology, and Effect on LI-RADS Categorization. <i>Radiographics</i> , 2021, 41, 1657-1675.	1.4	7
301	Effect of intravenous gadoxetate disodium and flip angle on hepatic proton density fat fraction estimation with six-echo, gradient-recalled-echo, magnitude-based MR imaging at 3T. <i>Abdominal Radiology</i> , 2017, 42, 1189-1198.	1.0	6
302	Agreement between region-of-interest- and parametric map-based hepatic proton density fat fraction estimation in adults with chronic liver disease. <i>Abdominal Radiology</i> , 2017, 42, 833-841.	1.0	6
303	Convolutional neural network-automated hepatobiliary phase adequacy evaluation may optimize examination time. <i>European Journal of Radiology</i> , 2020, 124, 108837.	1.2	6
304	Abbreviated Magnetic Resonance Imaging for HCC Surveillance. <i>Clinical Liver Disease</i> , 2021, 17, 133-138.	1.0	6
305	Magnetic resonance elastography biomarkers for detection of histologic alterations in nonalcoholic fatty liver disease in the absence of fibrosis. <i>European Radiology</i> , 2021, 31, 8408-8419.	2.3	6
306	Image Annotation by Eye Tracking: Accuracy and Precision of Centerlines of Obstructed Small-Bowel Segments Placed Using Eye Trackers. <i>Journal of Digital Imaging</i> , 2019, 32, 855-864.	1.6	5

#	ARTICLE	IF	CITATIONS
307	Repeatability and accuracy of various region-of-interest sampling strategies for hepatic MRI proton density fat fraction quantification. <i>Abdominal Radiology</i> , 2021, 46, 3105-3116.	1.0	5
308	Imaging of hepatocellular carcinoma: a pilot international survey. <i>Abdominal Radiology</i> , 2021, 46, 205-215.	1.0	4
309	The Golden Hour: How to Spend Your Time and Money in Trauma Radiology. <i>Radiology</i> , 2004, 232, 622-624.	3.6	3
310	Vascular Contact With Soft Tissue. <i>Journal of Computer Assisted Tomography</i> , 2008, 32, 185-190.	0.5	3
311	Abdominal Pain in a Young Man with Oral Pigmentations. <i>Journal of Emergency Medicine</i> , 2016, 50, 335-336.	0.3	3
312	Editorial for "Interreader Agreement of Liver Imaging Reporting and Data System on MRI: A Systematic Review and Meta Analysis" <i>Journal of Magnetic Resonance Imaging</i> , 2020, 52, 805-806.	1.9	3
313	Transient Homogeneously Enhancing Hepatic Masses: Can Size Predict Benignity?. <i>American Journal of Roentgenology</i> , 2008, 190, 300-307.	1.0	2
314	Technical report: gadoxetate-disodium-enhanced 2D R2* mapping: a novel approach for assessing bile ducts in living donors. <i>Abdominal Radiology</i> , 2018, 43, 1656-1660.	1.0	2
315	Examining LI-RADS recommendations: should observation size only be measured on non-arterial phases?. <i>Abdominal Radiology</i> , 2020, 45, 3144-3154.	1.0	2
316	Prospective comparison of longitudinal change in hepatic proton density fat fraction (PDFF) estimated by magnitude-based MRI (MRI-M) and complex-based MRI (MRI-C). <i>European Radiology</i> , 2020, 30, 5120-5129.	2.3	2
317	Invited Commentary. <i>Radiographics</i> , 2012, 32, 1995-1998.	1.4	1
318	Assessing liver fat fraction by ARFI induced shear wave attenuation: A preliminary result. , 2013, , .		1
319	Magnetic Resonance Elastography versus Transient Elastography in detection of fibrosis in nonalcoholic fatty liver disease: A systematic review and meta-analysis of individual participant data. <i>Journal of Hepatology</i> , 2018, 68, S560-S561.	1.8	1
320	PS-186-Functional liver imaging score derived from gadoteric acid-enhanced MRI predicts outcomes in patients with advanced chronic liver disease. <i>Journal of Hepatology</i> , 2019, 70, e115.	1.8	1
321	Insight into hepatocellular carcinoma biology with gadoxetate disodium-enhanced MRI. <i>Hepatic Oncology</i> , 2014, 1, 95-105.	4.2	0
322	Letter to the editor response. <i>Abdominal Radiology</i> , 2018, 43, 239-239.	1.0	0
323	Responsiveness of controlled attenuation parameter (CAP) and its correlation with magnetic resonance imaging-proton density fat fraction (MRI-PDFF) in a multi-center clinical trial of subjects with nonalcoholic steatohepatitis (NASH). <i>Journal of Hepatology</i> , 2018, 68, S564.	1.8	0