

# Hiroaki Ikezaki

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/5591420/publications.pdf>

Version: 2024-02-01

76  
papers

1,347  
citations

361388

20  
h-index

414395

32  
g-index

80  
all docs

80  
docs citations

80  
times ranked

2379  
citing authors

#	ARTICLE	IF	CITATIONS
1	Sex-specific Relationship Between Stress Coping Strategies and All-cause Mortality: Japan Multi-Institutional Collaborative Cohort Study. <i>Journal of Epidemiology</i> , 2023, 33, 236-245.	2.4	2
2	Association of perceived stress and coping strategies with the renal function in middle-aged and older Japanese men and women. <i>Scientific Reports</i> , 2022, 12, 291.	3.3	1
3	The association of reproductive history with hypertension and obesity according to menopausal status: the J-MICC Study. <i>Hypertension Research</i> , 2022, 45, 708-714.	2.7	2
4	Effect of the interaction between physical activity and estimated macronutrient intake on HbA1c: population-based cross-sectional and longitudinal studies. <i>BMJ Open Diabetes Research and Care</i> , 2022, 10, e002479.	2.8	1
5	Associations of breastfeeding history with metabolic syndrome and cardiovascular risk factors in community-dwelling parous women: The Japan Multi-Institutional Collaborative Cohort Study. <i>PLoS ONE</i> , 2022, 17, e0262252.	2.5	5
6	A genome-wide association study on adherence to low-carbohydrate diets in Japanese. <i>European Journal of Clinical Nutrition</i> , 2022, , .	2.9	1
7	Dynamics of anti-Spike IgG antibody level after the second BNT162b2 COVID-19 vaccination in health care workers. <i>Journal of Infection and Chemotherapy</i> , 2022, 28, 802-805.	1.7	18
8	A case of severe COVID-19 with pulmonary thromboembolism related to heparin-induced thrombocytopenia during prophylactic anticoagulation therapy. <i>Journal of Infection and Chemotherapy</i> , 2022, , .	1.7	3
9	Impact of the PNPLA3 genotype on the risk of hepatocellular carcinoma after hepatitis C virus eradication. <i>Journal of Medical Virology</i> , 2022, 94, 5007-5014.	5.0	1
10	Associations of Genome-Wide Polygenic Risk Score and Risk Factors With Hypertension in a Japanese Population. <i>Circulation Genomic and Precision Medicine</i> , 2022, 15, .	3.6	6
11	Differential Effect of Polymorphisms on Body Mass Index Across the Life Course of Japanese: The Japan Multi-Institutional Collaborative Cohort Study. <i>Journal of Epidemiology</i> , 2021, 31, 172-179.	2.4	5
12	A genome-wide association study in Japanese identified one variant associated with a preference for a Japanese dietary pattern. <i>European Journal of Clinical Nutrition</i> , 2021, 75, 937-945.	2.9	8
13	A genome-wide association study on fish consumption in a Japanese population—the Japan Multi-Institutional Collaborative Cohort study. <i>European Journal of Clinical Nutrition</i> , 2021, 75, 480-488.	2.9	5
14	Impact of <i>PSCA</i> Polymorphisms on the Risk of Duodenal Ulcer. <i>Journal of Epidemiology</i> , 2021, 31, 12-20.	2.4	9
15	Population-Based Impact of Smoking, Drinking, and Genetic Factors on HDL-Cholesterol Levels in J-MICC Study Participants. <i>Journal of Epidemiology</i> , 2021, , .	2.4	0
16	Assessing the Relationship Between High-sensitivity C-reactive Protein and Kidney Function Employing Mendelian Randomization in the Japanese Community-based J-MICC Study. <i>Journal of Epidemiology</i> , 2021, , .	2.4	0
17	A genome-wide association study on confection consumption in a Japanese population: the Japan Multi-Institutional Collaborative Cohort Study. <i>British Journal of Nutrition</i> , 2021, 126, 1843-1851.	2.3	6
18	Reproducibility and validity of food group intake in a short food frequency questionnaire for the middle-aged Japanese population. <i>Environmental Health and Preventive Medicine</i> , 2021, 26, 28.	3.4	29

#	ARTICLE	IF	CITATIONS
19	Small Dense Low-Density Lipoprotein Cholesterol Is the Most Atherogenic Lipoprotein Parameter in the Prospective Framingham Offspring Study. <i>Journal of the American Heart Association</i> , 2021, 10, e019140.	3.7	88
20	Do neuroticism and extraversion personality traits influence disease-specific risk factors for mortality from cancer and cardiovascular disease in a Japanese population?. <i>Journal of Psychosomatic Research</i> , 2021, 144, 110422.	2.6	7
21	Effect of Underlying Cardiometabolic Diseases on the Association Between Sedentary Time and All-Cause Mortality in a Large Japanese Population: A Cohort Analysis Based on the J-MICC Study. <i>Journal of the American Heart Association</i> , 2021, 10, e018293.	3.7	9
22	Association of lifestyle factors with osteoporosis and fracture in postmenopausal women: a Japanese cohort study. <i>Menopause</i> , 2021, 28, 1254-1263.	2.0	3
23	Study Profile of the Japan Multi-institutional Collaborative Cohort (J-MICC) Study. <i>Journal of Epidemiology</i> , 2021, 31, 660-668.	2.4	41
24	A genome-wide association study on meat consumption in a Japanese population: the Japan Multi-Institutional Collaborative Cohort study. <i>Journal of Nutritional Science</i> , 2021, 10, e61.	1.9	3
25	Association of skipping breakfast and short sleep duration with the prevalence of metabolic syndrome in the general Japanese population: Baseline data from the Japan Multi-Institutional Collaborative cohort study. <i>Preventive Medicine Reports</i> , 2021, 24, 101613.	1.8	6
26	Genome-wide association study of serum prostate-specific antigen levels based on 1000 Genomes imputed data in Japanese: the Japan Multi-Institutional Collaborative Cohort Study. <i>Nagoya Journal of Medical Science</i> , 2021, 83, 183-194.	0.3	1
27	Kyushu and Okinawa Population Study (KOPS): a large prospective cohort study in Japan. <i>BMJ Open</i> , 2021, 11, e053763.	1.9	2
28	Combined effect of weight gain within normal weight range and parental hypertension on the prevalence of hypertension; from the J-MICC Study. <i>Journal of Human Hypertension</i> , 2020, 34, 125-131.	2.2	0
29	The interaction between ABCA1 polymorphism and physical activity on the HDL-cholesterol levels in a Japanese population. <i>Journal of Lipid Research</i> , 2020, 61, 86-94.	4.2	11
30	Population-specific and trans-ancestry genome-wide analyses identify distinct and shared genetic risk loci for coronary artery disease. <i>Nature Genetics</i> , 2020, 52, 1169-1177.	21.4	206
31	Association between plasma levels of homocysteine, folate, and vitamin B12, and dietary folate intake and hypertension in a cross-sectional study. <i>Scientific Reports</i> , 2020, 10, 18499.	3.3	14
32	Association between alcohol intake pattern and metabolic syndrome components and simulated change by alcohol intake reduction: A cross-sectional study from the Japan Multi-Institutional Collaborative Cohort Study. <i>Alcohol</i> , 2020, 89, 129-138.	1.7	1
33	Transethnic Meta-Analysis of Genome-Wide Association Studies Identifies Three New Loci and Characterizes Population-Specific Differences for Coronary Artery Disease. <i>Circulation Genomic and Precision Medicine</i> , 2020, 13, e002670.	3.6	44
34	Association of Dietary Acid Load with the Prevalence of Metabolic Syndrome among Participants in Baseline Survey of the Japan Multi-Institutional Collaborative Cohort Study. <i>Nutrients</i> , 2020, 12, 1605.	4.1	23
35	Identification of a novel uterine leiomyoma GWAS locus in a Japanese population. <i>Scientific Reports</i> , 2020, 10, 1197.	3.3	14
36	Small Dense Low-Density Lipoprotein Cholesterol and Carotid Intimal Medial Thickness Progression. <i>Journal of Atherosclerosis and Thrombosis</i> , 2020, 27, 1108-1122.	2.0	13

#	ARTICLE	IF	CITATIONS
37	Sedentary Time is Associated with Cardiometabolic Diseases in A Large Japanese Population: A Cross-Sectional Study. <i>Journal of Atherosclerosis and Thrombosis</i> , 2020, 27, 1097-1107.	2.0	14
38	Objective tongue color and gastroesophageal reflux disease: Cross-sectional study. <i>Traditional &amp; Kampo Medicine</i> , 2019, 6, 19-25.	0.6	2
39	Direct Versus Calculated LDL Cholesterol and C-Reactive Protein in Cardiovascular Disease Risk Assessment in the Framingham Offspring Study. <i>Clinical Chemistry</i> , 2019, 65, 1102-1114.	3.2	18
40	Associations of Nutrient Patterns with the Prevalence of Metabolic Syndrome: Results from the Baseline Data of the Japan Multi-Institutional Collaborative Cohort Study. <i>Nutrients</i> , 2019, 11, 990.	4.1	24
41	Independent relationships of daily life activity and leisure-time exercise with metabolic syndrome and its traits in the general Japanese population. <i>Endocrine</i> , 2019, 64, 552-563.	2.3	8
42	Genome-wide meta-analysis identifies multiple novel loci associated with serum uric acid levels in Japanese individuals. <i>Communications Biology</i> , 2019, 2, 115.	4.4	66
43	A genome-wide association study in the Japanese population identifies the 12q24 locus for habitual coffee consumption: The J-MICC Study. <i>Scientific Reports</i> , 2018, 8, 1493.	3.3	32
44	Association of exposure level to passive smoking with hypertension among lifetime nonsmokers in Japan: a cross-sectional study. <i>Medicine (United States)</i> , 2018, 97, e13241.	1.0	14
45	Correlation between Thyroid Stimulating Hormone and Renal Function in Euthyroid Residents of Japan: Results from the Kyushu and Okinawa Population Study (KOPS). <i>Journal of Atherosclerosis and Thrombosis</i> , 2018, 25, 335-343.	2.0	13
46	Development and Population Results of a Fully Automated Homogeneous Assay for LDL Triglyceride. <i>Journal of Applied Laboratory Medicine</i> , 2018, 2, 746-756.	1.3	24
47	Genome-Wide Association Study of Renal Function Traits: Results from the Japan Multi-Institutional Collaborative Cohort Study. <i>American Journal of Nephrology</i> , 2018, 47, 304-316.	3.1	18
48	Association of the Serum Endostatin Level, Renal Function, and Carotid Atherosclerosis of Healthy Residents of Japan: Results from the Kyushu and Okinawa Population Study (KOPS). <i>Journal of Atherosclerosis and Thrombosis</i> , 2018, 25, 829-835.	2.0	3
49	Genomewide Association Study of Leisure-Time Exercise Behavior in Japanese Adults. <i>Medicine and Science in Sports and Exercise</i> , 2018, 50, 2433-2441.	0.4	36
50	Cardiovascular disease prevalence and insulin resistance in the Kyushu-Okinawa Population Study and the Framingham Offspring Study. <i>Journal of Clinical Lipidology</i> , 2017, 11, 348-356.	1.5	12
51	Higher dietary cholesterol and $\omega$ -3 fatty acid intakes are associated with a lower success rate of <i>Helicobacter pylori</i> eradication therapy in Japan. <i>American Journal of Clinical Nutrition</i> , 2017, 106, 581-588.	4.7	23
52	Genetic Variants of <i>RAMP2</i> and <i>CLR</i> are Associated with Stroke. <i>Journal of Atherosclerosis and Thrombosis</i> , 2017, 24, 1267-1281.	2.0	11
53	Impact of Food Consumption on <i>Helicobacter pylori</i> Eradication Therapy: A Japanese Community-Based Observational Study. <i>Open Forum Infectious Diseases</i> , 2016, 3, .	0.9	0
54	Efficacy of interferon- $\beta$ plus ribavirin combination treatment on the development of hepatocellular carcinoma in Japanese patients with chronic hepatitis C. <i>Hepatology Research</i> , 2016, 46, E174-80.	3.4	4

#	ARTICLE	IF	CITATIONS
55	Ethnic Differences in Glucose Homeostasis Markers between the Kyushu-Okinawa Population Study and the Framingham Offspring Study. <i>Scientific Reports</i> , 2016, 6, 36725.	3.3	9
56	The relation of postprandial plasma glucose and serum endostatin to the urinary albumin excretion of residents with prediabetes: results from the Kyushu and Okinawa Population Study (KOPS). <i>International Urology and Nephrology</i> , 2016, 48, 851-857.	1.4	3
57	Association of IL28B rs8099917 genotype and female sex with spontaneous clearance of hepatitis C virus infection: a Japanese cross-sectional study. <i>Archives of Virology</i> , 2016, 161, 641-648.	2.1	11
58	Glycated albumin as a diagnostic tool for diabetes in a general Japanese population. <i>Metabolism: Clinical and Experimental</i> , 2015, 64, 698-705.	3.4	42
59	Subclinical carotid atherosclerosis and triglycerides predict the incidence of chronic kidney disease in the Japanese general population: Results from the Kyushu and Okinawa Population Study (KOPS). <i>Atherosclerosis</i> , 2015, 238, 207-212.	0.8	33
60	Direct-acting antiviral-based triple therapy on alpha-fetoprotein level in chronic hepatitis C patients. <i>World Journal of Gastroenterology</i> , 2015, 21, 4696-4706.	3.3	4
61	Predictors of kidney tubular dysfunction induced by adefovir treatment for chronic hepatitis B. <i>World Journal of Gastroenterology</i> , 2015, 21, 2116-2123.	3.3	25
62	The utility of urinary myo-inositol as a marker of glucose intolerance. <i>Diabetes Research and Clinical Practice</i> , 2014, 103, 88-96.	2.8	14
63	A case of successful hepatitis C virus eradication by 24 weeks of telaprevir-based triple therapy for a hemophilia patient with hepatitis C virus/human immunodeficiency virus co-infection who previously failed pegylated interferon- $\alpha$ and ribavirin therapy. <i>Journal of Infection and Chemotherapy</i> , 2014, 20, 320-324.	1.7	4
64	Hepatitis B Virus-related Immune Reconstitution Inflammatory Syndrome in Two Patients Coinfected with Human Immunodeficiency Virus Diagnosed with a Liver Biopsy. <i>Internal Medicine</i> , 2014, 53, 2165-2170.	0.7	14
65	Efficacy and Tolerance of Interferon $\hat{I}^2$ Plus Ribavirin Treatment for Chronic Hepatitis C Patients with Depression or Thrombocytopenia Comparison with Pegylated Interferon $\hat{I}^1$ Plus Ribavirin Treatment. <i>Journal of Liver</i> , 2014, 03, .	0.3	1
66	The serum undercarboxylated osteocalcin level and the diet of a Japanese population: results from the Kyushu and Okinawa Population Study (KOPS). <i>Endocrine</i> , 2013, 43, 635-642.	2.3	14
67	Plasma glycated albumin level and atherosclerosis: Results from the Kyushu and Okinawa Population Study (KOPS). <i>International Journal of Cardiology</i> , 2013, 167, 2066-2072.	1.7	49
68	Serum cholesterol and triglyceride reference ranges of twenty lipoprotein subclasses for healthy Japanese men and women. <i>Atherosclerosis</i> , 2013, 231, 238-245.	0.8	28
69	Association between chronic hepatitis C virus infection and high levels of circulating N-terminal pro-brain natriuretic peptide. <i>Endocrine</i> , 2013, 43, 200-205.	2.3	3
70	Valuable antiviral therapeutic options for the treatment of chronic hepatitis C patients with thrombocytopenia. <i>Journal of Viral Hepatitis</i> , 2013, 20, 838-846.	2.0	2
71	Raloxifene hydrochloride is an adjuvant antiviral treatment of postmenopausal women with chronic hepatitis C: A randomized trial. <i>Journal of Hepatology</i> , 2012, 57, 1186-1192.	3.7	52
72	Insulin resistance undermines the advantages of IL28B polymorphism in the pegylated interferon alpha-2b and ribavirin treatment of chronic hepatitis C patients with genotype 1. <i>Journal of Hepatology</i> , 2012, 57, 534-540.	3.7	37

#	ARTICLE	IF	CITATIONS
73	Longitudinal assessment of liver stiffness by transient elastography for chronic hepatitis B patients treated with nucleoside analog. <i>Hepatology Research</i> , 2011, 41, 1178-1188.	3.4	55
74	Abbott RealTime PCR assay is useful for evaluating virological response to antiviral treatment for chronic hepatitis C. <i>Journal of Infection and Chemotherapy</i> , 2011, 17, 737-743.	1.7	4
75	Ribavirin concentration in the later stages of 48 week pegylated interferon- $\alpha$ 2b plus ribavirin therapy for chronic hepatitis C is useful for predicting virological response. <i>Journal of Antimicrobial Chemotherapy</i> , 2011, 66, 1127-1139.	3.0	20
76	Treatment for Eradication of <i>Helicobacter pylori</i> Infection among Chronic Hepatitis C Patients. <i>Gut and Liver</i> , 2011, 5, 447-453.	2.9	8