

Valentina Sabino

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/5592641/publications.pdf>

Version: 2024-02-01

77
papers

3,307
citations

117625

34
h-index

155660

55
g-index

77
all docs

77
docs citations

77
times ranked

2945
citing authors

#	ARTICLE	IF	CITATIONS
1	Pituitary adenylate cyclase-activating polypeptide type 1 receptor within the nucleus accumbens core mediates excessive alcohol drinking in alcohol-preferring rats. <i>Neuropharmacology</i> , 2022, 212, 109063.	4.1	3
2	Pituitary adenylate cyclase-activating polypeptide (PACAP) modulates dependence-induced alcohol drinking and anxiety-like behavior in male rats. <i>Neuropsychopharmacology</i> , 2021, 46, 509-518.	5.4	23
3	The Sigma-2 receptor / transmembrane protein 97 ($\sigma_2R/TMEM97$) modulator JW-1034 reduces heavy alcohol drinking and associated pain states in male mice. <i>Neuropharmacology</i> , 2021, 184, 108409.	4.1	27
4	Antagonism of σ_1 receptor blocks heavy alcohol drinking and associated hyperalgesia in male mice. <i>Alcoholism: Clinical and Experimental Research</i> , 2021, 45, 1398-1407.	2.4	10
5	Viral-Mediated Knockdown of Nucleus Accumbens Shell PAC1 Receptor Promotes Excessive Alcohol Drinking in Alcohol-Preferring Rats. <i>Frontiers in Behavioral Neuroscience</i> , 2021, 15, 787362.	2.0	5
6	PACAP regulation of central amygdala GABAergic synapses is altered by restraint stress. <i>Neuropharmacology</i> , 2020, 168, 107752.	4.1	26
7	Reward sensitivity deficits in a rat model of compulsive eating behavior. <i>Neuropsychopharmacology</i> , 2020, 45, 589-596.	5.4	17
8	Effect of different standard rodent diets on ethanol intake and associated allodynia in male mice. <i>Alcohol</i> , 2020, 87, 17-23.	1.7	15
9	Sigma receptor-induced heavy drinking in rats: Modulation by the opioid receptor system. <i>Pharmacology Biochemistry and Behavior</i> , 2020, 192, 172914.	2.9	5
10	Withdrawal from Extended, Intermittent Access to A Highly Palatable Diet Impairs Hippocampal Memory Function and Neurogenesis: Effects of Memantine. <i>Nutrients</i> , 2020, 12, 1520.	4.1	6
11	Consummatory, Feeding Microstructural, and Metabolic Effects Induced by Limiting Access to Either a High-Sucrose or a High-Fat Diet. <i>Nutrients</i> , 2020, 12, 1610.	4.1	2
12	The Alpha-1 Adrenergic Receptor Antagonist Prazosin Reduces Binge-Like Eating in Rats. <i>Nutrients</i> , 2020, 12, 1569.	4.1	8
13	Dissecting compulsive eating behavior into three elements. , 2019, , 41-81.		4
14	Habitual overeating. , 2019, , 83-95.		1
15	Modeling and testing compulsive eating behaviors in animals. , 2019, , 359-388.		2
16	Role of Sigma Receptors in Alcohol Addiction. <i>Frontiers in Pharmacology</i> , 2019, 10, 687.	3.5	19
17	Role of the PACAP system of the extended amygdala in the acoustic startle response in rats. <i>Neuropharmacology</i> , 2019, 160, 107761.	4.1	19
18	Opposing roles of σ_1 and σ_2 receptors in heavy alcohol drinking and associated mechanical allodynia in mice. <i>FASEB Journal</i> , 2019, 33, 499.8.	0.5	0

#	ARTICLE	IF	CITATIONS
19	Reward deficits in an animal model of compulsive eating. <i>FASEB Journal</i> , 2019, 33, 805.3.	0.5	0
20	Neuropharmacology of compulsive eating. <i>Philosophical Transactions of the Royal Society B: Biological Sciences</i> , 2018, 373, 20170024.	4.0	20
21	Small molecule modulators of Trf2R/Tmem97 reduce alcohol withdrawal-induced behaviors. <i>Neuropsychopharmacology</i> , 2018, 43, 1867-1875.	5.4	35
22	Impulsive choice does not predict binge-like eating in rats. <i>Behavioural Pharmacology</i> , 2018, 29, 726-731.	1.7	5
23	Trace Amine Associated Receptor 1 (TAAR1) Modulation of Food Reward. <i>Frontiers in Pharmacology</i> , 2018, 9, 129.	3.5	19
24	Evaluation of Alcohol Preference and Drinking in msP Rats Bearing a <i>Crhr1</i> Promoter Polymorphism. <i>Frontiers in Psychiatry</i> , 2018, 9, 28.	2.6	10
25	Sigma Receptors and Substance Use Disorders. <i>Advances in Experimental Medicine and Biology</i> , 2017, 964, 177-199.	1.6	15
26	Pathological Overeating: Emerging Evidence for a Compulsivity Construct. <i>Neuropsychopharmacology</i> , 2017, 42, 1375-1389.	5.4	92
27	A behavioral and pharmacological characterization of palatable diet alternation in mice. <i>Pharmacology Biochemistry and Behavior</i> , 2017, 163, 1-8.	2.9	14
28	The Trace Amine-Associated Receptor 1 Agonist RO5256390 Blocks Compulsive, Binge-like Eating in Rats. <i>Neuropsychopharmacology</i> , 2017, 42, 1458-1470.	5.4	63
29	Neuroscience of Compulsive Eating Behavior. <i>Frontiers in Neuroscience</i> , 2017, 11, 469.	2.8	41
30	Pituitary adenylate cyclase-activating polypeptide (PACAP) in the central nucleus of the amygdala induces anxiety via melanocortin receptors. <i>Psychopharmacology</i> , 2016, 233, 3269-3277.	3.1	37
31	Sigma Receptors and Alcohol Use Disorders. <i>Handbook of Experimental Pharmacology</i> , 2016, 244, 219-236.	1.8	7
32	Ethanol-related behaviors in mice lacking the sigma-1 receptor. <i>Behavioural Brain Research</i> , 2016, 297, 196-203.	2.2	13
33	Pituitary Adenylate Cyclase-Activating Peptide in the Central Amygdala Causes Anorexia and Body Weight Loss via the Melanocortin and the TrkB Systems. <i>Neuropsychopharmacology</i> , 2015, 40, 1846-1855.	5.4	32
34	Seeking behavior, place conditioning, and resistance to conditioned suppression of feeding in rats intermittently exposed to palatable food.. <i>Behavioral Neuroscience</i> , 2015, 129, 219-224.	1.2	35
35	Sigma-1 receptor mediates acquisition of alcohol drinking and seeking behavior in alcohol-preferring rats. <i>Behavioural Brain Research</i> , 2015, 287, 315-322.	2.2	29
36	Diet-induced obesity and diet-resistant rats: differences in the rewarding and anorectic effects of d-amphetamine. <i>Psychopharmacology</i> , 2015, 232, 3215-3226.	3.1	24

#	ARTICLE	IF	CITATIONS
37	Pituitary adenylate cyclase-activating polypeptide induces a depressive-like phenotype in rats. <i>Psychopharmacology</i> , 2015, 232, 3821-3831.	3.1	25
38	The Uncompetitive N-methyl-D-Aspartate Antagonist Memantine Reduces Binge-Like Eating, Food-Seeking Behavior, and Compulsive Eating: Role of the Nucleus Accumbens Shell. <i>Neuropsychopharmacology</i> , 2015, 40, 1163-1171.	5.4	47
39	High Trait Impulsivity Predicts Food Addiction-Like Behavior in the Rat. <i>FASEB Journal</i> , 2015, 29, 769.5.	0.5	0
40	Characterization of a shortened model of diet alternation in female rats. <i>Behavioural Pharmacology</i> , 2014, 25, 609-617.	1.7	22
41	High Trait Impulsivity Predicts Food Addiction-Like Behavior in the Rat. <i>Neuropsychopharmacology</i> , 2014, 39, 2463-2472.	5.4	116
42	Opioid system in the medial prefrontal cortex mediates binge-like eating. <i>Addiction Biology</i> , 2014, 19, 652-662.	2.6	77
43	Nicotine dependence produces hyperalgesia: Role of corticotropin-releasing factor-1 receptors (CRF1Rs) in the central amygdala (CeA). <i>Neuropharmacology</i> , 2014, 77, 217-223.	4.1	51
44	The inverse agonist of CB_1 receptor $SR141716$ blocks compulsive eating of palatable food. <i>Addiction Biology</i> , 2014, 19, 849-861.	2.6	42
45	The uncompetitive NMDA receptor antagonists ketamine and memantine preferentially increase the choice for a small, immediate reward in low-impulsive rats. <i>Psychopharmacology</i> , 2013, 226, 127-138.	3.1	42
46	Rimonabant Precipitates Anxiety in Rats Withdrawn from Palatable Food: Role of the Central Amygdala. <i>Neuropsychopharmacology</i> , 2013, 38, 2498-2507.	5.4	54
47	CRF-CRF1 Receptor System in the Central and Basolateral Nuclei of the Amygdala Differentially Mediates Excessive Eating of Palatable Food. <i>Neuropsychopharmacology</i> , 2013, 38, 2456-2466.	5.4	71
48	mTOR activation is required for the anti-alcohol effect of ketamine, but not memantine, in alcohol-preferring rats. <i>Behavioural Brain Research</i> , 2013, 247, 9-16.	2.2	51
49	Pharmacological Characterization of the 20% Alcohol Intermittent Access Model in S -ardinian Alcohol-Preferring Rats: A Model of Binge-Like Drinking. <i>Alcoholism: Clinical and Experimental Research</i> , 2013, 37, 635-643.	2.4	45
50	CRF Mediates the Anxiogenic and Anti-Rewarding, But Not the Anorectic Effects of PACAP. <i>Neuropsychopharmacology</i> , 2013, 38, 2160-2169.	5.4	68
51	Centrally administered urocortin 2 decreases gorging on high-fat diet in both diet-induced obesity-prone and -resistant rats. <i>International Journal of Obesity</i> , 2013, 37, 1515-1523.	3.4	13
52	Antagonism of Sigma-1 Receptors Blocks Compulsive-Like Eating. <i>Neuropsychopharmacology</i> , 2012, 37, 2593-2604.	5.4	72
53	Withdrawal from chronic, intermittent access to a highly palatable food induces depressive-like behavior in compulsive eating rats. <i>Behavioural Pharmacology</i> , 2012, 23, 593-602.	1.7	69
54	Effects of CB_1 and CRF1 receptor antagonists on binge-like eating in rats with limited access to a sweet fat diet: Lack of withdrawal-like responses. <i>Physiology and Behavior</i> , 2012, 107, 231-242.	2.1	60

#	ARTICLE	IF	CITATIONS
55	A modified adjusting delay task to assess impulsive choice between isocaloric reinforcers in non-deprived male rats: effects of 5-HT _{2A/C} and 5-HT _{1A} receptor agonists. <i>Psychopharmacology</i> , 2012, 219, 377-386.	3.1	22
56	Systemic urocortin 2, but not urocortin 1 or stressin1-A, suppresses feeding via CRF2 receptors without malaise and stress. <i>British Journal of Pharmacology</i> , 2011, 164, 1959-1975.	5.4	35
57	Activation of Υ -Receptors Induces Binge-like Drinking in Sardinian Alcohol-Preferring Rats. <i>Neuropsychopharmacology</i> , 2011, 36, 1207-1218.	5.4	53
58	Genome-wide gene expression analysis identifies K-ras as a regulator of alcohol intake. <i>Brain Research</i> , 2010, 1339, 1-10.	2.2	25
59	Corticotropin Releasing Factor-Induced Amygdala Gamma-Aminobutyric Acid Release Plays a Key Role in Alcohol Dependence. <i>Biological Psychiatry</i> , 2010, 67, 831-839.	1.3	303
60	The Υ -Receptor Antagonist BD-1063 Decreases Ethanol Intake and Reinforcement in Animal Models of Excessive Drinking. <i>Neuropsychopharmacology</i> , 2009, 34, 1482-1493.	5.4	69
61	Consummatory, anxiety-related and metabolic adaptations in female rats with alternating access to preferred food. <i>Psychoneuroendocrinology</i> , 2009, 34, 38-49.	2.7	92
62	Selective reduction of alcohol drinking in Sardinian alcohol-preferring rats by a sigma-1 receptor antagonist. <i>Psychopharmacology</i> , 2009, 205, 327-335.	3.1	38
63	Increased Periculomotor Urocortin 1 Immunoreactivity in Genetically Selected Alcohol Preferring Rats. <i>Alcoholism: Clinical and Experimental Research</i> , 2009, 33, 1956-1965.	2.4	29
64	Sigma-1 receptor knockout mice display a depressive-like phenotype. <i>Behavioural Brain Research</i> , 2009, 198, 472-476.	2.2	146
65	Social defeat stress activates medial amygdala cells that express type 2 corticotropin-releasing factor receptor mRNA. <i>Neuroscience</i> , 2009, 162, 5-13.	2.3	61
66	CRF system recruitment mediates dark side of compulsive eating. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2009, 106, 20016-20020.	7.1	168
67	Intermittent access to preferred food reduces the reinforcing efficacy of chow in rats. <i>American Journal of Physiology - Regulatory Integrative and Comparative Physiology</i> , 2008, 295, R1066-R1076.	1.8	70
68	Opioid-Dependent Anticipatory Negative Contrast and Binge-Like Eating in Rats with Limited Access to Highly Preferred Food. <i>Neuropsychopharmacology</i> , 2008, 33, 524-535.	5.4	130
69	FG 7142 Specifically Reduces Meal Size and the Rate and Regularity of Sustained Feeding in Female Rats: Evidence that Benzodiazepine Inverse Agonists Reduce Food Palatability. <i>Neuropsychopharmacology</i> , 2007, 32, 1069-1081.	5.4	26
70	Galanin type 1 receptor knockout mice show altered responses to high-fat diet and glucose challenge. <i>Physiology and Behavior</i> , 2007, 91, 479-485.	2.1	68
71	Feeding microstructure in diet-induced obesity susceptible versus resistant rats: central effects of urocortin 2. <i>Journal of Physiology</i> , 2007, 583, 487-504.	2.9	44
72	14-methoxymetopon, a highly potent Υ ₄ opioid agonist, biphasically affects ethanol intake in Sardinian alcohol-preferring rats. <i>Psychopharmacology</i> , 2007, 192, 537-546.	3.1	26

#	ARTICLE	IF	CITATIONS
73	Dissociation between opioid and CRF1 antagonist sensitive drinking in Sardinian alcohol-preferring rats. <i>Psychopharmacology</i> , 2006, 189, 175-186.	3.1	101
74	Increased Anxiety-Like Behavior and Ethanol Self-Administration in Dependent Rats: Reversal via Corticotropin-Releasing Factor-2 Receptor Activation. <i>Alcoholism: Clinical and Experimental Research</i> , 2004, 28, 865-872.	2.4	131
75	Long-term effects on cortical glutamate release induced by prenatal exposure to the cannabinoid receptor agonist (r)-(+)-[2,3-dihydro-5-methyl-3-(4-morpholinyl-methyl)pyrrolo[1,2,3-de]-1,4-benzoxazin-6-yl]-1-naphthalenylmethanone: an in vivo microdialysis study in the awake rat. <i>Neuroscience</i> , 2004, 124, 367-375.	2.3	43
76	Functional characterization of β_1 -adrenoceptor subtypes in vascular tissues using different experimental approaches: a comparative study. <i>British Journal of Pharmacology</i> , 2003, 138, 359-368.	5.4	18
77	Pituitary adenylate cyclase-activating polypeptide (PACAP) modulates dependence-induced alcohol drinking and anxiety-like behavior in male rats. , 0, .		1