

Ming-shunchen Chen

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/5631905/publications.pdf>

Version: 2024-02-01

68
papers

3,080
citations

218677

26
h-index

161849

54
g-index

69
all docs

69
docs citations

69
times ranked

2797
citing authors

#	ARTICLE	IF	CITATIONS
1	Identification of a major QTL for Hessian fly resistance in wheat cultivar "Chokwang". <i>Crop Journal</i> , 2022, 10, 775-782.	5.2	10
2	A new strategy for using historical imbalanced yield data to conduct genome-wide association studies and develop genomic prediction models for wheat breeding. <i>Molecular Breeding</i> , 2022, 42, 1.	2.1	0
3	Chromosome-level genome assembly for the horned gall aphid provides insights into interactions between gall-making insect and its host plant. <i>Ecology and Evolution</i> , 2022, 12, e8815.	1.9	8
4	"Gallagher" and "Iba" hard red winter wheat: Half-sibs inseparable by yield gain, separable by producer preference. <i>Journal of Plant Registrations</i> , 2021, 15, 177-195.	0.5	3
5	Proteomic and transcriptomic analyses of saliva and salivary glands from the Asian citrus psyllid, <i>Diaphorina citri</i> . <i>Journal of Proteomics</i> , 2021, 238, 104136.	2.4	16
6	Gall-Inducing Parasites: Convergent and Conserved Strategies of Plant Manipulation by Insects and Nematodes. <i>Annual Review of Phytopathology</i> , 2020, 58, 1-22.	7.8	37
7	Differential localization of Hessian fly candidate effectors in resistant and susceptible wheat plants. <i>Plant Direct</i> , 2020, 4, e00246.	1.9	9
8	Analyzing Molecular Basis of Heat-Induced Loss-of-Wheat Resistance to Hessian Fly (Diptera: Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50 467 1504-1512.	1.8	1
9	Identification of two novel Hessian fly resistance genes H35 and H36 in a hard winter wheat line SD06165. <i>Theoretical and Applied Genetics</i> , 2020, 133, 2343-2353.	3.6	31
10	The Hessian fly recessive resistance gene h4 mapped to chromosome 1A of the wheat cultivar "Java" using genotyping-by-sequencing. <i>Theoretical and Applied Genetics</i> , 2020, 133, 2927-2935.	3.6	7
11	A Horizontal Gene Transfer Led to the Acquisition of a Fructan Metabolic Pathway in a Gall Midge. <i>Advanced Biology</i> , 2020, 4, 1900275.	3.0	4
12	Cytokinins Are Abundant and Widespread among Insect Species. <i>Plants</i> , 2020, 9, 208.	3.5	31
13	Potential Pathways and Genes Involved in Lac Synthesis and Secretion in <i>Kerria chinensis</i> (Hemiptera: Tj ETQq1 1 0,784314 rgBT /Overlock 10 Tf 50 467 1504-1512.	2.2	5
14	Transcriptomic Analyses of Secreted Proteins From the Salivary Glands of Wheat Midge Larvae. <i>Journal of Insect Science</i> , 2018, 18, .	1.5	10
15	Increasing Temperature Reduces Wheat Resistance Mediated by Major Resistance Genes to <i>Mayetiola destructor</i> (Diptera: Cecidomyiidae). <i>Journal of Economic Entomology</i> , 2018, 111, 1433-1438.	1.8	11
16	Indirect plant defense against insect herbivores: a review. <i>Insect Science</i> , 2018, 25, 2-23.	3.0	225
17	Conserved and Unique Putative Effectors Expressed in the Salivary Glands of Three Related Gall Midge Species. <i>Journal of Insect Science</i> , 2018, 18, .	1.5	6
18	An insect nucleoside diphosphate kinase (NDK) functions as an effector protein in wheat - Hessian fly interactions. <i>Insect Biochemistry and Molecular Biology</i> , 2018, 100, 30-38.	2.7	17

#	ARTICLE	IF	CITATIONS
19	Genes Expressed Differentially in Hessian Fly Larvae Feeding in Resistant and Susceptible Plants. <i>International Journal of Molecular Sciences</i> , 2016, 17, 1324.	4.1	7
20	Massive Shift in Gene Expression during Transitions between Developmental Stages of the Gall Midge, <i>Mayetiola Destructor</i> . <i>PLoS ONE</i> , 2016, 11, e0155616.	2.5	8
21	Impact of Hessian fly, <i>Mayetiola destructor</i> , on Developmental Aspects of Hard Red Winter Wheat in Kansas. <i>Southwestern Entomologist</i> , 2016, 41, 321-330.	0.2	10
22	A Massive Expansion of Effector Genes Underlies Gall-Formation in the Wheat Pest <i>Mayetiola destructor</i> . <i>Current Biology</i> , 2015, 25, 613-620.	3.9	171
23	Unbalanced Activation of Glutathione Metabolic Pathways Suggests Potential Involvement in Plant Defense against the Gall Midge <i>Mayetiola destructor</i> in Wheat. <i>Scientific Reports</i> , 2015, 5, 8092.	3.3	38
24	Precisely mapping a major gene conferring resistance to Hessian fly in bread wheat using genotyping-by-sequencing. <i>BMC Genomics</i> , 2015, 16, 108.	2.8	36
25	Avirulence Effector Discovery in a Plant Galling and Plant Parasitic Arthropod, the Hessian Fly (<i>Mayetiola destructor</i>). <i>PLoS ONE</i> , 2014, 9, e100958.	2.5	54
26	Pyrosequencing Reveals the Predominance of Pseudomonadaceae in Gut Microbiome of a Gall Midge. <i>Pathogens</i> , 2014, 3, 459-472.	2.8	21
27	Molecular Markers for Species Identification of Hessian Fly Males Caught on Sticky Pheromone Traps. <i>Journal of Economic Entomology</i> , 2014, 107, 1110-1117.	1.8	8
28	Virulence and Biotype Analyses of Hessian Fly (Diptera: Cecidomyiidae) Populations From Texas, Louisiana, and Oklahoma. <i>Journal of Economic Entomology</i> , 2014, 107, 417-423.	1.8	22
29	Exogenous Salicylic Acid Enhances the Resistance of Wheat Seedlings to Hessian Fly (Diptera: Tj ETQq1 1 0.784314,rgBT /Overlock 10	1.8	8
30	Comparative gut transcriptome analysis reveals differences between virulent and avirulent Russian wheat aphids, <i>Diuraphis noxia</i> . <i>Arthropod-Plant Interactions</i> , 2014, 8, 79-88.	1.1	19
31	Wheat Mds-1 encodes a heat-shock protein and governs susceptibility towards the Hessian fly gall midge. <i>Nature Communications</i> , 2013, 4, 2070.	12.8	33
32	Mobilization of lipids and fortification of cell wall and cuticle are important in host defense against Hessian fly. <i>BMC Genomics</i> , 2013, 14, 423.	2.8	26
33	Genetic association of OPRgenes with resistance to Hessian fly in hexaploid wheat. <i>BMC Genomics</i> , 2013, 14, 369.	2.8	19
34	Deep sequencing and genome-wide analysis reveals the expansion of MicroRNA genes in the gall midge <i>Mayetiola destructor</i> . <i>BMC Genomics</i> , 2013, 14, 187.	2.8	17
35	Serine and cysteine protease-like genes in the genome of a gall midge and their interactions with host plant genotypes. <i>Insect Biochemistry and Molecular Biology</i> , 2013, 43, 701-711.	2.7	17
36	Gall Midges (Hessian Flies) as Plant Pathogens. <i>Annual Review of Phytopathology</i> , 2012, 50, 339-357.	7.8	130

#	ARTICLE	IF	CITATIONS
37	Rapid Mobilization of Membrane Lipids in Wheat Leaf Sheaths During Incompatible Interactions with Hessian Fly. <i>Molecular Plant-Microbe Interactions</i> , 2012, 25, 920-930.	2.6	25
38	Changes in Phytohormones and Fatty Acids in Wheat and Rice Seedlings in Response to Hessian Fly (Diptera: Cecidomyiidae) Infestation. <i>Journal of Economic Entomology</i> , 2011, 104, 1384-1392.	1.8	19
39	Hessian Fly-Associated Bacteria: Transmission, Essentiality, and Composition. <i>PLoS ONE</i> , 2011, 6, e23170.	2.5	55
40	Aphid Feeding Activates Expression of a Transcriptome of Oxylipin-based Defense Signals in Wheat Involved in Resistance to Herbivory. <i>Journal of Chemical Ecology</i> , 2010, 36, 260-276.	1.8	86
41	The gut transcriptome of a gall midge, <i>Mayetiola destructor</i> . <i>Journal of Insect Physiology</i> , 2010, 56, 1198-1206.	2.0	26
42	Unusual conservation among genes encoding small secreted salivary gland proteins from a gall midge. <i>BMC Evolutionary Biology</i> , 2010, 10, 296.	3.2	55
43	A Neo-Sex Chromosome That Drives Postzygotic Sex Determination in the Hessian Fly (<i>Mayetiola</i>) Tj ETQq1 1 0.784314 rgBT /Overlock 10	2.9	32
44	Differential Accumulation of Phytohormones in Wheat Seedlings Attacked by Avirulent and Virulent Hessian Fly (Diptera: Cecidomyiidae) Larvae. <i>Journal of Economic Entomology</i> , 2010, 103, 178-185.	1.8	17
45	Reactive Oxygen Species Are Involved in Plant Defense against a Gall Midge <i>Mayetiola destructor</i> . <i>Plant Physiology</i> , 2010, 152, 985-999.	4.8	161
46	Virulence Analysis of Hessian Fly Populations From Texas, Oklahoma, and Kansas. <i>Journal of Economic Entomology</i> , 2009, 102, 774-780.	1.8	97
47	Hessian Fly (Diptera: Cecidomyiidae) Interactions With Barley, Rice, and Wheat Seedlings. <i>Journal of Economic Entomology</i> , 2009, 102, 1663-1672.	1.8	21
48	Characterization and expression analysis of a gene encoding a secreted lipase-like protein expressed in the salivary glands of the larval Hessian fly, <i>Mayetiola destructor</i> (Say). <i>Journal of Insect Physiology</i> , 2009, 55, 105-112.	2.0	23
49	Expressed sequence tags from larval gut of the European corn borer (<i>Ostrinia nubilalis</i>): Exploring candidate genes potentially involved in <i>Bacillus thuringiensis</i> toxicity and resistance. <i>BMC Genomics</i> , 2009, 10, 286.	2.8	42
50	A BAC-based physical map of the Hessian fly genome anchored to polytene chromosomes. <i>BMC Genomics</i> , 2009, 10, 293.	2.8	20
51	Differential Responses of Wheat Inhibitor-like Genes to Hessian Fly, <i>Mayetiola destructor</i> , Attacks During Compatible and Incompatible Interactions. <i>Journal of Chemical Ecology</i> , 2008, 34, 1005-1012.	1.8	22
52	Inducible direct plant defense against insect herbivores: A review. <i>Insect Science</i> , 2008, 15, 101-114.	3.0	327
53	Analysis of transcripts and proteins expressed in the salivary glands of Hessian fly (<i>Mayetiola</i>) Tj ETQq1 1 0.784314 rgBT /Overlock 10	2.8	61
54	Hessian Fly. , 2008, , 93-102.		13

#	ARTICLE	IF	CITATIONS
55	A protein from the salivary glands of the pea aphid, <i>Acyrtosiphon pisum</i> , is essential in feeding on a host plant. Proceedings of the National Academy of Sciences of the United States of America, 2008, 105, 9965-9969.	7.1	339
56	Hessian Fly (<i>Mayetiola destructor</i>) Attack Causes a Dramatic Shift in Carbon and Nitrogen Metabolism in Wheat. Molecular Plant-Microbe Interactions, 2008, 21, 70-78.	2.6	77
57	Gene Expression of Different Wheat Genotypes During Attack by Virulent and Avirulent Hessian Fly (<i>Mayetiola destructor</i>) Larvae. Journal of Chemical Ecology, 2007, 33, 2171-2194.	1.8	105
58	Cloning and characterization of cDNAs encoding carboxypeptidase-like proteins from the gut of Hessian fly larvae [<i>Mayetiola destructor</i> (Say)]. Insect Biochemistry and Molecular Biology, 2006, 36, 665-673.	2.7	9
59	Genes encoding a group of related small secreted proteins from the gut of Hessian fly larvae [<i>Mayetiola destructor</i> (Say)]. Insect Science, 2006, 13, 339-348.	3.0	5
60	H22, a major resistance gene to the Hessian fly (<i>Mayetiola destructor</i>), is mapped to the distal region of wheat chromosome 1DS. Theoretical and Applied Genetics, 2006, 113, 1491-1496.	3.6	19
61	Genomic analysis of a 1 Mb region near the telomere of Hessian fly chromosome X2 and avirulence gene vH13. BMC Genomics, 2006, 7, 7.	2.8	24
62	A super-family of genes coding for secreted salivary gland proteins from the Hessian fly, <i>Mayetiola destructor</i> . Journal of Insect Science, 2006, 6, 1-11.	1.5	16
63	Genetic characterization and molecular mapping of a Hessian fly-resistance gene transferred from <i>T. turgidum</i> ssp. <i>dicoccum</i> to common wheat. Theoretical and Applied Genetics, 2005, 111, 1308-1315.	3.6	73
64	H9, H10, and H11 compose a cluster of Hessian fly-resistance genes in the distal gene-rich region of wheat chromosome 1AS. Theoretical and Applied Genetics, 2005, 110, 1473-1480.	3.6	54
65	Hessian fly resistance gene H13 is mapped to a distal cluster of resistance genes in chromosome 6DS of wheat. Theoretical and Applied Genetics, 2005, 111, 243-249.	3.6	56
66	Cloning and characterization of chymotrypsin- and trypsin-like cDNAs from the gut of the Hessian fly [<i>say</i>]. Insect Biochemistry and Molecular Biology, 2005, 35, 23-32.	2.7	52
67	A group of related cDNAs encoding secreted proteins from Hessian fly [<i>Mayetiola destructor</i> (Say)] salivary glands. Insect Molecular Biology, 2004, 13, 101-108.	2.0	61
68	Characterization of two genes expressed in the salivary glands of the Hessian fly, <i>Mayetiola destructor</i> (Say). Insect Biochemistry and Molecular Biology, 2004, 34, 229-237.	2.7	18