

Raimund Tenhaken

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/5632926/publications.pdf>

Version: 2024-02-01

64
papers

6,147
citations

172457

29
h-index

118850

62
g-index

64
all docs

64
docs citations

64
times ranked

6646
citing authors

#	ARTICLE	IF	CITATIONS
1	Nectaries in ferns: their taxonomic distribution, structure, function, and sugar composition. <i>American Journal of Botany</i> , 2022, 109, 46-57.	1.7	13
2	Characterization of an Arabidopsis Defensin-like Gene Conferring Resistance against Nematodes. <i>Plants</i> , 2022, 11, 280.	3.5	2
3	Overexpression of UDP-Glucose 4-epimerase leads to higher sensitivity towards galactose, providing new insights into the mechanisms of galactose toxicity in plants. <i>Plant Journal</i> , 2022, 109, 1416-1426.	5.7	4
4	Photodynamic Inactivation of plant pathogens part II: fungi. <i>Photochemical and Photobiological Sciences</i> , 2022, 21, 195-207.	2.9	9
5	Galactose induces formation of cell wall stubs and cell death in Arabidopsis roots. <i>Planta</i> , 2022, 256, .	3.2	6
6	Phosphoglucomutase Is Not the Target for Galactose Toxicity in Plants. <i>Frontiers in Plant Science</i> , 2020, 11, 167.	3.6	11
7	Ascorbate oxidation activates systemic defence against root-knot nematode <i>Meloidogyne graminicola</i> in rice. <i>Journal of Experimental Botany</i> , 2020, 71, 4271-4284.	4.8	26
8	The <i>Myo</i> -inositol pathway does not contribute to ascorbic acid synthesis. <i>Plant Biology</i> , 2019, 21, 95-102.	3.8	37
9	Bimodal Pollination Systems in Andean Melastomataceae Involving Birds, Bats, and Rodents. <i>American Naturalist</i> , 2019, 194, 104-116.	2.1	47
10	Arabidopsis MAP-Kinase 3 Phosphorylates UDP-Glucose Dehydrogenase: a Key Enzyme Providing UDP-Sugar for Cell Wall Biosynthesis. <i>Plant Molecular Biology Reporter</i> , 2018, 36, 870-877.	1.8	6
11	Transient alkalinization of the leaf apoplast stiffens the cell wall during onset of chloride salinity in corn leaves. <i>Journal of Biological Chemistry</i> , 2017, 292, 18800-18813.	3.4	34
12	Raffinose Family Oligosaccharides Act As Galactose Stores in Seeds and Are Required for Rapid Germination of Arabidopsis in the Dark. <i>Frontiers in Plant Science</i> , 2016, 7, 1115.	3.6	57
13	The role of arabinokinase in arabinose toxicity in plants. <i>Plant Journal</i> , 2016, 87, 376-390.	5.7	13
14	Generation of PHB from Spent Sulfite Liquor Using Halophilic Microorganisms. <i>Microorganisms</i> , 2015, 3, 268-289.	3.6	17
15	Molecular cloning of AtRS4, a seed specific multifunctional RFO synthase/galactosylhydrolase in Arabidopsis thaliana. <i>Frontiers in Plant Science</i> , 2015, 6, 789.	3.6	27
16	Molecular Cloning of a Novel Glucuronokinase/Putative Pyrophosphorylase from Zebrafish Acting in an UDP-Glucuronic Acid Salvage Pathway. <i>PLoS ONE</i> , 2014, 9, e89690.	2.5	8
17	Myo-inositol oxygenase is important for the removal of excess myo-inositol from syncytia induced by <i>Heterodera schachtii</i> in Arabidopsis roots. <i>New Phytologist</i> , 2014, 201, 476-485.	7.3	46
18	Quantitative HPLC-MS analysis of nucleotide sugars in plant cells following off-line SPE sample preparation. <i>Analytical and Bioanalytical Chemistry</i> , 2014, 406, 3229-3237.	3.7	34

#	ARTICLE	IF	CITATIONS
19	Cell wall remodeling under abiotic stress. <i>Frontiers in Plant Science</i> , 2014, 5, 771.	3.6	509
20	The effect of Translationally Controlled Tumour Protein (TCTP) on programmed cell death in plants. <i>BMC Plant Biology</i> , 2013, 13, 135.	3.6	47
21	A mutation in the <i>Arabidopsis thaliana</i> cell wall biosynthesis gene <i>pectin methylesterase 3</i> as well as its aberrant expression cause hypersensitivity specifically to Zn. <i>Plant Journal</i> , 2013, 76, 151-164.	5.7	36
22	UDP-sugar pyrophosphorylase controls the activity of proceeding sugar-1-kinases enzymes. <i>Plant Signaling and Behavior</i> , 2013, 8, e25478.	2.4	8
23	Molecular and biochemical analysis of the first ARA6 homologue, a RAB5 GTPase, from green algae. <i>Journal of Experimental Botany</i> , 2013, 64, 5553-5568.	4.8	26
24	UDP-sugar pyrophosphorylase is essential for arabinose and xylose recycling, and is required during vegetative and reproductive growth in <i>Arabidopsis</i> . <i>Plant Journal</i> , 2013, 74, 239-247.	5.7	51
25	An emerging role of pectic rhamnogalacturonanII for cell wall integrity. <i>Plant Signaling and Behavior</i> , 2012, 7, 298-299.	2.4	5
26	Cell Wall Ingrowths in Nematode Induced Syncytia Require UGD2 and UGD3. <i>PLoS ONE</i> , 2012, 7, e41515.	2.5	37
27	Down-regulation of the myo-inositol oxygenase gene family has no effect on cell wall composition in <i>Arabidopsis</i> . <i>Planta</i> , 2011, 234, 157-169.	3.2	41
28	Investigations on N-rich protein (NRP) of <i>Arabidopsis thaliana</i> under different stress conditions. <i>Plant Physiology and Biochemistry</i> , 2011, 49, 293-302.	5.8	21
29	Down-regulation of UDP-glucuronic Acid Biosynthesis Leads to Swollen Plant Cell Walls and Severe Developmental Defects Associated with Changes in Pectic Polysaccharides. <i>Journal of Biological Chemistry</i> , 2011, 286, 39982-39992.	3.4	104
30	Characterization of GDP-mannose Dehydrogenase from the Brown Alga <i>Ectocarpus siliculosus</i> Providing the Precursor for the Alginate Polymer. <i>Journal of Biological Chemistry</i> , 2011, 286, 16707-16715.	3.4	29
31	Cloning of Glucuronokinase from <i>Arabidopsis thaliana</i> , the Last Missing Enzyme of the myo-Inositol Oxygenase Pathway to Nucleotide Sugars. <i>Journal of Biological Chemistry</i> , 2010, 285, 2902-2910.	3.4	46
32	Myo-inositol Oxygenase Controls the Level of Myo-inositol in <i>Arabidopsis</i> , But Does Not Increase Ascorbic Acid. <i>Plant Physiology</i> , 2009, 149, 1042-1049.	4.8	108
33	Myo-inositol oxygenase genes are involved in the development of syncytia induced by <i>Heterodera schachtii</i> in <i>Arabidopsis</i> roots. <i>New Phytologist</i> , 2009, 184, 457-472.	7.3	51
34	Nonradioactive enzyme measurement by high-performance liquid chromatography of partially purified sugar-1-kinase (glucuronokinase) from pollen of <i>Lilium longiflorum</i> . <i>Analytical Biochemistry</i> , 2009, 388, 254-259.	2.4	8
35	OCCURRENCE AND CHARACTERIZATION OF ARABINOGLACTAN-LIKE PROTEINS AND HEMICELLULOSES IN <i>MICRASTERIAS</i> (STREPTOPHYTA). <i>Journal of Phycology</i> , 2008, 44, 1221-1234.	2.3	73
36	Genome-wide analysis of the UDP-glucose dehydrogenase gene family in <i>Arabidopsis</i> , a key enzyme for matrix polysaccharides in cell walls. <i>Journal of Experimental Botany</i> , 2007, 58, 3609-3621.	4.8	95

#	ARTICLE	IF	CITATIONS
37	Isolation of a novel ABC-transporter gene from soybean induced by salicylic acid. <i>Journal of Experimental Botany</i> , 2006, 57, 2193-2201.	4.8	73
38	DCD - a novel plant specific domain in proteins involved in development and programmed cell death. <i>BMC Bioinformatics</i> , 2005, 6, 169.	2.6	58
39	The inositol oxygenase gene family of Arabidopsis is involved in the biosynthesis of nucleotide sugar precursors for cell-wall matrix polysaccharides. <i>Planta</i> , 2005, 221, 243-254.	3.2	135
40	A mitogen-activated-protein kinase from soybean is activated by a pathogen and novel functional analogs of salicylic acid. <i>Plant Physiology and Biochemistry</i> , 2003, 41, 929-934.	5.8	2
41	Purification, characterization and functional cloning of inositol oxygenase from <i>Cryptococcus</i> . <i>Yeast</i> , 2003, 20, 1317-1329.	1.7	24
42	Recombinant UDP-glucose dehydrogenase from soybean. <i>Plant Physiology and Biochemistry</i> , 2002, 40, 1011-1017.	5.8	28
43	Transgenic Plants with Enhanced Tolerance against Microbial Pathogens. , 2002, , .		0
44	Suppression of the ribosomal L2 gene reveals a novel mechanism for stress adaptation in soybean. <i>Planta</i> , 2001, 212, 792-798.	3.2	27
45	WY-14,643 and other agonists of the peroxisome proliferator-activated receptor reveal a new mode of action for salicylic acid in soybean disease resistance. <i>Planta</i> , 2001, 212, 888-895.	3.2	11
46	Title is missing!. <i>European Journal of Plant Pathology</i> , 2001, 107, 323-336.	1.7	31
47	Matrix polysaccharide precursors in Arabidopsis cell walls are synthesized by alternate pathways with organ-specific expression patterns. <i>Plant Journal</i> , 2000, 21, 537-546.	5.7	79
48	Cloning, Expression and Characterization of Protein Elicitors from the Soybean Pathogenic Fungus <i>Phytophthora sojae</i> . <i>Journal of Phytopathology</i> , 2000, 148, 161-167.	1.0	14
49	Defence gene expression in soybean is linked to the status of the cell death program. , 2000, 44, 209-218.		13
50	Cloning of genes by mRNA differential display induced during the hypersensitive reaction of soybean after inoculation with <i>Pseudomonas syringae</i> pv. <i>glycinea</i> . <i>Plant Molecular Biology</i> , 1998, 38, 1225-1234.	3.9	75
51	Cloning of putative subunits of the soybean plasma membrane NADPH oxidase involved in the oxidative burst by antibody expression screening. <i>Protoplasma</i> , 1998, 205, 21-28.	2.1	18
52	Induction of alkalization and an oxidative burst by low doses of cycloheximide in soybean cells. <i>Planta</i> , 1998, 206, 666-672.	3.2	22
53	Chitinase in cucumber hypocotyls is induced by germinating fungal spores and by fungal elicitor in synergism with inducers of acquired resistance. <i>Plant Journal</i> , 1998, 13, 447-454.	5.7	50
54	Characterization of a Diffusible Signal Capable of Inducing Defense Gene Expression in Tobacco. <i>Plant Physiology</i> , 1997, 113, 621-629.	4.8	30

#	ARTICLE	IF	CITATIONS
55	Salicylic Acid Is Needed in Hypersensitive Cell Death in Soybean but Does Not Act as a Catalase Inhibitor. <i>Plant Physiology</i> , 1997, 115, 291-298.	4.8	88
56	Characterization and Cloning of Cutinase from <i>Ascochyta rabiei</i> . <i>Zeitschrift Fur Naturforschung - Section C Journal of Biosciences</i> , 1997, 52, 197-208.	1.4	22
57	Cloning of an Enzyme That Synthesizes a Key Nucleotide-Sugar Precursor of Hemicellulose Biosynthesis from Soybean:UDP-Glucose Dehydrogenase. <i>Plant Physiology</i> , 1996, 112, 1127-1134.	4.8	122
58	Function of the oxidative burst in hypersensitive disease resistance.. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1995, 92, 4158-4163.	7.1	377
59	Function of Oxidative Cross-Linking of Cell Wall Structural Proteins in Plant Disease Resistance.. <i>Plant Cell</i> , 1994, 6, 1703-1712.	6.6	484
60	H ₂ O ₂ from the oxidative burst orchestrates the plant hypersensitive disease resistance response. <i>Cell</i> , 1994, 79, 583-593.	28.9	2,602
61	Characterization of metabolic changes involved in hypersensitive-like browning reactions of chickpea (<i>Cicer arietinum</i> L.) cell cultures following challenge by <i>Ascochyta rabiei</i> culture filtrate. <i>Physiological and Molecular Plant Pathology</i> , 1994, 44, 141-155.	2.5	15
62	Characterization of Pectic Enzymes from the Chickpea Pathogen <i>Ascochyta rabiei</i> . <i>Zeitschrift Fur Naturforschung - Section C Journal of Biosciences</i> , 1991, 46, 51-57.	1.4	14
63	Purification and characterization of pterocarpan hydroxylase, a flavoprotein monooxygenase from the fungus <i>Ascochyta rabiei</i> involved in pterocarpan phytoalexin metabolism. <i>Archives of Microbiology</i> , 1991, 155, 353.	2.2	18
64	Medicarpin and maackiain 3-O-glucoside-6?-O-malonate conjugates are constitutive compounds in chickpea (<i>Cicer arietinum</i> L.) cell cultures. <i>Plant Cell Reports</i> , 1991, 10, 371-4.	5.6	23