

Naoyuki Tsuchiya

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/5638743/publications.pdf>

Version: 2024-02-01

189
papers

7,169
citations

50276

46
h-index

74163

75
g-index

194
all docs

194
docs citations

194
times ranked

7119
citing authors

#	ARTICLE	IF	CITATIONS
1	<i>MUC5B</i> Promoter Variant and Rheumatoid Arthritis with Interstitial Lung Disease. <i>New England Journal of Medicine</i> , 2018, 379, 2209-2219.	27.0	326
2	Sex-specific association of X-linked Toll-like receptor 7 (TLR7) with male systemic lupus erythematosus. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2010, 107, 15838-15843.	7.1	324
3	Fc γ receptor gene polymorphisms in Japanese patients with systemic lupus erythematosus: Contribution of FCGR2B to genetic susceptibility. <i>Arthritis and Rheumatism</i> , 2002, 46, 1242-1254.	6.7	301
4	Fc γ RIIB Ile232Thr transmembrane polymorphism associated with human systemic lupus erythematosus decreases affinity to lipid rafts and attenuates inhibitory effects on B cell receptor signaling. <i>Human Molecular Genetics</i> , 2005, 14, 2881-2892.	2.9	216
5	BAFF/BLyS can potentiate B-cell selection with the B-cell coreceptor complex. <i>Blood</i> , 2004, 103, 2257-2265.	1.4	151
6	Association of Fc γ receptor IIb and IIIb polymorphisms with susceptibility to systemic lupus erythematosus in Thais. <i>Tissue Antigens</i> , 2003, 61, 374-383.	1.0	146
7	Association of Fc γ receptor IIb polymorphism with susceptibility to systemic lupus erythematosus in Chinese: a common susceptibility gene in the Asian populations. <i>Tissue Antigens</i> , 2004, 63, 21-27.	1.0	142
8	Effects of galactose depletion from oligosaccharide chains on immunological activities of human IgG. <i>Journal of Rheumatology</i> , 1989, 16, 285-90.	2.0	141
9	Association of tumor necrosis factor receptor 2 (TNFR2) polymorphism with susceptibility to systemic lupus erythematosus. <i>Tissue Antigens</i> , 1999, 53, 527-533.	1.0	127
10	Association of hla-b39 with hla-b27-negative ankylosing spondylitis and pauciarticular juvenile rheumatoid arthritis in Japanese patients. <i>Arthritis and Rheumatism</i> , 1995, 38, 1672-1677.	6.7	124
11	Association of Fc γ receptor IIIb, but not of Fc γ receptor IIA and IIIA, polymorphisms with systemic lupus erythematosus in Japanese. <i>Genes and Immunity</i> , 1999, 1, 53-60.	4.1	121
12	Association of a functional polymorphism in the <i>IRF5</i> region with systemic sclerosis in a Japanese population. <i>Arthritis and Rheumatism</i> , 2009, 60, 1845-1850.	6.7	115
13	Analysis on the association of human BLYS (BAFF, TNFSF13B) polymorphisms with systemic lupus erythematosus and rheumatoid arthritis. <i>Genes and Immunity</i> , 2002, 3, 424-429.	4.1	100
14	Association of <i>IRF5</i> polymorphisms with systemic lupus erythematosus in a Japanese population: Support for a crucial role of intron 1 polymorphisms. <i>Arthritis and Rheumatism</i> , 2008, 58, 826-834.	6.7	100
15	TLR7 single-nucleotide polymorphisms in the 3' untranslated region and intron 2 independently contribute to systemic lupus erythematosus in Japanese women: a case-control association study. <i>Arthritis Research and Therapy</i> , 2011, 13, R41.	3.5	93
16	Comparison of statistical power between 2 \times 2 allele frequency and allele positivity tables in case-control studies of complex disease genes. <i>Annals of Human Genetics</i> , 2001, 65, 197-206.	0.8	93
17	Comparison of statistical power between 2x2 allele frequency and allele positivity tables in case-control studies of complex disease genes. <i>Annals of Human Genetics</i> , 2001, 65, 197-206.	0.8	91
18	Crucial Role of Inhibitor of DNA Binding/Differentiation in the Vascular Endothelial Growth Factor-Induced Activation and Angiogenic Processes of Human Endothelial Cells. <i>Journal of Immunology</i> , 2004, 173, 5801-5809.	0.8	88

#	ARTICLE	IF	CITATIONS
19	Role of STAT4 polymorphisms in systemic lupus erythematosus in a Japanese population: a case-control association study of the STAT1-STAT4 region. <i>Arthritis Research and Therapy</i> , 2008, 10, R113.	3.5	88
20	A compass that points to lupus: genetic studies on type I interferon pathway. <i>Genes and Immunity</i> , 2007, 8, 445-455.	4.1	84
21	MIC-A polymorphism in Japanese and a MIC-A-MIC-B null haplotype. <i>Immunogenetics</i> , 1999, 49, 620-628.	2.4	82
22	Association of a functional CD19 polymorphism with susceptibility to systemic sclerosis. <i>Arthritis and Rheumatism</i> , 2004, 50, 4002-4007.	6.7	82
23	Transethnic meta-analysis identifies <i>GSDMA</i> and <i>PRDM1</i> as susceptibility genes to systemic sclerosis. <i>Annals of the Rheumatic Diseases</i> , 2017, 76, 1150-1158.	0.9	77
24	Association of the <i>FAM167A</i> "BLK" region with systemic sclerosis. <i>Arthritis and Rheumatism</i> , 2010, 62, 890-895.	6.7	76
25	The Q223R polymorphism in <i>LEPR</i> is associated with obesity in Pacific Islanders. <i>Human Genetics</i> , 2010, 127, 287-294.	3.8	74
26	Molecular genetic analyses of human <i>NKG2C</i> (<i>KLRC2</i>) gene deletion. <i>International Immunology</i> , 2004, 16, 163-168.	4.0	73
27	Association of TNFAIP3 interacting protein 1, <i>TNIP1</i> with systemic lupus erythematosus in a Japanese population: a case-control association study. <i>Arthritis Research and Therapy</i> , 2010, 12, R174.	3.5	70
28	Association of Human Leukocyte Antigen with Interstitial Lung Disease in Rheumatoid Arthritis: A Protective Role for Shared Epitope. <i>PLoS ONE</i> , 2012, 7, e33133.	2.5	70
29	Genetic background of Japanese patients with antineutrophil cytoplasmic antibody-associated vasculitis: association of HLA-DRB1*0901 with microscopic polyangiitis. <i>Journal of Rheumatology</i> , 2003, 30, 1534-40.	2.0	70
30	Analysis of the association of HLA-DRB1, <i>TNF</i> \pm promoter and <i>TNFR2</i> (<i>TNFRSF1B</i>) polymorphisms with SLE using transmission disequilibrium test. <i>Genes and Immunity</i> , 2001, 2, 317-322.	4.1	69
31	Extensive polymorphisms of <i>LILRB1</i> (<i>ILT2</i> , <i>LIR1</i>) and their association with HLA-DRB1 shared epitope negative rheumatoid arthritis. <i>Human Molecular Genetics</i> , 2005, 14, 2469-2480.	2.9	69
32	Identification of the gene variations in human <i>CD22</i> . <i>Immunogenetics</i> , 1999, 49, 280-286.	2.4	68
33	Cross-reactive epitope with <i>Klebsiella pneumoniae</i> nitrogenase in articular tissue of HLA "B27+ patients with ankylosing spondylitis. <i>Arthritis and Rheumatism</i> , 1989, 32, 437-445.	6.7	66
34	Association of <i>STAT4</i> polymorphism with systemic sclerosis in a Japanese population. <i>Annals of the Rheumatic Diseases</i> , 2009, 68, 1375-1376.	0.9	64
35	Studies on the association of <i>Fcγ3 receptor IIA, IIB, IIIA and IIIB polymorphisms with rheumatoid arthritis in the Japanese: evidence for a genetic interaction between HLA-DRB1 and <i>FCGR3A</i>. <i>Genes and Immunity</i>, 2002, 3, 488-493.</i>	4.1	62
36	<i>PLD4</i> as a novel susceptibility gene for systemic sclerosis in a Japanese population. <i>Arthritis and Rheumatism</i> , 2013, 65, 472-480.	6.7	62

#	ARTICLE	IF	CITATIONS
37	Protective Effect of the HLA-DRB1*13:02 Allele in Japanese Rheumatoid Arthritis Patients. <i>PLoS ONE</i> , 2014, 9, e99453.	2.5	60
38	Lack of a strong association of CTLA-4 exon 1 polymorphism with the susceptibility to rheumatoid arthritis and systemic lupus erythematosus in Japanese: an association study using a novel variation screening method. <i>Tissue Antigens</i> , 1999, 54, 578-584.	1.0	57
39	Replication of the association between the C8orf13-BLK region and systemic lupus erythematosus in a Japanese population. <i>Arthritis and Rheumatism</i> , 2009, 60, 553-558.	6.7	57
40	Tumor necrosis factor \pm 5 ϵ 2-flanking region, tumor necrosis factor receptor II, and HLA ϵ DRB1 polymorphisms in Japanese patients with rheumatoid arthritis. <i>Arthritis and Rheumatism</i> , 2000, 43, 753.	6.7	56
41	Targeting Id1 and Id3 inhibits peritoneal metastasis of gastric cancer. <i>Cancer Science</i> , 2005, 96, 784-790.	3.9	53
42	Variations of human killer cell lectin-like receptors: common occurrence of NKG2-C deletion in the general population. <i>Genes and Immunity</i> , 2003, 4, 160-167.	4.1	51
43	Genome, epigenome and transcriptome analyses of a pair of monozygotic twins discordant for systemic lupus erythematosus. <i>Human Immunology</i> , 2013, 74, 170-175.	2.4	51
44	Association of a functional polymorphism in the 3 ϵ 2-untranslated region of SPI1 with systemic lupus erythematosus. <i>Arthritis and Rheumatism</i> , 2011, 63, 755-763.	6.7	50
45	Human Leukocyte Antigens and Systemic Lupus Erythematosus: A Protective Role for the HLA-DR6 Alleles DRB1*13:02 and *14:03. <i>PLoS ONE</i> , 2014, 9, e87792.	2.5	50
46	Role of the Fc γ 3 receptor IIA polymorphism in the antiphospholipid syndrome: An international meta-analysis. <i>Arthritis and Rheumatism</i> , 2003, 48, 1930-1938.	6.7	49
47	Association of HLA-DRB1*0901-DQB1*0303 haplotype with microscopic polyangiitis in Japanese. <i>Genes and Immunity</i> , 2006, 7, 81-84.	4.1	49
48	Independent contribution of HLA-DRB1 and TNF \pm promoter polymorphisms to the susceptibility to Crohn ϵ ™s disease. <i>Genes and Immunity</i> , 2000, 1, 351-357.	4.1	48
49	Detection of glycosylation abnormality in rheumatoid IgG using N-acetylglucosamine-specific <i>Psathyrella velutina</i> lectin. <i>Journal of Immunology</i> , 1993, 151, 1137-46.	0.8	46
50	Elevated serum level of soluble HLA class I antigens in patients with systemic lupus erythematosus. <i>Arthritis and Rheumatism</i> , 1996, 39, 792-796.	6.7	45
51	MICA allele typing of HLA-B27 positive Japanese patients with seronegative spondylarthropathies and healthy individuals: Differential linkage disequilibrium with HLA-B27 subtypes. <i>Arthritis and Rheumatism</i> , 1998, 41, 68-73.	6.7	45
52	Variations in immune response genes and their associations with multifactorial immune disorders. <i>Immunological Reviews</i> , 2002, 190, 169-181.	6.0	45
53	The Impact of Natural Selection on an ABCC11 SNP Determining Earwax Type. <i>Molecular Biology and Evolution</i> , 2011, 28, 849-857.	8.9	44
54	The role of common protective alleles HLA-DRB1*13 among systemic autoimmune diseases. <i>Genes and Immunity</i> , 2017, 18, 1-7.	4.1	44

#	ARTICLE	IF	CITATIONS
55	Immunogenetic features in 120 Japanese patients with idiopathic inflammatory myopathy. <i>Journal of Rheumatology</i> , 2004, 31, 1768-74.	2.0	44
56	New polymorphisms of human CD80 and CD86: lack of association with rheumatoid arthritis and systemic lupus erythematosus. <i>Genes and Immunity</i> , 2000, 1, 428-434.	4.1	43
57	CD72 polymorphisms associated with alternative splicing modify susceptibility to human systemic lupus erythematosus through epistatic interaction with FCGR2B. <i>Human Molecular Genetics</i> , 2004, 13, 2907-2917.	2.9	43
58	HLA-A*31:01 and methotrexate-induced interstitial lung disease in Japanese rheumatoid arthritis patients: a multidrug hypersensitivity marker? <i>Annals of the Rheumatic Diseases</i> , 2013, 72, 153-155.	0.9	43
59	Association of HLA-A*3303-B*4403-DRB1*1302 haplotype, but not of TNFA promoter and NKp30 polymorphism, with postherpetic neuralgia (PHN) in the Japanese population. <i>Genes and Immunity</i> , 2002, 3, 477-481.	4.1	42
60	Association study of a polymorphism of the CTGF gene and susceptibility to systemic sclerosis in the Japanese population. <i>Annals of the Rheumatic Diseases</i> , 2009, 68, 1921-1924.	0.9	42
61	Association of HLA-DRB1*1502-DQB1*0501 haplotype with susceptibility to systemic lupus erythematosus in Thais. <i>Tissue Antigens</i> , 2002, 59, 113-117.	1.0	40
62	Association of killer cell immunoglobulin-like receptor genotypes with microscopic polyangiitis. <i>Arthritis and Rheumatism</i> , 2006, 54, 992-997.	6.7	39
63	Allele typing of human TNFA 5' flanking region using polymerase chain reaction-preferential homoduplex formation assay (PCR-PHFA): linkage disequilibrium with HLA class I and class II genes in Japanese. <i>Tissue Antigens</i> , 1999, 54, 478-484.	1.0	38
64	Association of CD22 gene polymorphism with susceptibility to limited cutaneous systemic sclerosis. <i>Tissue Antigens</i> , 2007, 69, 242-249.	1.0	38
65	Overlapping peptide-binding specificities of HLA-B27 and B39: Evidence for a role of peptide supermotif in the pathogenesis of spondylarthropathies. <i>Arthritis and Rheumatism</i> , 1999, 42, 175-181.	6.7	36
66	Autoantibodies to the HLA-B27 sequence cross-react with the hypothetical peptide from the arthritis-associated Shigella plasmid. <i>Journal of Clinical Investigation</i> , 1990, 86, 1193-1203.	8.2	36
67	Association of IL-10 receptor 2 (IL10RB) SNP with systemic sclerosis. <i>Biochemical and Biophysical Research Communications</i> , 2008, 373, 403-407.	2.1	35
68	Cumulative association of eight susceptibility genes with systemic lupus erythematosus in a Japanese female population. <i>Journal of Human Genetics</i> , 2011, 56, 503-507.	2.3	35
69	Protective Role of HLA-DRB1*13:02 against Microscopic Polyangiitis and MPO-ANCA-Positive Vasculitides in a Japanese Population: A Case-Control Study. <i>PLoS ONE</i> , 2016, 11, e0154393.	2.5	35
70	Association of Fcγ receptor IIA, but not IIB and IIIA, polymorphisms with systemic lupus erythematosus: A family-based association study in Caucasians. <i>Arthritis and Rheumatism</i> , 2004, 50, 671-673.	6.7	34
71	The human histocompatibility leukocyte antigen (HLA) haplotype is associated with the onset of postherpetic neuralgia after herpes zoster. <i>Pain</i> , 2004, 110, 329-336.	4.2	33
72	Expression of ID Family Genes in the Synovia from Patients with Rheumatoid Arthritis. <i>Biochemical and Biophysical Research Communications</i> , 2001, 284, 436-442.	2.1	32

#	ARTICLE	IF	CITATIONS
73	Role of Fc γ 3 receptor IIb polymorphism in the genetic background of systemic lupus erythematosus: Insights from Asia. <i>Autoimmunity</i> , 2005, 38, 347-352.	2.6	32
74	Differential association of HLA-DRB1 alleles in Japanese patients with early rheumatoid arthritis in relationship to autoantibodies to cyclic citrullinated peptide. <i>Clinical and Experimental Rheumatology</i> , 2007, 25, 219-24.	0.8	32
75	Association of TNFAIP3 Polymorphism with Susceptibility to Systemic Lupus Erythematosus in a Japanese Population. <i>Journal of Biomedicine and Biotechnology</i> , 2010, 2010, 1-5.	3.0	31
76	Induction of alternative splicing of HLA-B27 by bacterial invasion. <i>Arthritis and Rheumatism</i> , 1997, 40, 694-703.	6.7	30
77	Association of LILRA2 (ILT1, LIR7) splice site polymorphism with systemic lupus erythematosus and microscopic polyangiitis. <i>Genes and Immunity</i> , 2008, 9, 214-223.	4.1	30
78	Single nucleotide polymorphisms in the coding regions of human CXC-chemokine receptors CXCR1, CXCR2 and CXCR3. <i>Genes and Immunity</i> , 2000, 1, 330-337.	4.1	29
79	New single nucleotide polymorphisms in the coding region of human TNFR2: association with systemic lupus erythematosus. <i>Genes and Immunity</i> , 2000, 1, 501-503.	4.1	29
80	Variations in the human Th2-specific chemokine TARC gene. <i>Immunogenetics</i> , 2003, 54, 742-745.	2.4	29
81	Association of human leukocyte antigen alleles with chronic lung diseases in rheumatoid arthritis. <i>Rheumatology</i> , 2016, 55, 1301-1307.	1.9	29
82	Plasma miRNA expression profiles in rheumatoid arthritis associated interstitial lung disease. <i>BMC Musculoskeletal Disorders</i> , 2017, 18, 21.	1.9	29
83	Lambert-Eaton Myasthenic Syndrome Associated with Sjögren's Syndrome and Discoid Lupus Erythematosus. <i>Scandinavian Journal of Rheumatology</i> , 1993, 22, 302-304.	1.1	27
84	Polymorphisms of human CD19 gene: possible association with susceptibility to systemic lupus erythematosus in Japanese. <i>Genes and Immunity</i> , 2002, 3, S21-S30.	4.1	26
85	Serum biomarker analysis of collagen disease patients with acute-onset diffuse interstitial lung disease. <i>BMC Immunology</i> , 2013, 14, 9.	2.2	26
86	A replication study confirms the association of GWAS-identified SNPs at MICB and PLCE1 in Thai patients with dengue shock syndrome. <i>BMC Medical Genetics</i> , 2014, 15, 58.	2.1	26
87	HLA-DRB1 and DQB1 alleles in Japanese type 1 autoimmune hepatitis: The predisposing role of the DR4/DR8 heterozygous genotype. <i>PLoS ONE</i> , 2017, 12, e0187325.	2.5	26
88	Effects of HLA-DRB1 alleles on susceptibility and clinical manifestations in Japanese patients with adult onset Still's disease. <i>Arthritis Research and Therapy</i> , 2017, 19, 199.	3.5	25
89	Human Leukocyte Antigen and Systemic Sclerosis in Japanese: The Sign of the Four Independent Protective Alleles, DRB1*13:02, DRB1*14:06, DQB1*03:01, and DPB1*02:01. <i>PLoS ONE</i> , 2016, 11, e0154255.	2.5	25
90	Studies of humoral and cell-mediated immunity to peptides shared by HLA-27.1 and Klebsiella pneumoniae nitrogenase in ankylosing spondylitis. <i>Clinical and Experimental Immunology</i> , 1989, 76, 354-60.	2.6	25

#	ARTICLE	IF	CITATIONS
91	Presence of four major haplotypes in human BCMA gene: lack of association with systemic lupus erythematosus and rheumatoid arthritis. <i>Genes and Immunity</i> , 2001, 2, 276-279.	4.1	24
92	Variations in the human CC chemokine eotaxin gene. <i>Genes and Immunity</i> , 2001, 2, 461-463.	4.1	24
93	Replication of association between FAM167A(C8orf13)-BLK region and rheumatoid arthritis in a Japanese population. <i>Annals of the Rheumatic Diseases</i> , 2010, 69, 936-937.	0.9	24
94	Identification of secreted phosphoprotein 1 gene as a new rheumatoid arthritis susceptibility gene. <i>Annals of the Rheumatic Diseases</i> , 2015, 74, e19-e19.	0.9	24
95	Epistatic Interaction between BANK1 and BLK in Rheumatoid Arthritis: Results from a Large Trans-Ethnic Meta-Analysis. <i>PLoS ONE</i> , 2013, 8, e61044.	2.5	24
96	New variations of human CC-chemokine receptors CCR3 and CCR4. <i>Genes and Immunity</i> , 1999, 1, 97-104.	4.1	23
97	Genetics of ANCA-associated vasculitis in Japan: a role for HLA-DRB1*09:01 haplotype. <i>Clinical and Experimental Nephrology</i> , 2013, 17, 628-630.	1.6	23
98	Association of <i>MUC5B</i> promoter polymorphism with interstitial lung disease in myeloperoxidase-antineutrophil cytoplasmic antibody-associated vasculitis. <i>Annals of the Rheumatic Diseases</i> , 2019, 78, 1144-1146.	0.9	23
99	Distribution of Glycosylation Abnormality among Serum IgG Subclasses from Patients with Rheumatoid Arthritis. <i>Clinical Immunology and Immunopathology</i> , 1994, 70, 47-50.	2.0	22
100	Role of APRIL (TNFSF13) polymorphisms in the susceptibility to systemic lupus erythematosus in Japanese. <i>Rheumatology</i> , 2007, 46, 776-782.	1.9	22
101	Rheumatoid factors may bear the internal image of the Fc gamma-binding protein of herpes simplex virus type 1. <i>Journal of Immunology</i> , 1990, 144, 4742-8.	0.8	22
102	Association of Increased Frequencies of HLA-DPB1*05:01 with the Presence of Anti-Ro/SS-A and Anti-La/SS-B Antibodies in Japanese Rheumatoid Arthritis and Systemic Lupus Erythematosus Patients. <i>PLoS ONE</i> , 2013, 8, e53910.	2.5	21
103	Identification of a haplotype block in the 5q31 cytokine gene cluster associated with the susceptibility to severe malaria. <i>Malaria Journal</i> , 2009, 8, 232.	2.3	20
104	Association of ADAMTS13 polymorphism with cerebral malaria. <i>Malaria Journal</i> , 2011, 10, 366.	2.3	19
105	Association of HLA-C 3' UTR Untranslated Region Polymorphisms with Systemic Lupus Erythematosus in a Japanese Population: A Case-Control Association Study. <i>PLoS ONE</i> , 2016, 11, e0158065.	2.5	19
106	Role of B cell inhibitory receptor polymorphisms in systemic lupus erythematosus: a negative times a negative makes a positive. <i>Journal of Human Genetics</i> , 2006, 51, 741-750.	2.3	18
107	Association of <i>PHRF1-IRF7</i> region polymorphism with clinical manifestations of systemic lupus erythematosus in a Japanese population. <i>Lupus</i> , 2012, 21, 890-895.	1.6	18
108	HLA-B27 subtypes in Japanese with seronegative spondyloarthropathies and healthy controls. <i>Journal of Rheumatology</i> , 1996, 23, 1189-93.	2.0	18

#	ARTICLE	IF	CITATIONS
109	Diversity of Human Immune System Multigene Families and its Implication in the Genetic Background of Rheumatic Diseases. <i>Current Medicinal Chemistry</i> , 2007, 14, 431-439.	2.4	17
110	Anti-citrullinated glucose-6-phosphate isomerase peptide antibodies in patients with rheumatoid arthritis are associated with <i>HLA-DRB1</i> shared epitope alleles and disease activity. <i>Clinical and Experimental Immunology</i> , 2013, 172, 44-53.	2.6	17
111	Differential mapping of Fc gamma-binding and monoclonal antibody-reactive epitopes on gE, the Fc gamma-binding glycoprotein of herpes simplex virus type 1. <i>Journal of Immunology</i> , 1992, 149, 2415-27.	0.8	17
112	Association of IRF5, STAT4 and BLK with systemic lupus erythematosus and other rheumatic diseases.. <i>Japanese Journal of Clinical Immunology</i> , 2010, 33, 57-65.	0.0	16
113	Genetics of Interstitial Lung Disease: <i>Vol de Nuit</i> (Night Flight). <i>Clinical Medicine Insights: Circulatory, Respiratory and Pulmonary Medicine</i> , 2015, 9s1, CCRPM.S23283.	0.9	16
114	Effects of <i>APRIL</i> (<i>TNFSF13</i>) polymorphisms and splicing isoforms on the secretion of soluble APRIL. <i>Modern Rheumatology</i> , 2012, 22, 541-549.	1.8	15
115	Association of NCF1 polymorphism with systemic lupus erythematosus and systemic sclerosis but not with ANCA-associated vasculitis in a Japanese population. <i>Scientific Reports</i> , 2019, 9, 16366.	3.3	15
116	New variations of human SHP-1. <i>Immunogenetics</i> , 1999, 49, 577-579.	2.4	14
117	Association of ETS1 polymorphism with granulomatosis with polyangiitis and proteinase 3-anti-neutrophil cytoplasmic antibody positive vasculitis in a Japanese population. <i>Journal of Human Genetics</i> , 2018, 63, 55-62.	2.3	14
118	Cytomegalovirus genomes demonstrated by polymerase chain reaction in synovial fluid from rheumatoid arthritis patients. <i>Clinical and Experimental Rheumatology</i> , 1992, 10, 161-4.	0.8	14
119	Elevated Cytokine Levels in Synovial Fluid of Rheumatoid Arthritis Correlates with the Presence of Cytomegalovirus Genome. <i>Autoimmunity</i> , 1994, 17, 333-337.	2.6	13
120	HLA-B associations of HLA-B27 negative ankylosing spondylitis: Comment on the article by Yamaguchi et al. <i>Arthritis and Rheumatism</i> , 1996, 39, 1768-1769.	6.7	13
121	Human CD72 splicing isoform responsible for resistance to systemic lupus erythematosus regulates serum immunoglobulin level and is localized in endoplasmic reticulum. <i>BMC Immunology</i> , 2012, 13, 72.	2.2	13
122	Antibodies to Human Cytomegalovirus 65-Kilodalton Fc Binding Protein in Rheumatoid Arthritis: Idiotypic Mimicry Hypothesis of Rheumatoid Factor Production. <i>Autoimmunity</i> , 1993, 15, 39-48.	2.6	12
123	Evaluation of microsatellite markers in association studies: a search for an immune-related susceptibility gene in sarcoidosis. <i>Immunogenetics</i> , 2005, 56, 861-870.	2.4	12
124	Advances in the genomics of ANCA-associated vasculitis—a view from East Asia. <i>Genes and Immunity</i> , 2021, 22, 1-11.	4.1	12
125	Rheumatoid factors react with fab fragments of monoclonal antibodies to herpes simplex virus types 1 and 2 $\text{Fc}\gamma$ -binding proteins. <i>Arthritis and Rheumatism</i> , 1991, 34, 846-855.	6.7	11
126	New variations in human OX40 ligand (CD134L) gene. <i>Genes and Immunity</i> , 2000, 1, 521-522.	4.1	11

#	ARTICLE	IF	CITATIONS
127	Identification of novel single nucleotide substitutions in the NKp30 gene expressed in human natural killer cells. <i>Tissue Antigens</i> , 2001, 58, 255-258.	1.0	11
128	Antibodies to the peptide from the plasmid-coded <i>Yersinia</i> outer membrane protein (YOP1) in patients with ankylosing spondylitis. <i>Clinical and Experimental Immunology</i> , 2008, 82, 493-498.	2.6	11
129	Association of a single nucleotide polymorphism in the <i>SH2D1A</i> intronic region with systemic lupus erythematosus. <i>Lupus</i> , 2013, 22, 497-503.	1.6	11
130	Molecular mimicry—hypothesis or reality?. <i>Western Journal of Medicine</i> , 1992, 157, 133-8.	0.3	11
131	2-dimensional flow cytometric analysis of peripheral blood T lymphocytes from patients with systemic lupus erythematosus: preferential expression of HLA-DR antigen on the surface of Leu 2a+ cells. <i>Journal of Rheumatology</i> , 1988, 15, 946-51.	2.0	11
132	A novel method for isolation of endothelial cells and macrophages from murine tumors based on Ac-LDL uptake and CD16 expression. <i>Journal of Immunological Methods</i> , 2004, 295, 183-193.	1.4	10
133	IFNGR1 polymorphisms in Thai malaria patients. <i>Infection, Genetics and Evolution</i> , 2009, 9, 1406-1409.	2.3	10
134	A replication study of the association between the IL12B promoter allele CTCTAA and susceptibility to cerebral malaria in Thai population. <i>Malaria Journal</i> , 2009, 8, 290.	2.3	10
135	Association of IRF5 polymorphism with MPO-ANCA-positive vasculitis in a Japanese population. <i>Genes and Immunity</i> , 2013, 14, 527-529.	4.1	10
136	A functional SNP upstream of the beta-2 adrenergic receptor gene (ADRB2) is associated with obesity in Oceanic populations. <i>International Journal of Obesity</i> , 2013, 37, 1204-1210.	3.4	10
137	Autoantibody Profiles in Collagen Disease Patients with Interstitial Lung Disease (ILD): Antibodies to Major Histocompatibility Complex Class I-Related Chain a (MICA) as Markers of ILD. <i>Biomarker Insights</i> , 2015, 10, BMI.S28209.	2.5	10
138	Association of a single nucleotide polymorphism upstream of ICOS with Japanese autoimmune hepatitis type 1. <i>Journal of Human Genetics</i> , 2017, 62, 481-484.	2.3	10
139	Association of HLA-DRB1 genotype with younger age onset and elder age onset rheumatoid arthritis in Japanese populations. <i>Medicine (United States)</i> , 2019, 98, e18218.	1.0	10
140	Association of TERT and DSP variants with microscopic polyangiitis and myeloperoxidase-ANCA positive vasculitis in a Japanese population: a genetic association study. <i>Arthritis Research and Therapy</i> , 2020, 22, 246.	3.5	10
141	TNFR2 position 196 polymorphism in Japanese patients with rheumatoid arthritis: Comment on the article by Dieudonné et al. <i>Arthritis and Rheumatism</i> , 2003, 48, 273-274.	6.7	9
142	HLA-DRB1*08:02 Is Associated with Bucillamine-Induced Proteinuria in Japanese Rheumatoid Arthritis Patients. <i>Biomarker Insights</i> , 2014, 9, BMI.S13654.	2.5	9
143	Association of a single nucleotide polymorphism in TNIP1 with type-1 autoimmune hepatitis in the Japanese population. <i>Journal of Human Genetics</i> , 2018, 63, 739-744.	2.3	9
144	Effects of APRIL (TNFSF13) polymorphisms and splicing isoforms on the secretion of soluble APRIL. <i>Modern Rheumatology</i> , 2012, 22, 541-549.	1.8	9

#	ARTICLE	IF	CITATIONS
145	Significant association of HLA-B and HLA-DRB1 alleles with cleft lip with or without cleft palate. <i>Tissue Antigens</i> , 1999, 53, 147-152.	1.0	8
146	Effects of down-regulating the Id genes in human colorectal cancer cells on early steps of haematogenous metastasis. <i>European Journal of Cancer</i> , 2006, 42, 668-673.	2.8	8
147	Association of BAK1 single nucleotide polymorphism with a risk for dengue hemorrhagic fever. <i>BMC Medical Genetics</i> , 2016, 17, 43.	2.1	8
148	The pattern of GPI-80 expression is a useful marker for unusual myeloid maturation in peripheral blood. <i>Clinical and Experimental Immunology</i> , 2016, 186, 373-386.	2.6	8
149	Exacerbation of Lambert-Eaton Myasthenic Syndrome Caused by an L-type Ca ²⁺ Channel Antagonist.. <i>International Heart Journal</i> , 2003, 44, 139-144.	0.6	8
150	Immunologic Mechanisms in Common Rheumatologic Diseases. <i>Clinical Orthopaedics and Related Research</i> , 1996, 326, 43-54.	1.5	7
151	Comparative study of the haplotype structure and linkage disequilibrium of chromosome 1p36.2 region in the Korean and Japanese populations. <i>Journal of Human Genetics</i> , 2004, 49, 603-609.	2.3	7
152	Association of UBE2L3 polymorphisms with diffuse cutaneous systemic sclerosis in a Japanese population: Table 1. <i>Annals of the Rheumatic Diseases</i> , 2012, 71, 1259-1260.	0.9	7
153	Association of Functional Polymorphisms in Interferon Regulatory Factor 2 (IRF2) with Susceptibility to Systemic Lupus Erythematosus: A Case-Control Association Study. <i>PLoS ONE</i> , 2014, 9, e109764.	2.5	7
154	Identification of Genes Upregulated in the Inflamed Colonic Lesions of Crohn's Disease. <i>Biochemical and Biophysical Research Communications</i> , 2001, 283, 130-135.	2.1	6
155	Identification of a novel HLA allele, HLA-DQB1*06:51, in a Japanese rheumatoid arthritis patient. <i>Tissue Antigens</i> , 2012, 80, 386-387.	1.0	6
156	HLA-DQB1 DPB1 alleles in Japanese patients with adult-onset Still's disease. <i>Modern Rheumatology</i> , 2019, 29, 843-847.	1.8	6
157	Human leukocyte antigen in Japanese patients with idiopathic inflammatory myopathy. <i>Modern Rheumatology</i> , 2020, 30, 696-702.	1.8	6
158	Modulation of methotrexate-induced intestinal mucosal injury by dietary factors. <i>Human and Experimental Toxicology</i> , 2020, 39, 500-513.	2.2	6
159	MICA allele typing of HLA-B27 positive Japanese patients with seronegative spondylarthropathies and healthy individuals: Differential linkage disequilibrium with HLA-B27 subtypes. <i>Arthritis and Rheumatism</i> , 1998, 41, 68-73.	6.7	6
160	Association of functional (GA) _n microsatellite polymorphism in the FLI1 gene with susceptibility to human systemic sclerosis. <i>Rheumatology</i> , 2020, 59, 3553-3562.	1.9	5
161	Molecular Mimicry, Ankylosing Spondylitis and Reactive Arthritis -Something Missing?. <i>Scandinavian Journal of Rheumatology</i> , 1992, 21, 105-108.	1.1	4
162	Genes Highly Expressed in the Early Phase of Murine Graft-versus-Host Reaction. <i>Biochemical and Biophysical Research Communications</i> , 2001, 282, 200-206.	2.1	4

#	ARTICLE	IF	CITATIONS
163	A novel HLA-DQB1*04 allele, DQB1*04:10, identified in a Japanese individual. <i>Tissue Antigens</i> , 2013, 82, 148-149.	1.0	4
164	Genetics of Microscopic Polyangiitis in the Japanese Population. <i>Annals of Vascular Diseases</i> , 2012, 5, 289-295.	0.5	3
165	Human immune system diversity and its implications in diseases. <i>Journal of Human Genetics</i> , 2015, 60, 655-656.	2.3	3
166	Biomarker for nontuberculous mycobacterial pulmonary disease in patients with rheumatoid arthritis: Anti-glycopeptidolipid core antigen immunoglobulin A antibodies. <i>Modern Rheumatology</i> , 2018, 28, 271-275.	1.8	3
167	Combining effects of polymorphism of tumor necrosis factor \pm 5'-flanking region and HLA-DRB1 on radiological progression in patients with rheumatoid arthritis. <i>Modern Rheumatology</i> , 2009, 19, 134-139.	1.8	3
168	Use of Lectin for Detection of Agalactosyl IgG. , 1998, 9, 195-206.		2
169	Identification of the gene variations in human IKKA. <i>Immunogenetics</i> , 1999, 50, 363-365.	2.4	2
170	HLA-DRB1 alleles encoding the shared epitope associated with rheumatoid arthritis confer additional susceptibility to seronegative spondyloarthropathies in HLA-B27-positive Japanese individuals. <i>Japanese Journal of Rheumatology</i> , 1999, 9, 55-64.	0.0	2
171	Genetics of Systemic Sclerosis. , 2016, , 81-92.		2
172	Successful Catheter Intervention for Acute Coronary Syndrome in a Patient with Antiphospholipid Syndrome.. <i>International Heart Journal</i> , 2001, 42, 627-631.	0.6	2
173	Synovial Expression of HLA-B27 Related Antigens in Ankylosing Spodilitis. <i>Scandinavian Journal of Rheumatology</i> , 1988, 17, 23-25.	1.1	1
174	Possible Relationship between Bacterial or Viral Fc γ -Binding Proteins and Rheumatoid Factor. <i>ImmunoMethods</i> , 1993, 2, 65-70.	0.8	1
175	Association of CYP17 with HLA-B27-negative seronegative spondyloarthropathy in Japanese males. <i>American Journal of Medical Genetics Part A</i> , 2004, 130A, 169-171.	2.4	1
176	Combining effects of polymorphism of tumor necrosis factor \pm 5'-flanking region and HLA-DRB1 on radiological progression in patients with rheumatoid arthritis. <i>Modern Rheumatology</i> , 2009, 19, 134-139.	1.8	1
177	Response to: \sim HLA-A* 31:01 is not associated with the development of methotrexate pneumonitis in the UK population: results from a genome wide association study \hat{e} ™ by Bluett et al. <i>Annals of the Rheumatic Diseases</i> , 2017, 76, e52-e52.	0.9	1
178	Enlargement of multiple cavernous hemangioma of the liver in patients with systemic lupus erythematosus: a report of four cases. <i>Japanese Journal of Rheumatology</i> , 1998, 8, 445-452.	0.0	0
179	Enlargement of multiple cavernous hemangioma of the liver in patients with systemic lupus erythematosus: a report of four cases. <i>Japanese Journal of Rheumatology</i> , 1998, 8, 445-452.	0.0	0
180	HLA-DRB1 alleles encoding the shared epitope associated with rheumatoid arthritis confer additional susceptibility to seronegative spondyloarthropathies in HLA-B27-positive Japanese individuals. <i>Japanese Journal of Rheumatology</i> , 1999, 9, 55-64.	0.0	0

#	ARTICLE	IF	CITATIONS
181	The human histocompatibility leukocyte antigen (HLA) haplotype is associated with the onset of postherpetic neuralgia after herpes zoster. <i>Pain</i> , 2004, 110, 329-329.	4.2	0
182	OP0120 Specific Identification of Anti-Citrullinated Glucose-6-Phosphate Isomerase Peptide (CCG) Antibodies Associated with HLA-DRB1 SE and Disease Activity in Patients with RA. <i>Annals of the Rheumatic Diseases</i> , 2013, 72, A92.1-A92.	0.9	0
183	HLA and autoimmune rheumatic diseases: association studies in Japan and recent progress in research. <i>Major Histocompatibility Complex</i> , 2015, 22, 74-83.	0.1	0
184	P2_17 Association of TNFSF4 Polymorphism with Proteinase 3 - ANCA Positive Vasculitis in a Japanese Population. <i>Rheumatology</i> , 2017, 56, iii109-iii110.	1.9	0
185	HLA-DRB1 and FCGR2B: highlights of the first genome-wide association study of IgG4-related disease. <i>Lancet Rheumatology</i> , The, 2019, 1, e2-e3.	3.9	0
186	Role of $IRF5$, $STAT4$ and BLK polymorphisms for the genetic predisposition to systemic lupus erythematosus in Japanese. <i>Inflammation and Regeneration</i> , 2009, 29, 190-197.	3.7	0
187	Dietary Factors Modulate Gastrointestinal Adverse Effects of Methotrexate. <i>Proceedings for Annual Meeting of the Japanese Pharmacological Society</i> , 2018, WCP2018, PO3-13-22.	0.0	0
188	Diagnostic significance of fluorocytometric DNA ploidy analysis of gastric cancer with special reference to its comparison with clinical and histopathological diagnosis. <i>Nihon Geka Hokan Archiv F&A/r Japanische Chirurgie</i> , 1990, 59, 116-33.	0.0	0
189	Expression of membrane-type matrix metalloproteinases in synovial tissue from patients with rheumatoid arthritis or osteoarthritis. <i>Modern Rheumatology</i> , 2001, 11, 34-39.	1.8	0