

Naoyuki Tsuchiya

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/5638743/publications.pdf>

Version: 2024-02-01

189
papers

7,169
citations

50276

46
h-index

74163

75
g-index

194
all docs

194
docs citations

194
times ranked

7119
citing authors

#	ARTICLE	IF	CITATIONS
1	Advances in the genomics of ANCA-associated vasculitis—a view from East Asia. <i>Genes and Immunity</i> , 2021, 22, 1-11.	4.1	12
2	Human leukocyte antigen in Japanese patients with idiopathic inflammatory myopathy. <i>Modern Rheumatology</i> , 2020, 30, 696-702.	1.8	6
3	Modulation of methotrexate-induced intestinal mucosal injury by dietary factors. <i>Human and Experimental Toxicology</i> , 2020, 39, 500-513.	2.2	6
4	Association of functional (GA) _n microsatellite polymorphism in the FLI1 gene with susceptibility to human systemic sclerosis. <i>Rheumatology</i> , 2020, 59, 3553-3562.	1.9	5
5	Association of TERT and DSP variants with microscopic polyangiitis and myeloperoxidase-ANCA positive vasculitis in a Japanese population: a genetic association study. <i>Arthritis Research and Therapy</i> , 2020, 22, 246.	3.5	10
6	HLA-DQB1 DPB1 alleles in Japanese patients with adult-onset Still's disease. <i>Modern Rheumatology</i> , 2019, 29, 843-847.	1.8	6
7	Association of NCF1 polymorphism with systemic lupus erythematosus and systemic sclerosis but not with ANCA-associated vasculitis in a Japanese population. <i>Scientific Reports</i> , 2019, 9, 16366.	3.3	15
8	HLA-DRB1 and FCGR2B: highlights of the first genome-wide association study of IgG4-related disease. <i>Lancet Rheumatology</i> , The, 2019, 1, e2-e3.	3.9	0
9	Association of <i>MUC5B</i> promoter polymorphism with interstitial lung disease in myeloperoxidase-antineutrophil cytoplasmic antibody-associated vasculitis. <i>Annals of the Rheumatic Diseases</i> , 2019, 78, 1144-1146.	0.9	23
10	Association of HLA-DRB1 genotype with younger age onset and elder age onset rheumatoid arthritis in Japanese populations. <i>Medicine (United States)</i> , 2019, 98, e18218.	1.0	10
11	Association of a single nucleotide polymorphism in TNIP1 with type-1 autoimmune hepatitis in the Japanese population. <i>Journal of Human Genetics</i> , 2018, 63, 739-744.	2.3	9
12	Biomarker for nontuberculous mycobacterial pulmonary disease in patients with rheumatoid arthritis: Anti-glycopeptidolipid core antigen immunoglobulin A antibodies. <i>Modern Rheumatology</i> , 2018, 28, 271-275.	1.8	3
13	Association of ETS1 polymorphism with granulomatosis with polyangiitis and proteinase 3-anti-neutrophil cytoplasmic antibody positive vasculitis in a Japanese population. <i>Journal of Human Genetics</i> , 2018, 63, 55-62.	2.3	14
14	<i>MUC5B</i> Promoter Variant and Rheumatoid Arthritis with Interstitial Lung Disease. <i>New England Journal of Medicine</i> , 2018, 379, 2209-2219.	27.0	326
15	Dietary Factors Modulate Gastrointestinal Adverse Effects of Methotrexate. <i>Proceedings for Annual Meeting of the Japanese Pharmacological Society</i> , 2018, WCP2018, PO3-13-22.	0.0	0
16	P2_17 Association of TNFSF4 Polymorphism with Proteinase 3 - ANCA Positive Vasculitis in a Japanese Population. <i>Rheumatology</i> , 2017, 56, iii109-iii110.	1.9	0
17	Response to: HLA-A* 31:01 is not associated with the development of methotrexate pneumonitis in the UK population: results from a genome wide association study™ by Bluett et al. <i>Annals of the Rheumatic Diseases</i> , 2017, 76, e52-e52.	0.9	1
18	Transethnic meta-analysis identifies <i>GSDMA</i> and <i>PRDM1</i> as susceptibility genes to systemic sclerosis. <i>Annals of the Rheumatic Diseases</i> , 2017, 76, 1150-1158.	0.9	77

#	ARTICLE	IF	CITATIONS
19	Association of a single nucleotide polymorphism upstream of ICOS with Japanese autoimmune hepatitis type 1. <i>Journal of Human Genetics</i> , 2017, 62, 481-484.	2.3	10
20	Plasma miRNA expression profiles in rheumatoid arthritis associated interstitial lung disease. <i>BMC Musculoskeletal Disorders</i> , 2017, 18, 21.	1.9	29
21	The role of common protective alleles HLA-DRB1*13 among systemic autoimmune diseases. <i>Genes and Immunity</i> , 2017, 18, 1-7.	4.1	44
22	HLA-DRB1 and DQB1 alleles in Japanese type 1 autoimmune hepatitis: The predisposing role of the DR4/DR8 heterozygous genotype. <i>PLoS ONE</i> , 2017, 12, e0187325.	2.5	26
23	Effects of HLA-DRB1 alleles on susceptibility and clinical manifestations in Japanese patients with adult onset Still's disease. <i>Arthritis Research and Therapy</i> , 2017, 19, 199.	3.5	25
24	Association of HLA-G 3' Untranslated Region Polymorphisms with Systemic Lupus Erythematosus in a Japanese Population: A Case-Control Association Study. <i>PLoS ONE</i> , 2016, 11, e0158065.	2.5	19
25	Association of BAK1 single nucleotide polymorphism with a risk for dengue hemorrhagic fever. <i>BMC Medical Genetics</i> , 2016, 17, 43.	2.1	8
26	Association of human leukocyte antigen alleles with chronic lung diseases in rheumatoid arthritis. <i>Rheumatology</i> , 2016, 55, 1301-1307.	1.9	29
27	The pattern of GPI-80 expression is a useful marker for unusual myeloid maturation in peripheral blood. <i>Clinical and Experimental Immunology</i> , 2016, 186, 373-386.	2.6	8
28	Genetics of Systemic Sclerosis. , 2016, , 81-92.		2
29	Human Leukocyte Antigen and Systemic Sclerosis in Japanese: The Sign of the Four Independent Protective Alleles, DRB1*13:02, DRB1*14:06, DQB1*03:01, and DPB1*02:01. <i>PLoS ONE</i> , 2016, 11, e0154255.	2.5	25
30	Protective Role of HLA-DRB1*13:02 against Microscopic Polyangiitis and MPO-ANCA-Positive Vasculitides in a Japanese Population: A Case-Control Study. <i>PLoS ONE</i> , 2016, 11, e0154393.	2.5	35
31	Genetics of Interstitial Lung Disease: <i>Vol de Nuit</i> (Night Flight). <i>Clinical Medicine Insights: Circulatory, Respiratory and Pulmonary Medicine</i> , 2015, 9s1, CCRPM.S23283.	0.9	16
32	HLA and autoimmune rheumatic diseases: association studies in Japan and recent progress in research. <i>Major Histocompatibility Complex</i> , 2015, 22, 74-83.	0.1	0
33	Autoantibody Profiles in Collagen Disease Patients with Interstitial Lung Disease (ILD): Antibodies to Major Histocompatibility Complex Class I-Related Chain a (MICA) as Markers of ILD. <i>Biomarker Insights</i> , 2015, 10, BMI.S28209.	2.5	10
34	Human immune system diversity and its implications in diseases. <i>Journal of Human Genetics</i> , 2015, 60, 655-656.	2.3	3
35	Identification of secreted phosphoprotein 1 gene as a new rheumatoid arthritis susceptibility gene. <i>Annals of the Rheumatic Diseases</i> , 2015, 74, e19-e19.	0.9	24
36	Protective Effect of the HLA-DRB1*13:02 Allele in Japanese Rheumatoid Arthritis Patients. <i>PLoS ONE</i> , 2014, 9, e99453.	2.5	60

#	ARTICLE	IF	CITATIONS
37	Association of Functional Polymorphisms in Interferon Regulatory Factor 2 (IRF2) with Susceptibility to Systemic Lupus Erythematosus: A Case-Control Association Study. <i>PLoS ONE</i> , 2014, 9, e109764.	2.5	7
38	A replication study confirms the association of GWAS-identified SNPs at MICB and PLCE1 in Thai patients with dengue shock syndrome. <i>BMC Medical Genetics</i> , 2014, 15, 58.	2.1	26
39	HLA-DRB1*08:02 Is Associated with Bucillamine-Induced Proteinuria in Japanese Rheumatoid Arthritis Patients. <i>Biomarker Insights</i> , 2014, 9, BMI.S13654.	2.5	9
40	Human Leukocyte Antigens and Systemic Lupus Erythematosus: A Protective Role for the HLA-DR6 Alleles DRB1*13:02 and *14:03. <i>PLoS ONE</i> , 2014, 9, e87792.	2.5	50
41	Serum biomarker analysis of collagen disease patients with acute-onset diffuse interstitial lung disease. <i>BMC Immunology</i> , 2013, 14, 9.	2.2	26
42	Association of IRF5 polymorphism with MPO-ANCA-positive vasculitis in a Japanese population. <i>Genes and Immunity</i> , 2013, 14, 527-529.	4.1	10
43	Genome, epigenome and transcriptome analyses of a pair of monozygotic twins discordant for systemic lupus erythematosus. <i>Human Immunology</i> , 2013, 74, 170-175.	2.4	51
44	Genetics of ANCA-associated vasculitis in Japan: a role for HLA-DRB1*09:01 haplotype. <i>Clinical and Experimental Nephrology</i> , 2013, 17, 628-630.	1.6	23
45	<i>PLD4</i> as a novel susceptibility gene for systemic sclerosis in a Japanese population. <i>Arthritis and Rheumatism</i> , 2013, 65, 472-480.	6.7	62
46	HLA-A*31:01 and methotrexate-induced interstitial lung disease in Japanese rheumatoid arthritis patients: a multidrug hypersensitivity marker? <i>Table A1. Annals of the Rheumatic Diseases</i> , 2013, 72, 153-155.	0.9	43
47	Association of a single nucleotide polymorphism in the <i>SH2D1A</i> intronic region with systemic lupus erythematosus. <i>Lupus</i> , 2013, 22, 497-503.	1.6	11
48	Anti-citrullinated glucose-6-phosphate isomerase peptide antibodies in patients with rheumatoid arthritis are associated with <i>HLA-DRB1</i> shared epitope alleles and disease activity. <i>Clinical and Experimental Immunology</i> , 2013, 172, 44-53.	2.6	17
49	A novel <i>HLA-DQB1*04</i> allele, <i>DQB1*04:10</i> , identified in a Japanese individual. <i>Tissue Antigens</i> , 2013, 82, 148-149.	1.0	4
50	A functional SNP upstream of the beta-2 adrenergic receptor gene (<i>ADRB2</i>) is associated with obesity in Oceanic populations. <i>International Journal of Obesity</i> , 2013, 37, 1204-1210.	3.4	10
51	OP0120...Specific Identification of Anti-Citrullinated Glucose-6-Phosphate Isomerase Peptide (CCG) Antibodies Associated with HLA-DRB1 SE and Disease Activity in Patients with RA. <i>Annals of the Rheumatic Diseases</i> , 2013, 72, A92.1-A92.	0.9	0
52	Association of Increased Frequencies of HLA-DPB1*05:01 with the Presence of Anti-Ro/SS-A and Anti-La/SS-B Antibodies in Japanese Rheumatoid Arthritis and Systemic Lupus Erythematosus Patients. <i>PLoS ONE</i> , 2013, 8, e53910.	2.5	21
53	Epistatic Interaction between BANK1 and BLK in Rheumatoid Arthritis: Results from a Large Trans-Ethnic Meta-Analysis. <i>PLoS ONE</i> , 2013, 8, e61044.	2.5	24
54	Association of <i>PHRF1-IRF7</i> region polymorphism with clinical manifestations of systemic lupus erythematosus in a Japanese population. <i>Lupus</i> , 2012, 21, 890-895.	1.6	18

#	ARTICLE	IF	CITATIONS
55	Association of UBE2L3 polymorphisms with diffuse cutaneous systemic sclerosis in a Japanese population: Table 1. <i>Annals of the Rheumatic Diseases</i> , 2012, 71, 1259-1260.	0.9	7
56	Effects of APRIL (TNFSF13) polymorphisms and splicing isoforms on the secretion of soluble APRIL. <i>Modern Rheumatology</i> , 2012, 22, 541-549.	1.8	15
57	Human CD72 splicing isoform responsible for resistance to systemic lupus erythematosus regulates serum immunoglobulin level and is localized in endoplasmic reticulum. <i>BMC Immunology</i> , 2012, 13, 72.	2.2	13
58	Association of Human Leukocyte Antigen with Interstitial Lung Disease in Rheumatoid Arthritis: A Protective Role for Shared Epitope. <i>PLoS ONE</i> , 2012, 7, e33133.	2.5	70
59	Genetics of Microscopic Polyangiitis in the Japanese Population. <i>Annals of Vascular Diseases</i> , 2012, 5, 289-295.	0.5	3
60	Identification of a novel HLA allele, HLA-DQB1*06:51, in a Japanese rheumatoid arthritis patient. <i>Tissue Antigens</i> , 2012, 80, 386-387.	1.0	6
61	Effects of APRIL (TNFSF13) polymorphisms and splicing isoforms on the secretion of soluble APRIL. <i>Modern Rheumatology</i> , 2012, 22, 541-549.	1.8	9
62	TLR7 single-nucleotide polymorphisms in the 3' untranslated region and intron 2 independently contribute to systemic lupus erythematosus in Japanese women: a case-control association study. <i>Arthritis Research and Therapy</i> , 2011, 13, R41.	3.5	93
63	Association of ADAMTS13 polymorphism with cerebral malaria. <i>Malaria Journal</i> , 2011, 10, 366.	2.3	19
64	Association of a functional polymorphism in the 3'-untranslated region of SPI1 with systemic lupus erythematosus. <i>Arthritis and Rheumatism</i> , 2011, 63, 755-763.	6.7	50
65	The Impact of Natural Selection on an ABCC11 SNP Determining Earwax Type. <i>Molecular Biology and Evolution</i> , 2011, 28, 849-857.	8.9	44
66	Cumulative association of eight susceptibility genes with systemic lupus erythematosus in a Japanese female population. <i>Journal of Human Genetics</i> , 2011, 56, 503-507.	2.3	35
67	The Q223R polymorphism in LEPR is associated with obesity in Pacific Islanders. <i>Human Genetics</i> , 2010, 127, 287-294.	3.8	74
68	Association of the FAM167A-BLK region with systemic sclerosis. <i>Arthritis and Rheumatism</i> , 2010, 62, 890-895.	6.7	76
69	Association of IRF5, STAT4 and BLK with systemic lupus erythematosus and other rheumatic diseases.. <i>Japanese Journal of Clinical Immunology</i> , 2010, 33, 57-65.	0.0	16
70	Sex-specific association of X-linked Toll-like receptor 7 (TLR7) with male systemic lupus erythematosus. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2010, 107, 15838-15843.	7.1	324
71	Association of TNFAIP3 Polymorphism with Susceptibility to Systemic Lupus Erythematosus in a Japanese Population. <i>Journal of Biomedicine and Biotechnology</i> , 2010, 2010, 1-5.	3.0	31
72	Replication of association between FAM167A(C8orf13)-BLK region and rheumatoid arthritis in a Japanese population. <i>Annals of the Rheumatic Diseases</i> , 2010, 69, 936-937.	0.9	24

#	ARTICLE	IF	CITATIONS
73	Association of TNFAIP3 interacting protein 1, TNIP1 with systemic lupus erythematosus in a Japanese population: a case-control association study. <i>Arthritis Research and Therapy</i> , 2010, 12, R174.	3.5	70
74	Association of STAT4 polymorphism with systemic sclerosis in a Japanese population. <i>Annals of the Rheumatic Diseases</i> , 2009, 68, 1375-1376.	0.9	64
75	Replication of the association between the C8orf13-BLK region and systemic lupus erythematosus in a Japanese population. <i>Arthritis and Rheumatism</i> , 2009, 60, 553-558.	6.7	57
76	Association of a functional polymorphism in the IRF5 region with systemic sclerosis in a Japanese population. <i>Arthritis and Rheumatism</i> , 2009, 60, 1845-1850.	6.7	115
77	IFNGR1 polymorphisms in Thai malaria patients. <i>Infection, Genetics and Evolution</i> , 2009, 9, 1406-1409.	2.3	10
78	A replication study of the association between the IL12B promoter allele CTCTAA and susceptibility to cerebral malaria in Thai population. <i>Malaria Journal</i> , 2009, 8, 290.	2.3	10
79	Identification of a haplotype block in the 5q31 cytokine gene cluster associated with the susceptibility to severe malaria. <i>Malaria Journal</i> , 2009, 8, 232.	2.3	20
80	Association study of a polymorphism of the CTGF gene and susceptibility to systemic sclerosis in the Japanese population. <i>Annals of the Rheumatic Diseases</i> , 2009, 68, 1921-1924.	0.9	42
81	Combining effects of polymorphism of tumor necrosis factor \pm 5'-flanking region and HLA-DRB1 on radiological progression in patients with rheumatoid arthritis. <i>Modern Rheumatology</i> , 2009, 19, 134-139.	1.8	1
82	Role of IRF5, STAT4 and BLK polymorphisms for the genetic predisposition to systemic lupus erythematosus in Japanese. <i>Inflammation and Regeneration</i> , 2009, 29, 190-197.	3.7	0
83	Combining effects of polymorphism of tumor necrosis factor \pm 5'-flanking region and HLA-DRB1 on radiological progression in patients with rheumatoid arthritis. <i>Modern Rheumatology</i> , 2009, 19, 134-139.	1.8	3
84	Association of IRF5 polymorphisms with systemic lupus erythematosus in a Japanese population: Support for a crucial role of intron 1 polymorphisms. <i>Arthritis and Rheumatism</i> , 2008, 58, 826-834.	6.7	100
85	Association of LILRA2 (ILT1, LIR7) splice site polymorphism with systemic lupus erythematosus and microscopic polyangiitis. <i>Genes and Immunity</i> , 2008, 9, 214-223.	4.1	30
86	Antibodies to the peptide from the plasmid-coded Yersinia outer membrane protein (YOP1) in patients with ankylosing spondylitis. <i>Clinical and Experimental Immunology</i> , 2008, 82, 493-498.	2.6	11
87	Role of STAT4 polymorphisms in systemic lupus erythematosus in a Japanese population: a case-control association study of the STAT1-STAT4 region. <i>Arthritis Research and Therapy</i> , 2008, 10, R113.	3.5	88
88	Association of IL-10 receptor 2 (IL10RB) SNP with systemic sclerosis. <i>Biochemical and Biophysical Research Communications</i> , 2008, 373, 403-407.	2.1	35
89	Diversity of Human Immune System Multigene Families and its Implication in the Genetic Background of Rheumatic Diseases. <i>Current Medicinal Chemistry</i> , 2007, 14, 431-439.	2.4	17
90	Role of APRIL (TNFSF13) polymorphisms in the susceptibility to systemic lupus erythematosus in Japanese. <i>Rheumatology</i> , 2007, 46, 776-782.	1.9	22

#	ARTICLE	IF	CITATIONS
91	A compass that points to lupus: genetic studies on type I interferon pathway. <i>Genes and Immunity</i> , 2007, 8, 445-455.	4.1	84
92	Association of CD22 gene polymorphism with susceptibility to limited cutaneous systemic sclerosis. <i>Tissue Antigens</i> , 2007, 69, 242-249.	1.0	38
93	Differential association of HLA-DRB1 alleles in Japanese patients with early rheumatoid arthritis in relationship to autoantibodies to cyclic citrullinated peptide. <i>Clinical and Experimental Rheumatology</i> , 2007, 25, 219-24.	0.8	32
94	Effects of down-regulating the Id genes in human colorectal cancer cells on early steps of haematogenous metastasis. <i>European Journal of Cancer</i> , 2006, 42, 668-673.	2.8	8
95	Association of HLA-DRB1*0901-DQB1*0303 haplotype with microscopic polyangiitis in Japanese. <i>Genes and Immunity</i> , 2006, 7, 81-84.	4.1	49
96	Role of B cell inhibitory receptor polymorphisms in systemic lupus erythematosus: a negative times a negative makes a positive. <i>Journal of Human Genetics</i> , 2006, 51, 741-750.	2.3	18
97	Association of killer cell immunoglobulin-like receptor genotypes with microscopic polyangiitis. <i>Arthritis and Rheumatism</i> , 2006, 54, 992-997.	6.7	39
98	Targeting Id1 and Id3 inhibits peritoneal metastasis of gastric cancer. <i>Cancer Science</i> , 2005, 96, 784-790.	3.9	53
99	Evaluation of microsatellite markers in association studies: a search for an immune-related susceptibility gene in sarcoidosis. <i>Immunogenetics</i> , 2005, 56, 861-870.	2.4	12
100	Role of Fc γ 3 receptor IIb polymorphism in the genetic background of systemic lupus erythematosus: Insights from Asia. <i>Autoimmunity</i> , 2005, 38, 347-352.	2.6	32
101	Extensive polymorphisms of LILRB1 (ILT2, LIR1) and their association with HLA-DRB1 shared epitope negative rheumatoid arthritis. <i>Human Molecular Genetics</i> , 2005, 14, 2469-2480.	2.9	69
102	Fc γ 3RIIB Ile232Thr transmembrane polymorphism associated with human systemic lupus erythematosus decreases affinity to lipid rafts and attenuates inhibitory effects on B cell receptor signaling. <i>Human Molecular Genetics</i> , 2005, 14, 2881-2892.	2.9	216
103	Molecular genetic analyses of human NKG2C (KLRC2) gene deletion. <i>International Immunology</i> , 2004, 16, 163-168.	4.0	73
104	Crucial Role of Inhibitor of DNA Binding/Differentiation in the Vascular Endothelial Growth Factor-Induced Activation and Angiogenic Processes of Human Endothelial Cells. <i>Journal of Immunology</i> , 2004, 173, 5801-5809.	0.8	88
105	CD72 polymorphisms associated with alternative splicing modify susceptibility to human systemic lupus erythematosus through epistatic interaction with FCGR2B. <i>Human Molecular Genetics</i> , 2004, 13, 2907-2917.	2.9	43
106	Association of Fc γ 3 receptor IIb polymorphism with susceptibility to systemic lupus erythematosus in Chinese: a common susceptibility gene in the Asian populations. <i>Tissue Antigens</i> , 2004, 63, 21-27.	1.0	142
107	A novel method for isolation of endothelial cells and macrophages from murine tumors based on Ac-LDL uptake and CD16 expression. <i>Journal of Immunological Methods</i> , 2004, 295, 183-193.	1.4	10
108	Comparative study of the haplotype structure and linkage disequilibrium of chromosome 1p36.2 region in the Korean and Japanese populations. <i>Journal of Human Genetics</i> , 2004, 49, 603-609.	2.3	7

#	ARTICLE	IF	CITATIONS
109	Association of Fc γ receptor IIA, but not IIB and IIIA, polymorphisms with systemic lupus erythematosus: A family-based association study in Caucasians. <i>Arthritis and Rheumatism</i> , 2004, 50, 671-673.	6.7	34
110	Association of a functional CD19 polymorphism with susceptibility to systemic sclerosis. <i>Arthritis and Rheumatism</i> , 2004, 50, 4002-4007.	6.7	82
111	Association of CYP17 with HLA-B27-negative seronegative spondyloarthritis in Japanese males. <i>American Journal of Medical Genetics Part A</i> , 2004, 130A, 169-171.	2.4	1
112	The human histocompatibility leukocyte antigen (HLA) haplotype is associated with the onset of postherpetic neuralgia after herpes zoster. <i>Pain</i> , 2004, 110, 329-329.	4.2	0
113	The human histocompatibility leukocyte antigen (HLA) haplotype is associated with the onset of postherpetic neuralgia after herpes zoster. <i>Pain</i> , 2004, 110, 329-336.	4.2	33
114	BAFF/BLyS can potentiate B-cell selection with the B-cell coreceptor complex. <i>Blood</i> , 2004, 103, 2257-2265.	1.4	151
115	Immunogenetic features in 120 Japanese patients with idiopathic inflammatory myopathy. <i>Journal of Rheumatology</i> , 2004, 31, 1768-74.	2.0	44
116	Variations in the human Th2-specific chemokine TARC gene. <i>Immunogenetics</i> , 2003, 54, 742-745.	2.4	29
117	TNFR2 position 196 polymorphism in Japanese patients with rheumatoid arthritis: Comment on the article by Dieudonné et al. <i>Arthritis and Rheumatism</i> , 2003, 48, 273-274.	6.7	9
118	Role of the Fc γ 3 receptor IIA polymorphism in the antiphospholipid syndrome: An international meta-analysis. <i>Arthritis and Rheumatism</i> , 2003, 48, 1930-1938.	6.7	49
119	Association of Fc γ 3 receptor IIb and IIIb polymorphisms with susceptibility to systemic lupus erythematosus in Thais. <i>Tissue Antigens</i> , 2003, 61, 374-383.	1.0	146
120	Variations of human killer cell lectin-like receptors: common occurrence of NKG2-C deletion in the general population. <i>Genes and Immunity</i> , 2003, 4, 160-167.	4.1	51
121	Exacerbation of Lambert-Eaton Myasthenic Syndrome Caused by an L-type Ca ²⁺ -Channel Antagonist.. <i>International Heart Journal</i> , 2003, 44, 139-144.	0.6	8
122	Genetic background of Japanese patients with antineutrophil cytoplasmic antibody-associated vasculitis: association of HLA-DRB1*0901 with microscopic polyangiitis. <i>Journal of Rheumatology</i> , 2003, 30, 1534-40.	2.0	70
123	Fc γ receptor gene polymorphisms in Japanese patients with systemic lupus erythematosus: Contribution of FCGR2B to genetic susceptibility. <i>Arthritis and Rheumatism</i> , 2002, 46, 1242-1254.	6.7	301
124	Association of HLA-DRB1*1502 -DQB1*0501 haplotype with susceptibility to systemic lupus erythematosus in Thais. <i>Tissue Antigens</i> , 2002, 59, 113-117.	1.0	40
125	Variations in immune response genes and their associations with multifactorial immune disorders. <i>Immunological Reviews</i> , 2002, 190, 169-181.	6.0	45
126	Association of HLA-A*3303-B*4403-DRB1*1302 haplotype, but not of TNFA promoter and NKp30 polymorphism, with postherpetic neuralgia (PHN) in the Japanese population. <i>Genes and Immunity</i> , 2002, 3, 477-481.	4.1	42

#	ARTICLE	IF	CITATIONS
127	Polymorphisms of human CD19 gene: possible association with susceptibility to systemic lupus erythematosus in Japanese. <i>Genes and Immunity</i> , 2002, 3, S21-S30.	4.1	26
128	Studies on the association of Fc γ 3 receptor IIA, IIB, IIIA and IIIB polymorphisms with rheumatoid arthritis in the Japanese: evidence for a genetic interaction between HLA-DRB1 and FCGR3A. <i>Genes and Immunity</i> , 2002, 3, 488-493.	4.1	62
129	Analysis on the association of human BLYS (BAFF, TNFSF13B) polymorphisms with systemic lupus erythematosus and rheumatoid arthritis. <i>Genes and Immunity</i> , 2002, 3, 424-429.	4.1	100
130	Genes Highly Expressed in the Early Phase of Murine Graft-versus-Host Reaction. <i>Biochemical and Biophysical Research Communications</i> , 2001, 282, 200-206.	2.1	4
131	Identification of Genes Upregulated in the Inflamed Colonic Lesions of Crohn's Disease. <i>Biochemical and Biophysical Research Communications</i> , 2001, 283, 130-135.	2.1	6
132	Expression of ID Family Genes in the Synovia from Patients with Rheumatoid Arthritis. <i>Biochemical and Biophysical Research Communications</i> , 2001, 284, 436-442.	2.1	32
133	Comparison of statistical power between 2x2 allele frequency and allele positivity tables in case-control studies of complex disease genes. <i>Annals of Human Genetics</i> , 2001, 65, 197-206.	0.8	91
134	Identification of novel single nucleotide substitutions in the NKp30 gene expressed in human natural killer cells. <i>Tissue Antigens</i> , 2001, 58, 255-258.	1.0	11
135	Presence of four major haplotypes in human BCMA gene: lack of association with systemic lupus erythematosus and rheumatoid arthritis. <i>Genes and Immunity</i> , 2001, 2, 276-279.	4.1	24
136	Analysis of the association of HLA-DRB1, TNF β promoter and TNFR2 (TNFRSF1B) polymorphisms with SLE using transmission disequilibrium test. <i>Genes and Immunity</i> , 2001, 2, 317-322.	4.1	69
137	Variations in the human CC chemokine eotaxin gene. <i>Genes and Immunity</i> , 2001, 2, 461-463.	4.1	24
138	Comparison of statistical power between 2 \times 2 allele frequency and allele positivity tables in case-control studies of complex disease genes. <i>Annals of Human Genetics</i> , 2001, 65, 197-206.	0.8	93
139	Successful Catheter Intervention for Acute Coronary Syndrome in a Patient with Antiphospholipid Syndrome.. <i>International Heart Journal</i> , 2001, 42, 627-631.	0.6	2
140	Expression of membrane-type matrix metalloproteinases in synovial tissue from patients with rheumatoid arthritis or osteoarthritis. <i>Modern Rheumatology</i> , 2001, 11, 34-39.	1.8	0
141	Tumor necrosis factor β 5' flanking region, tumor necrosis factor receptor II, and HLA-DRB1 polymorphisms in Japanese patients with rheumatoid arthritis. <i>Arthritis and Rheumatism</i> , 2000, 43, 753.	6.7	56
142	Single nucleotide polymorphisms in the coding regions of human CXC-chemokine receptors CXCR1, CXCR2 and CXCR3. <i>Genes and Immunity</i> , 2000, 1, 330-337.	4.1	29
143	Independent contribution of HLA-DRB1 and TNF β promoter polymorphisms to the susceptibility to Crohn's disease. <i>Genes and Immunity</i> , 2000, 1, 351-357.	4.1	48
144	New single nucleotide polymorphisms in the coding region of human TNFR2: association with systemic lupus erythematosus. <i>Genes and Immunity</i> , 2000, 1, 501-503.	4.1	29

#	ARTICLE	IF	CITATIONS
145	New polymorphisms of human CD80 and CD86: lack of association with rheumatoid arthritis and systemic lupus erythematosus. <i>Genes and Immunity</i> , 2000, 1, 428-434.	4.1	43
146	New variations in human OX40 ligand (CD134L) gene. <i>Genes and Immunity</i> , 2000, 1, 521-522.	4.1	11
147	HLA-DRB1 alleles encoding the shared epitope associated with rheumatoid arthritis confer additional susceptibility to seronegative spondyloarthropathies in HLA-B27-positive Japanese individuals. <i>Japanese Journal of Rheumatology</i> , 1999, 9, 55-64.	0.0	0
148	Significant association of HLA-B and HLA-DRB1 alleles with cleft lip with or without cleft palate. <i>Tissue Antigens</i> , 1999, 53, 147-152.	1.0	8
149	Association of tumor necrosis factor receptor 2 (TNFR2) polymorphism with susceptibility to systemic lupus erythematosus. <i>Tissue Antigens</i> , 1999, 53, 527-533.	1.0	127
150	Allele typing of human TNFA 5' flanking region using polymerase chain reaction-preferential homoduplex formation assay (PCR-PHFA): linkage disequilibrium with HLA class I and class II genes in Japanese. <i>Tissue Antigens</i> , 1999, 54, 478-484.	1.0	38
151	Lack of a strong association of CTLA-4 exon 1 polymorphism with the susceptibility to rheumatoid arthritis and systemic lupus erythematosus in Japanese: an association study using a novel variation screening method. <i>Tissue Antigens</i> , 1999, 54, 578-584.	1.0	57
152	New variations of human CC-chemokine receptors CCR3 and CCR4. <i>Genes and Immunity</i> , 1999, 1, 97-104.	4.1	23
153	Association of Fcγ3 receptor IIIB, but not of Fcγ3 receptor IIA and IIIA, polymorphisms with systemic lupus erythematosus in Japanese. <i>Genes and Immunity</i> , 1999, 1, 53-60.	4.1	121
154	Identification of the gene variations in human CD22. <i>Immunogenetics</i> , 1999, 49, 280-286.	2.4	68
155	New variations of human SHP-1. <i>Immunogenetics</i> , 1999, 49, 577-579.	2.4	14
156	Identification of the gene variations in human IKKA. <i>Immunogenetics</i> , 1999, 50, 363-365.	2.4	2
157	MIC-A polymorphism in Japanese and a MIC-A-MIC-B null haplotype. <i>Immunogenetics</i> , 1999, 49, 620-628.	2.4	82
158	HLA-DRB1 alleles encoding the shared epitope associated with rheumatoid arthritis confer additional susceptibility to seronegative spondyloarthropathies in HLA-B27-positive Japanese individuals. <i>Japanese Journal of Rheumatology</i> , 1999, 9, 55-64.	0.0	2
159	Overlapping peptide-binding specificities of HLA-B27 and B39: Evidence for a role of peptide supermotif in the pathogenesis of spondylarthropathies. <i>Arthritis and Rheumatism</i> , 1999, 42, 175-181.	6.7	36
160	Use of Lectin for Detection of Agalactosyl IgG. , 1998, 9, 195-206.		2
161	MICA allele typing of HLA-B27 positive Japanese patients with seronegative spondylarthropathies and healthy individuals: Differential linkage disequilibrium with HLA-B27 subtypes. <i>Arthritis and Rheumatism</i> , 1998, 41, 68-73.	6.7	45
162	Enlargement of multiple cavernous hemangioma of the liver in patients with systemic lupus erythematosus: a report of four cases. <i>Japanese Journal of Rheumatology</i> , 1998, 8, 445-452.	0.0	0

#	ARTICLE	IF	CITATIONS
163	Enlargement of multiple cavernous hemangioma of the liver in patients with systemic lupus erythematosus: a report of four cases. Japanese Journal of Rheumatology, 1998, 8, 445-452.	0.0	0
164	MICA allele typing of HLA-B27 positive japanese patients with seronegative spondylarthropathies and healthy individuals: Differential linkage disequilibrium with HLA-B27 subtypes. Arthritis and Rheumatism, 1998, 41, 68-73.	6.7	6
165	Induction of alternative splicing of HLA-B27 by bacterial invasion. Arthritis and Rheumatism, 1997, 40, 694-703.	6.7	30
166	Immunologic Mechanisms in Common Rheumatologic Diseases. Clinical Orthopaedics and Related Research, 1996, 326, 43-54.	1.5	7
167	Elevated serum level of soluble HLA class I antigens in patients with systemic lupus erythematosus. Arthritis and Rheumatism, 1996, 39, 792-796.	6.7	45
168	HLA-B associations of HLA-B27 negative ankylosing spondylitis: Comment on the article by Yamaguchi et al. Arthritis and Rheumatism, 1996, 39, 1768-1769.	6.7	13
169	HLA-B27 subtypes in Japanese with seronegative spondyloarthropathies and healthy controls. Journal of Rheumatology, 1996, 23, 1189-93.	2.0	18
170	Association of hla-b39 with hla-b27-negative ankylosing spondylitis and pauciarticular juvenile rheumatoid arthritis in japanese patients. Arthritis and Rheumatism, 1995, 38, 1672-1677.	6.7	124
171	Elevated Cytokine Levels in Synovial Fluid of Rheumatoid Arthritis Correlates with the Presence of Cytomegalovirus Genome. Autoimmunity, 1994, 17, 333-337.	2.6	13
172	Distribution of Glycosylation Abnormality among Serum IgG Subclasses from Patients with Rheumatoid Arthritis. Clinical Immunology and Immunopathology, 1994, 70, 47-50.	2.0	22
173	Possible Relationship between Bacterial or Viral Fc γ 3-Binding Proteins and Rheumatoid Factor. ImmunoMethods, 1993, 2, 65-70.	0.8	1
174	Lambert-Eaton Myasthenic Syndrome Associated with Sjögren's Syndrome and Discoid Lupus Erythematosus. Scandinavian Journal of Rheumatology, 1993, 22, 302-304.	1.1	27
175	Antibodies to Human Cytomegalovirus 65-Kilodalton Fc Binding Protein in Rheumatoid Arthritis: Idiotypic Mimicry Hypothesis of Rheumatoid Factor Production. Autoimmunity, 1993, 15, 39-48.	2.6	12
176	Detection of glycosylation abnormality in rheumatoid IgG using N-acetylglucosamine-specific Psathyrella velutina lectin. Journal of Immunology, 1993, 151, 1137-46.	0.8	46
177	Molecular Mimicry, Ankylosing Spondylitis and Reactive Arthritis -Something Missing?. Scandinavian Journal of Rheumatology, 1992, 21, 105-108.	1.1	4
178	Molecular mimicry-hypothesis or reality?. Western Journal of Medicine, 1992, 157, 133-8.	0.3	11
179	Cytomegalovirus genomes demonstrated by polymerase chain reaction in synovial fluid from rheumatoid arthritis patients. Clinical and Experimental Rheumatology, 1992, 10, 161-4.	0.8	14
180	Differential mapping of Fc gamma-binding and monoclonal antibody-reactive epitopes on gE, the Fc gamma-binding glycoprotein of herpes simplex virus type 1. Journal of Immunology, 1992, 149, 2415-27.	0.8	17

#	ARTICLE	IF	CITATIONS
181	Rheumatoid factors react with Fab fragments of monoclonal antibodies to herpes simplex virus types 1 and 2 Fc-binding proteins. <i>Arthritis and Rheumatism</i> , 1991, 34, 846-855.	6.7	11
182	Autoantibodies to the HLA-B27 sequence cross-react with the hypothetical peptide from the arthritis-associated Shigella plasmid. <i>Journal of Clinical Investigation</i> , 1990, 86, 1193-1203.	8.2	36
183	Diagnostic significance of fluorocytometric DNA ploidy analysis of gastric cancer with special reference to its comparison with clinical and histopathological diagnosis. <i>Nihon Geka Hokan Archiv FÅ¼r Japanische Chirurgie</i> , 1990, 59, 116-33.	0.0	0
184	Rheumatoid factors may bear the internal image of the Fc gamma-binding protein of herpes simplex virus type 1. <i>Journal of Immunology</i> , 1990, 144, 4742-8.	0.8	22
185	Cross-reactive epitope with <i>Klebsiella pneumoniae</i> nitrogenase in articular tissue of HLA-B27+ patients with ankylosing spondylitis. <i>Arthritis and Rheumatism</i> , 1989, 32, 437-445.	6.7	66
186	Studies of humoral and cell-mediated immunity to peptides shared by HLA-27.1 and <i>Klebsiella pneumoniae</i> nitrogenase in ankylosing spondylitis. <i>Clinical and Experimental Immunology</i> , 1989, 76, 354-60.	2.6	25
187	Effects of galactose depletion from oligosaccharide chains on immunological activities of human IgG. <i>Journal of Rheumatology</i> , 1989, 16, 285-90.	2.0	141
188	Synovial Expression of HLA-B27 Related Antigens in Ankylosing Spondylitis. <i>Scandinavian Journal of Rheumatology</i> , 1988, 17, 23-25.	1.1	1
189	2-dimensional flow cytometric analysis of peripheral blood T lymphocytes from patients with systemic lupus erythematosus: preferential expression of HLA-DR antigen on the surface of Leu 2a+ cells. <i>Journal of Rheumatology</i> , 1988, 15, 946-51.	2.0	11