
## Sandra Nolte

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/567533/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	Disability-adjusted life years (DALYs) for 291 diseases and injuries in 21 regions, 1990–2010: a systematic analysis for the Global Burden of Disease Study 2010. Lancet, The, 2012, 380, 2197-2223.	13.7	7,061
2	Years lived with disability (YLDs) for 1160 sequelae of 289 diseases and injuries 1990–2010: a systematic analysis for the Global Burden of Disease Study 2010. Lancet, The, 2012, 380, 2163-2196.	13.7	6,376
3	Global, regional, and national age–sex specific all-cause and cause-specific mortality for 240 causes of death, 1990–2013: a systematic analysis for the Global Burden of Disease Study 2013. Lancet, The, 2015, 385, 117-171.	13.7	5,847
4	Global, regional, and national incidence, prevalence, and years lived with disability for 328 diseases and injuries for 195 countries, 1990–2016: a systematic analysis for the Global Burden of Disease Study 2016. Lancet, The, 2017, 390, 1211-1259.	13.7	5,578
5	Global, regional, and national incidence, prevalence, and years lived with disability for 301 acute and chronic diseases and injuries in 188 countries, 1990–2013: a systematic analysis for the Global Burden of Disease Study 2013. Lancet, The, 2015, 386, 743-800.	13.7	4,951
6	Global, Regional, and National Cancer Incidence, Mortality, Years of Life Lost, Years Lived With Disability, and Disability-Adjusted Life-years for 32 Cancer Groups, 1990 to 2015. JAMA Oncology, 2017, 3, 524.	7.1	4,254
7	The global burden of hip and knee osteoarthritis: estimates from the Global Burden of Disease 2010 study. Annals of the Rheumatic Diseases, 2014, 73, 1323-1330.	0.9	2,433
8	The Global Burden of Cancer 2013. JAMA Oncology, 2015, 1, 505.	7.1	2,269
9	Global, regional, and national comparative risk assessment of 79 behavioural, environmental and occupational, and metabolic risks or clusters of risks in 188 countries, 1990–2013: a systematic analysis for the Global Burden of Disease Study 2013. Lancet, The, 2015, 386, 2287-2323.	13.7	2,184
10	Global, regional, and national disability-adjusted life years (DALYs) for 306 diseases and injuries and healthy life expectancy (HALE) for 188 countries, 1990–2013: quantifying the epidemiological transition. Lancet, The, 2015, 386, 2145-2191.	13.7	1,544
11	Global, regional, and national levels and causes of maternal mortality during 1990–2013: a systematic analysis for the Global Burden of Disease Study 2013. Lancet, The, 2014, 384, 980-1004.	13.7	1,230
12	Global, regional, and national incidence and mortality for HIV, tuberculosis, and malaria during 1990–2013: a systematic analysis for the Global Burden of Disease Study 2013. Lancet, The, 2014, 384, 1005-1070.	13.7	786
13	Healthcare Access and Quality Index based on mortality from causes amenable to personal health care in 195 countries and territories, 1990–2015: a novel analysis from the Global Burden of Disease Study 2015. Lancet, The, 2017, 390, 231-266.	13.7	480
14	Global and National Burden of Diseases and Injuries Among Children and Adolescents Between 1990 and 2013. JAMA Pediatrics, 2016, 170, 267.	6.2	479
15	Measuring the health-related Sustainable Development Goals in 188 countries: a baseline analysis from the Global Burden of Disease Study 2015. Lancet, The, 2016, 388, 1813-1850.	13.7	413
16	General population normative data for the EORTC QLQ-C30 health-related quality of life questionnaire based on 15,386 persons across 13 European countries, Canada and the Unites States. European Journal of Cancer, 2019, 107, 153-163.	2.8	259
17	Systematic skin cancer screening in Northern Germany. Journal of the American Academy of Dermatology, 2012, 66, 201-211.	1.2	256
18	Understanding the quality of life (QOL) issues in survivors of cancer: towards the development of an EORTC QOL cancer survivorship questionnaire. Health and Quality of Life Outcomes, 2018, 16, 114.	2.4	133

#	Article	IF	CITATIONS
19	Effects of a Psychological Internet Intervention in the Treatment of Mild to Moderate Depressive Symptoms: Results of the EVIDENT Study, a Randomized Controlled Trial. Psychotherapy and Psychosomatics, 2016, 85, 218-228.	8.8	124
20	The extent and breadth of benefits from participating in chronic disease self-management courses: A national patient-reported outcomes survey. Patient Education and Counseling, 2007, 65, 351-360.	2.2	113
21	A systematic review of outcomes of chronic disease self-management interventions. Quality of Life Research, 2013, 22, 1805-1816.	3.1	107
22	Skin cancer screening participation and impact on melanoma incidence in Germany – an observational study on incidence trends in regions with and without population-based screening. British Journal of Cancer, 2012, 106, 970-974.	6.4	100
23	Validation of patient health questionnaire (PHQ) for major depression in Chinese outpatients with multiple somatic symptoms: A multicenter cross-sectional study. Journal of Affective Disorders, 2015, 174, 636-643.	4.1	78
24	German translation, cultural adaptation, and validation of the Health Literacy Questionnaire (HLQ). PLoS ONE, 2017, 12, e0172340.	2.5	75
25	Standardization of health outcomes assessment for depression and anxiety: recommendations from the ICHOM Depression and Anxiety Working Group. Quality of Life Research, 2017, 26, 3211-3225.	3.1	70
26	Assessment of proximal outcomes of self-management programs: translation and psychometric evaluation of a German version of the Health Education Impact Questionnaire (heiQâ"¢). Quality of Life Research, 2013, 22, 1391-1403.	3.1	67
27	Frequency of Excisions and Yields of Malignant Skin Tumors in a Population-Based Screening Intervention of 360Â288 Whole-Body Examinations. Archives of Dermatology, 2012, 148, 903-10.	1.4	66
28	Tests of measurement invariance failed to support the application of the "then-test― Journal of Clinical Epidemiology, 2009, 62, 1173-1180.	5.0	47
29	Establishing the European Norm for the health-related quality of life domains of the computer-adaptive test EORTC CAT Core. European Journal of Cancer, 2019, 107, 133-141.	2.8	38
30	Implications of response shift for micro-, meso-, and macro-level healthcare decision-making using results of patient-reported outcome measures. Quality of Life Research, 2021, 30, 3343-3357.	3.1	38
31	Using structural equation modeling to detect response shift in performance and health-related quality of life scores of multiple sclerosis patients. Quality of Life Research, 2011, 20, 1527-1540.	3.1	34
32	Updated EORTC QLQ-C30 general population norm data for Germany. European Journal of Cancer, 2020, 137, 161-170.	2.8	32
33	Implementation of the Kids-CAT in clinical settings: a newly developed computer-adaptive test to facilitate the assessment of patient-reported outcomes of children and adolescents in clinical practice in Germany. Quality of Life Research, 2016, 25, 585-594.	3.1	31
34	Defining and Predicting Patterns of Early Response in a Web-Based Intervention for Depression. Journal of Medical Internet Research, 2017, 19, e206.	4.3	31
35	Measurement issues in the evaluation of chronic disease self-management programs. Quality of Life Research, 2013, 22, 1655-1664.	3.1	30
36	Comparison of the Factor Structure of the Patient Health Questionnaire for Somatic Symptoms (PHQ-15) in Germany, the Netherlands, and China. A Transcultural Structural Equation Modeling (SEM) Study. Frontiers in Psychiatry, 2018, 9, 240.	2.6	29

#	Article	IF	CITATIONS
37	Absence of social desirability bias in the evaluation of chronic disease self-management interventions. Health and Quality of Life Outcomes, 2013, 11, 114.	2.4	26
38	A new computerized adaptive test advancing the measurement of health-related quality of life (HRQoL) in children: the Kids-CAT. Quality of Life Research, 2015, 24, 871-884.	3.1	26
39	Interval cancers after skin cancer screening: incidence, tumour characteristics and risk factors for cutaneous melanoma. British Journal of Cancer, 2017, 116, 253-259.	6.4	26
40	The validation of a computer-adaptive test (CAT) for assessing health-related quality of life in children and adolescents in a clinical sample: study design, methods and first results of the Kids-CAT study. Quality of Life Research, 2017, 26, 1105-1117.	3.1	24
41	Psychometric evaluation of the PROMIS® Depression Item Bank: an illustration of classical test theory methods. Journal of Patient-Reported Outcomes, 2019, 3, 46.	1.9	24
42	Factor structure and measurement invariance of the Health Education Impact Questionnaire: Does the subjectivity of the response perspective threaten the contextual validity of inferences?. SAGE Open Medicine, 2015, 3, 205031211558504.	1.8	22
43	Predictors of self-reported health-related quality of life according to the EQ-5D-Y in chronically ill children and adolescents with asthma, diabetes, and juvenile arthritis: longitudinal results. Quality of Life Research, 2018, 27, 879-890.	3.1	22
44	The inclusion of â€~then-test' questions in post-test questionnaires alters post-test responses: a randomized study of bias in health program evaluation. Quality of Life Research, 2012, 21, 487-494.	3.1	21
45	Adaptation and validation of the Japanese version of the Health Education Impact Questionnaire (heiQâ€J) for the evaluation of selfâ€management education interventions. Japan Journal of Nursing Science, 2013, 10, 255-266.	1.3	21
46	Adult Measures of General Health and Healthâ€Related Quality of Life. Arthritis Care and Research, 2020, 72, 522-564.	3.4	21
47	International validation of the EORTC CAT Core: a new adaptive instrument for measuring core quality of life domains in cancer. Quality of Life Research, 2020, 29, 1405-1417.	3.1	21
48	Characteristics of participants in a randomized trial of an Internet intervention for depression (EVIDENT) in comparison to a national sample (DECS1). Internet Interventions, 2017, 9, 46-50.	2.7	20
49	An initial psychometric evaluation of the German PROMIS v1.2 Physical Function item bank in patients with a wide range of health conditions. Clinical Rehabilitation, 2018, 32, 84-93.	2.2	20
50	Normative data for the EORTC QLQ-C30 from the Austrian general population. Health and Quality of Life Outcomes, 2020, 18, 275.	2.4	20
51	Measurement invariance across chronic conditions: a systematic review and an empirical investigation of the Health Education Impact Questionnaire (heiQâ,,¢). Health and Quality of Life Outcomes, 2014, 12, 56.	2.4	19
52	Predictors of health-related quality of life in chronically ill children and adolescents over time. Journal of Psychosomatic Research, 2018, 109, 63-70.	2.6	19
53	The ProCaSP study: quality of life outcomes of prostate cancer patients after radiotherapy or radical prostatectomy in a cohort study. BMC Urology, 2015, 15, 28.	1.4	18
54	Making robust decisions about the impact of health education programs: Psychometric evaluation of the Health Education Impact Questionnaire (heiQ) in diverse patient groups in Norway. Patient Education and Counseling, 2016, 99, 1733-1738.	2.2	18

#	Article	IF	CITATIONS
55	Using Patient Health Questionnaire-9 item parameters of a common metric resulted in similar depression scores compared to independent item response theory model reestimation. Journal of Clinical Epidemiology, 2016, 71, 25-34.	5.0	17
56	The regression discontinuity design showed to be a valid alternative to a randomized controlled trial for estimating treatment effects. Journal of Clinical Epidemiology, 2017, 82, 94-102.	5.0	16
57	Validity arguments for patient-reported outcomes: justifying the intended interpretation and use of data. Journal of Patient-Reported Outcomes, 2021, 5, 64.	1.9	15
58	Languageâ€related differential item functioning between English and German PROMIS Depression items is negligible. International Journal of Methods in Psychiatric Research, 2017, 26, .	2.1	14
59	Is regular use of physiotherapy treatment associated with health locus of control and self-management competency? A study of patients with musculoskeletal disorders undergoing physiotherapy in primary health care. Musculoskeletal Science and Practice, 2018, 36, 43-47.	1.3	14
60	Quality of life in patients with relapsed/refractory multiple myeloma during ixazomib-thalidomide-dexamethasone induction and ixazomib maintenance therapy and comparison to the general population. Leukemia and Lymphoma, 2020, 61, 377-386.	1.3	14
61	The German SCREEN project – design and evaluation of the communication strategy. European Journal of Public Health, 2015, 25, 150-155.	0.3	13
62	Comparison of patient-reported need of psycho-oncologic support and the doctor's perspective: how do they relate to disease severity in melanoma patients?. Psycho-Oncology, 2016, 25, 1271-1277.	2.3	12
63	EORTC QLQ-C30 general population normative data for Italy by sex, age and health condition: an analysis of 1,036 individuals. BMC Public Health, 2022, 22, .	2.9	12
64	Measurement of stable changes of self-management skills after rehabilitation: a latent state–trait analysis of the Health Education Impact Questionnaire (heiQâ"¢). Quality of Life Research, 2014, 23, 2531-2543.	3.1	11
65	Varying the item format improved the range of measurement in patient-reported outcome measures assessing physical function. Arthritis Research and Therapy, 2017, 19, 66.	3.5	11
66	General Spanish population normative data analysis for the EORTC QLQ-C30 by sex, age, and health condition. Health and Quality of Life Outcomes, 2021, 19, 208.	2.4	11
67	On the validity of measuring change over time in routine clinical assessment: a close examination of item-level response shifts in psychosomatic inpatients. Quality of Life Research, 2016, 25, 1339-1347.	3.1	10
68	Different Aspects of Self-Reported Quality of Life in 450 German Melanoma Survivors. Cancers, 2011, 3, 2316-2332.	3.7	9
69	Critical Appraisal of Clinical Practice Guidelines for Adaptation in the Evidence-Based Guideline "Prevention of Skin Cancer― JAMA Dermatology, 2013, 149, 466.	4.1	7
70	A 67-item stress resilience item bank showing high content validity wasÂdeveloped in a psychosomatic sample. Journal of Clinical Epidemiology, 2018, 100, 1-12.	5.0	7
71	Update on the psychometric properties and minimal important difference (MID) thresholds of the FACT-M questionnaire for use in treatment-naÃ <sup>-</sup> ve and previously treated patients with metastatic Merkel cell carcinoma. Health and Quality of Life Outcomes, 2020, 18, 145.	2.4	6
72	Norm-based comparison of the quality-of-life impact of ravulizumab and eculizumab in paroxysmal nocturnal hemoglobinuria. Orphanet Journal of Rare Diseases, 2021, 16, 389.	2.7	6

#	Article	IF	CITATIONS
73	Peripheral injected cholecystokinin-8S modulates the concentration of serotonin in nerve fibers of the rat brainstem. Peptides, 2014, 59, 25-33.	2.4	5
74	Using subdomain-specific item sets affected PROMIS physical function scores differently in cardiology and rheumatology patients. Journal of Clinical Epidemiology, 2020, 127, 151-160.	5.0	5
75	Trajectories of Health-Related Quality of Life and HbA1c Values of Children and Adolescents With Diabetes Mellitus Type 1 Over 6 Months: A Longitudinal Observational Study. Frontiers in Pediatrics, 2019, 7, 566.	1.9	5
76	Do sociodemographic variables moderate effects of an internet intervention for mild to moderate depressive symptoms? An exploratory analysis of a randomised controlled trial (EVIDENT) including 1013 participants. BMJ Open, 2021, 11, e041389.	1.9	5
77	The dopamine antagonist flupentixol does not alter ghrelin-induced food intake in rats. Neuropeptides, 2015, 53, 19-27.	2.2	4
78	Health-related quality of life scores of metastatic pancreatic cancer patients responsive to first line chemotherapy compared to newly derived EORTC QLQ-C30 reference values. BMC Cancer, 2022, 22, .	2.6	4
79	Severity of osteoarthritis in global populations. Osteoarthritis and Cartilage, 2014, 22, S208.	1.3	3
80	Minimal Associations Between Clinical Data and Children's Self-Reported Health-Related Quality of Life in Children With Chronic Conditions—A Cross-Sectional Study. Frontiers in Pediatrics, 2019, 7, 17.	1.9	3
81	Frequently Used Patient-Reported Outcome Measures of General Physical Function Were Highly Correlated With a Multitask Performance Outcome Test Battery. Value in Health, 2022, 25, 1752-1759.	0.3	3
82	Hand-related physical function in rheumatic hand conditions: a protocol for developing a patient-reported outcome measurement instrument: TableÂ1. BMJ Open, 2016, 6, e011174.	1.9	2
83	Course of self-reported symptoms of 342 outpatients receiving medium- versus long-term psychodynamic psychotherapy. BioPsychoSocial Medicine, 2016, 10, 23.	2.1	2
84	Health-related quality of life trajectory of treatment-naive patients with Merkel cell carcinoma receiving avelumab. Future Oncology, 2020, 16, 2089-2099.	2.4	2
85	Investigating the response scale of the EORTC QLQ-C30 in German cancer patients and a population survey. Health and Quality of Life Outcomes, 2021, 19, 235.	2.4	2
86	Reference values for EORTC QLQ-C30, for metastatic pancreatic cancer (mPC): Enhancing the interpretation of HRQoL scores in the POLO trial Journal of Clinical Oncology, 2021, 39, 393-393.	1.6	1
87	Evaluation of an electronic psycho-oncological adaptive screening program (EPAS) with immediate patient feedback: findings from a German cluster intervention study. Journal of Cancer Survivorship, 2021, , 1.	2.9	1
88	Health-related quality of life in patients with metastatic Merkel cell carcinoma receiving second-line or later avelumab treatment: 36-month follow-up data. Annals of Oncology, 2019, 30, v538.	1.2	0
89	1092P Non-progression with avelumab treatment in patients with metastatic Merkel cell carcinoma (mMCC) is associated with a clinically meaningful better health-related quality of life compared with progressive disease. Annals of Oncology, 2020, 31, S740-S741.	1.2	0