

Evan D Paul

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/5751432/publications.pdf>

Version: 2024-02-01

26
papers

2,784
citations

394421

19
h-index

552781

26
g-index

27
all docs

27
docs citations

27
times ranked

3268
citing authors

| # | ARTICLE | IF | CITATIONS |
|----|---|------|-----------|
| 1 | Medial prefrontal cortex determines how stressor controllability affects behavior and dorsal raphe nucleus. <i>Nature Neuroscience</i> , 2005, 8, 365-371. | 14.8 | 823 |
| 2 | MicroRNA 135 Is Essential for Chronic Stress Resiliency, Antidepressant Efficacy, and Intact Serotonergic Activity. <i>Neuron</i> , 2014, 83, 344-360. | 8.1 | 321 |
| 3 | Previous Experience with Behavioral Control over Stress Blocks the Behavioral and Dorsal Raphe Nucleus Activating Effects of Later Uncontrollable Stress: Role of the Ventral Medial Prefrontal Cortex. <i>Journal of Neuroscience</i> , 2006, 26, 13264-13272. | 3.6 | 203 |
| 4 | Behavioral control, the medial prefrontal cortex, and resilience. <i>Dialogues in Clinical Neuroscience</i> , 2006, 8, 397-406. | 3.7 | 182 |
| 5 | 5-Hydroxytryptamine 2C Receptors in the Basolateral Amygdala Are Involved in the Expression of Anxiety After Uncontrollable Traumatic Stress. <i>Biological Psychiatry</i> , 2010, 67, 339-345. | 1.3 | 173 |
| 6 | Activation of the ventral medial prefrontal cortex during an uncontrollable stressor reproduces both the immediate and long-term protective effects of behavioral control. <i>Neuroscience</i> , 2008, 154, 1178-1186. | 2.3 | 134 |
| 7 | Functional topography of serotonergic systems supports the Deakin/Graeff hypothesis of anxiety and affective disorders. <i>Journal of Psychopharmacology</i> , 2013, 27, 1090-1106. | 4.0 | 117 |
| 8 | Microinjection of urocortin 2 into the dorsal raphe nucleus activates serotonergic neurons and increases extracellular serotonin in the basolateral amygdala. <i>Neuroscience</i> , 2004, 129, 509-519. | 2.3 | 115 |
| 9 | The role of prior stressor controllability and the dorsal raphe nucleus in sucrose preference and social exploration. <i>Behavioural Brain Research</i> , 2008, 193, 87-93. | 2.2 | 91 |
| 10 | The Sensory Insular Cortex Mediates the Stress-Buffering Effects of Safety Signals But Not Behavioral Control. <i>Journal of Neuroscience</i> , 2008, 28, 13703-13711. | 3.6 | 86 |
| 11 | Behavioral control over shock blocks behavioral and neurochemical effects of later social defeat. <i>Neuroscience</i> , 2010, 165, 1031-1038. | 2.3 | 80 |
| 12 | Repeated social defeat increases reactive emotional coping behavior and alters functional responses in serotonergic neurons in the rat dorsal raphe nucleus. <i>Physiology and Behavior</i> , 2011, 104, 272-282. | 2.1 | 78 |
| 13 | The Deakin/Graeff hypothesis: Focus on serotonergic inhibition of panic. <i>Neuroscience and Biobehavioral Reviews</i> , 2014, 46, 379-396. | 6.1 | 69 |
| 14 | Bacterial infection early in life protects against stressor-induced depressive-like symptoms in adult rats. <i>Psychoneuroendocrinology</i> , 2008, 33, 261-269. | 2.7 | 64 |
| 15 | MicroRNA-19b Associates with Ago2 in the Amygdala Following Chronic Stress and Regulates the Adrenergic Receptor Beta 1. <i>Journal of Neuroscience</i> , 2014, 34, 15070-15082. | 3.6 | 56 |
| 16 | Chronic Activation of Corticotropin-Releasing Factor Type 2 Receptors Reveals a Key Role for 5-HT1A Receptor Responsiveness in Mediating Behavioral and Serotonergic Responses to Stressful Challenge. <i>Biological Psychiatry</i> , 2012, 72, 437-447. | 1.3 | 33 |
| 17 | A SARS-CoV-2 mutant from B.1.258 lineage with Δ H69/ Δ V70 deletion in the Spike protein circulating in Central Europe in the fall 2020. <i>Virus Genes</i> , 2021, 57, 556-560. | 1.6 | 27 |
| 18 | Increased anxiety in corticotropin-releasing factor type 2 receptor-null mice requires recent acute stress exposure and is associated with dysregulated serotonergic activity in limbic brain areas. <i>Biology of Mood & Anxiety Disorders</i> , 2014, 4, 1. | 4.7 | 26 |

| # | ARTICLE | IF | CITATIONS |
|----|---|-----|-----------|
| 19 | Inescapable but not escapable stress leads to increased struggling behavior and basolateral amygdala c-fos gene expression in response to subsequent novel stress challenge. <i>Neuroscience</i> , 2010, 170, 138-148. | 2.3 | 24 |
| 20 | Surveillance of SARS-CoV-2 lineage B.1.1.7 in Slovakia using a novel, multiplexed RT-qPCR assay. <i>Scientific Reports</i> , 2021, 11, 20494. | 3.3 | 24 |
| 21 | Role of the dorsomedial hypothalamus in glucocorticoid-mediated feedback inhibition of the hypothalamicâ€“pituitaryâ€“adrenal axis. <i>Stress</i> , 2015, 18, 76-87. | 1.8 | 15 |
| 22 | Adenosine-to-Inosine RNA Editing Within Corticolimbic Brain Regions Is Regulated in Response to Chronic Social Defeat Stress in Mice. <i>Frontiers in Psychiatry</i> , 2019, 10, 277. | 2.6 | 15 |
| 23 | Whole-body hyperthermia and a subthreshold dose of citalopram act synergistically to induce antidepressant-like behavioral responses in adolescent rats. <i>Progress in Neuro-Psychopharmacology and Biological Psychiatry</i> , 2017, 79, 162-168. | 4.8 | 8 |
| 24 | Sequential development of several RTâ€“qPCR tests using LNA nucleotides and dual probe technology to differentiate SARSâ€“CoVâ€“2 from influenza A and B. <i>Microbial Biotechnology</i> , 2022, 15, 1995-2021. | 4.2 | 6 |
| 25 | Fibroblast Growth Factor 8 Deficiency Compromises the Functional Response of the Serotonergic System to Stress. <i>PLoS ONE</i> , 2014, 9, e101420. | 2.5 | 4 |
| 26 | Interactions between whole-body heating and citalopram on body temperature, antidepressant-like behaviour, and neurochemistry in adolescent male rats. <i>Behavioural Brain Research</i> , 2019, 359, 428-439. | 2.2 | 3 |