

Volker Hans Haase

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/5770689/publications.pdf>

Version: 2024-02-01

109
papers

17,020
citations

17405

63
h-index

27345

106
g-index

109
all docs

109
docs citations

109
times ranked

18825
citing authors

#	ARTICLE	IF	CITATIONS
1	Disruption of mitochondrial complex III in cap mesenchyme but not in ureteric progenitors results in defective nephrogenesis associated with amino acid deficiency. <i>Kidney International</i> , 2022, , .	2.6	0
2	The ins and outs of ferric citrate. <i>Kidney International</i> , 2022, 101, 668-670.	2.6	1
3	EPO synthesis induced by HIF-1 α inhibition is dependent on myofibroblast transdifferentiation and colocalizes with non-injured nephron segments in murine kidney fibrosis. <i>Acta Physiologica</i> , 2022, 235, e13826.	1.8	18
4	Inhibition of hypoxia-inducible factor-prolyl hydroxylation protects from cyclophosphamide-induced bladder injury and urinary dysfunction. <i>American Journal of Physiology - Renal Physiology</i> , 2022, 323, F81-F91.	1.3	1
5	Stabilization of hypoxia-inducible factor ameliorates glomerular injury sensitization after tubulointerstitial injury. <i>Kidney International</i> , 2021, 99, 620-631.	2.6	13
6	Kidney epithelial targeted mitochondrial transcription factor A deficiency results in progressive mitochondrial depletion associated with severe cystic disease. <i>Kidney International</i> , 2021, 99, 657-670.	2.6	16
7	Inactivation of HIF-1 α prolyl 4-hydroxylases 1, 2 and 3 in NG2-expressing cells induces HIF-1 α -mediated neurovascular expansion independent of erythropoietin. <i>Acta Physiologica</i> , 2021, 231, e13547.	1.8	9
8	Hypoxia-inducible factor-1 α prolyl hydroxylase inhibitors in the treatment of anemia of chronic kidney disease. <i>Kidney International Supplements</i> , 2021, 11, 8-25.	4.6	75
9	Over-Generalizing About GC (Hypoxia): Pitfalls of Limiting Breadth of Experimental Systems and Analyses in Framing Informatics Conclusions. <i>Frontiers in Immunology</i> , 2021, 12, 664249.	2.2	8
10	Pharmacological HIF-1 α inhibition reduces renovascular resistance and increases glomerular filtration by stimulating nitric oxide generation. <i>Acta Physiologica</i> , 2021, 233, e13668.	1.8	14
11	Controversies in optimal anemia management: conclusions from a Kidney Disease: Improving Global Outcomes (KDIGO) Conference. <i>Kidney International</i> , 2021, 99, 1280-1295.	2.6	103
12	Got glycogen? An energy resource in HIF-mediated prevention of ischemic kidney injury. <i>Kidney International</i> , 2020, 97, 645-647.	2.6	2
13	Hypoxia-Inducible Factor Activators in Renal Anemia: Current Clinical Experience. <i>Advances in Chronic Kidney Disease</i> , 2019, 26, 253-266.	0.6	135
14	Hypoxia-inducible factors in CD4 ⁺ T cells promote metabolism, switch cytokine secretion, and T cell help in humoral immunity. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2019, 116, 8975-8984.	3.3	100
15	A Unilateral Facial Rash with Eye Involvement. <i>American Journal of Medicine</i> , 2019, 132, 823-825.	0.6	0
16	ARNT as a Novel Antifibrotic Target in CKD. <i>American Journal of Kidney Diseases</i> , 2019, 73, 281-284.	2.1	4
17	Effects of vadadustat on hemoglobin concentrations in patients receiving hemodialysis previously treated with erythropoiesis-stimulating agents. <i>Nephrology Dialysis Transplantation</i> , 2019, 34, 90-99.	0.4	62
18	Therapeutic targeting of the HIF oxygen-sensing pathway: Lessons learned from clinical studies. <i>Experimental Cell Research</i> , 2017, 356, 160-165.	1.2	44

#	ARTICLE	IF	CITATIONS
19	Inflamed fat and mitochondrial dysfunction in end-stage renal disease links to hypoxia—could curcumin be of benefit?. <i>Nephrology Dialysis Transplantation</i> , 2017, 32, 909-912.	0.4	7
20	HIF—prolyl hydroxylases as therapeutic targets in erythropoiesis and iron metabolism. <i>Hemodialysis International</i> , 2017, 21, S110-S124.	0.4	120
21	Hypoxia-inducible factor prolyl-4-hydroxylation in FOXD1 lineage cells is essential for normal kidney development. <i>Kidney International</i> , 2017, 92, 1370-1383.	2.6	22
22	Oxygen sensors as therapeutic targets in kidney disease. <i>Nephrologie Et Therapeutique</i> , 2017, 13, S29-S34.	0.2	15
23	Prolyl-4-hydroxylase 2 and 3 coregulate murine erythropoietin in brain pericytes. <i>Blood</i> , 2016, 128, 2550-2560.	0.6	32
24	Germinal centre hypoxia and regulation of antibody qualities by a hypoxia response system. <i>Nature</i> , 2016, 537, 234-238.	13.7	215
25	Vadadustat, a novel oral HIF stabilizer, provides effective anemia treatment in nondialysis-dependent chronic kidney disease. <i>Kidney International</i> , 2016, 90, 1115-1122.	2.6	187
26	The Endothelial Prolyl-4-Hydroxylase Domain 2/Hypoxia-Inducible Factor 2 Axis Regulates Pulmonary Artery Pressure in Mice. <i>Molecular and Cellular Biology</i> , 2016, 36, 1584-1594.	1.1	110
27	Endothelial HIF signaling regulates pulmonary fibrosis-associated pulmonary hypertension. <i>American Journal of Physiology - Lung Cellular and Molecular Physiology</i> , 2016, 310, L249-L262.	1.3	65
28	Renal epithelium regulates erythropoiesis via HIF-dependent suppression of erythropoietin. <i>Journal of Clinical Investigation</i> , 2016, 126, 1425-1437.	3.9	47
29	Distinct subpopulations of FOXD1 stroma-derived cells regulate renal erythropoietin. <i>Journal of Clinical Investigation</i> , 2016, 126, 1926-1938.	3.9	91
30	Anaemia in kidney disease: harnessing hypoxia responses for therapy. <i>Nature Reviews Nephrology</i> , 2015, 11, 394-410.	4.1	235
31	Molecular mechanisms of ischemic preconditioning in the kidney. <i>American Journal of Physiology - Renal Physiology</i> , 2015, 309, F821-F834.	1.3	67
32	A Breath of Fresh Air for Diabetic Nephropathy. <i>Journal of the American Society of Nephrology: JASN</i> , 2015, 26, 239-241.	3.0	12
33	Inflammation and hypoxia in the kidney: friends or foes?. <i>Kidney International</i> , 2015, 88, 213-215.	2.6	16
34	Muc1 is protective during kidney ischemia-reperfusion injury. <i>American Journal of Physiology - Renal Physiology</i> , 2015, 308, F1452-F1462.	1.3	35
35	FO015AKB-6548, A NOVEL HYPOXIA-INDUCIBLE FACTOR PROLYL-HYDROXYLASE INHIBITOR (HIF-PHI) FOR THE TREATMENT OF ANEMIA IN PATIENTS WITH CHRONIC KIDNEY DISEASE NOT ON DIALYSIS (ND-CKD). <i>Nephrology Dialysis Transplantation</i> , 2015, 30, iii8-iii8.	0.4	4
36	Endothelial HIF-2 mediates protection and recovery from ischemic kidney injury. <i>Journal of Clinical Investigation</i> , 2014, 124, 2396-2409.	3.9	150

#	ARTICLE	IF	CITATIONS
37	CD73-Dependent Generation of Adenosine and Endothelial Adora2b Signaling Attenuate Diabetic Nephropathy. <i>Journal of the American Society of Nephrology: JASN</i> , 2014, 25, 547-563.	3.0	40
38	Regulation of erythropoiesis by hypoxia-inducible factors. <i>Blood Reviews</i> , 2013, 27, 41-53.	2.8	522
39	Mechanisms of Hypoxia Responses in Renal Tissue. <i>Journal of the American Society of Nephrology: JASN</i> , 2013, 24, 537-541.	3.0	121
40	Activation of Hypoxia-Inducible Factor-1 in Adipocytes Results in Pathological Cardiac Hypertrophy. <i>Journal of the American Heart Association</i> , 2013, 2, e000548.	1.6	34
41	Preischemic targeting of HIF prolyl hydroxylation inhibits fibrosis associated with acute kidney injury. <i>American Journal of Physiology - Renal Physiology</i> , 2012, 302, F1172-F1179.	1.3	104
42	Proximal tubule sphingosine kinase-1 has a critical role in A1 adenosine receptor-mediated renal protection from ischemia. <i>Kidney International</i> , 2012, 82, 878-891.	2.6	36
43	Myeloid Cell-Derived Hypoxia-Inducible Factor Attenuates Inflammation in Unilateral Ureteral Obstruction-Induced Kidney Injury. <i>Journal of Immunology</i> , 2012, 188, 5106-5115.	0.4	86
44	Hypoxia-inducible factor regulates hepcidin via erythropoietin-induced erythropoiesis. <i>Journal of Clinical Investigation</i> , 2012, 122, 4635-4644.	3.9	263
45	Equilibrative nucleoside transporter 1 (ENT1) regulates postischemic blood flow during acute kidney injury in mice. <i>Journal of Clinical Investigation</i> , 2012, 122, 693-710.	3.9	99
46	Renal cancer: Oxygen meets metabolism. <i>Experimental Cell Research</i> , 2012, 318, 1057-1067.	1.2	28
47	Hypoxia-inducible factor signaling in the development of kidney fibrosis. <i>Fibrogenesis and Tissue Repair</i> , 2012, 5, S16.	3.4	54
48	AT 1A Angiotensin Receptors in the Renal Proximal Tubule Regulate Blood Pressure. <i>Cell Metabolism</i> , 2011, 13, 469-475.	7.2	220
49	Angiotensin II: breathtaking in the renal medulla. <i>Kidney International</i> , 2011, 79, 269-271.	2.6	3
50	Oxygen-Dependent Regulation of Erythropoiesis. , 2011, , 437-463.		0
51	Hepatic HIF-2 regulates erythropoietic responses to hypoxia in renal anemia. <i>Blood</i> , 2010, 116, 3039-3048.	0.6	264
52	Astrocyte hypoxic response is essential for pathological but not developmental angiogenesis of the retina. <i>Glia</i> , 2010, 58, 1177-1185.	2.5	142
53	Renal Oxygenation Suppresses VHL Loss-Induced Senescence That Is Caused by Increased Sensitivity to Oxidative Stress. <i>Molecular and Cellular Biology</i> , 2010, 30, 4595-4603.	1.1	38
54	Activation of Sphingosine-1-Phosphate 1 Receptor in the Proximal Tubule Protects Against Ischemia-Reperfusion Injury. <i>Journal of the American Society of Nephrology: JASN</i> , 2010, 21, 955-965.	3.0	109

#	ARTICLE	IF	CITATIONS
55	Sirtuins and Their Relevance to the Kidney. <i>Journal of the American Society of Nephrology: JASN</i> , 2010, 21, 1620-1627.	3.0	103
56	Targeted Deletion of Dicer from Proximal Tubules Protects against Renal Ischemia-Reperfusion Injury. <i>Journal of the American Society of Nephrology: JASN</i> , 2010, 21, 756-761.	3.0	207
57	Hypoxia activates the cyclooxygenase-2/prostaglandin E synthase axis. <i>Carcinogenesis</i> , 2010, 31, 427-434.	1.3	104
58	HO-1 in Control of a Self-Eating Kidney. <i>Journal of the American Society of Nephrology: JASN</i> , 2010, 21, 1600-1602.	3.0	2
59	The sweet side of HIF. <i>Kidney International</i> , 2010, 78, 10-13.	2.6	18
60	VHL Deletion Impairs Mammary Alveologenesis but Is Not Sufficient for Mammary Tumorigenesis. <i>American Journal of Pathology</i> , 2010, 176, 2269-2282.	1.9	12
61	Hypoxic regulation of erythropoiesis and iron metabolism. <i>American Journal of Physiology - Renal Physiology</i> , 2010, 299, F1-F13.	1.3	266
62	Epithelial Notch signaling regulates interstitial fibrosis development in the kidneys of mice and humans. <i>Journal of Clinical Investigation</i> , 2010, 120, 4040-4054.	3.9	306
63	Hypoxia-Inducible Factor 2 Regulates Hepatic Lipid Metabolism. <i>Molecular and Cellular Biology</i> , 2009, 29, 4527-4538.	1.1	283
64	Pathophysiological Consequences of HIF Activation. <i>Annals of the New York Academy of Sciences</i> , 2009, 1177, 57-65.	1.8	68
65	Oxygen regulates epithelial-to-mesenchymal transition: insights into molecular mechanisms and relevance to disease. <i>Kidney International</i> , 2009, 76, 492-499.	2.6	91
66	The VHL Tumor Suppressor: Master Regulator of HIF. <i>Current Pharmaceutical Design</i> , 2009, 15, 3895-3903.	0.9	125
67	The glial cell response is an essential component of hypoxia-induced erythropoiesis in mice. <i>Journal of Clinical Investigation</i> , 2009, 119, 3373-83.	3.9	82
68	Hypoxia-inducible factor-2 regulates vascular tumorigenesis in mice. <i>Oncogene</i> , 2008, 27, 5354-5358.	2.6	136
69	The VHL tumor suppressor and HIF: insights from genetic studies in mice. <i>Cell Death and Differentiation</i> , 2008, 15, 650-659.	5.0	125
70	Hypoxia-Inducible Factor Augments Experimental Colitis Through an MIF-Dependent Inflammatory Signaling Cascade. <i>Gastroenterology</i> , 2008, 134, 2036-2048.e3.	0.6	146
71	Epidermal Sensing of Oxygen Is Essential for Systemic Hypoxic Response. <i>Cell</i> , 2008, 133, 223-234.	13.5	160
72	Low oxygen stimulates the immune system. <i>Kidney International</i> , 2008, 73, 797-799.	2.6	9

#	ARTICLE	IF	CITATIONS
73	Hypoxia-inducible factor signaling in the development of tissue fibrosis. <i>Cell Cycle</i> , 2008, 7, 1128-1132.	1.3	174
74	Hemoglobin in the Kidney: Breaking with Traditional Dogma. <i>Journal of the American Society of Nephrology: JASN</i> , 2008, 19, 1440-1441.	3.0	7
75	Stable expression of HIF-1 α in tubular epithelial cells promotes interstitial fibrosis. <i>American Journal of Physiology - Renal Physiology</i> , 2008, 295, F1023-F1029.	1.3	234
76	Primary Coenzyme Q Deficiency in <i>Pdss2</i> Mutant Mice Causes Isolated Renal Disease. <i>PLoS Genetics</i> , 2008, 4, e1000061.	1.5	109
77	Hypoxia and podocyte-specific <i>Vhlh</i> deletion confer risk of glomerular disease. <i>American Journal of Physiology - Renal Physiology</i> , 2007, 293, F1397-F1407.	1.3	54
78	Hypoxia-inducible factor-2 (HIF-2) regulates hepatic erythropoietin in vivo. <i>Journal of Clinical Investigation</i> , 2007, 117, 1068-1077.	3.9	496
79	The hypoxia-inducible factor α pathway couples angiogenesis to osteogenesis during skeletal development. <i>Journal of Clinical Investigation</i> , 2007, 117, 1616-1626.	3.9	616
80	Regulation of iron homeostasis by the hypoxia-inducible transcription factors (HIFs). <i>Journal of Clinical Investigation</i> , 2007, 117, 1926-1932.	3.9	538
81	Hypoxia promotes fibrogenesis in vivo via HIF-1 stimulation of epithelial-to-mesenchymal transition. <i>Journal of Clinical Investigation</i> , 2007, 117, 3810-20.	3.9	778
82	Loss of vascular endothelial growth factor expression reduces vascularization, but not growth, of tumors lacking the Von Hippel-Lindau tumor suppressor gene. <i>Oncogene</i> , 2007, 26, 4531-4540.	2.6	10
83	Suppression of Fas-FasL coexpression by erythropoietin mediates erythroblast expansion during the erythropoietic stress response in vivo. <i>Blood</i> , 2006, 108, 123-133.	0.6	192
84	Loss of the tumor suppressor <i>Vhlh</i> leads to upregulation of <i>Cxcr4</i> and rapidly progressive glomerulonephritis in mice. <i>Nature Medicine</i> , 2006, 12, 1081-1087.	15.2	191
85	The VHL/HIF oxygen-sensing pathway and its relevance to kidney disease. <i>Kidney International</i> , 2006, 69, 1302-1307.	2.6	133
86	Hypoxia-inducible factors in the kidney. <i>American Journal of Physiology - Renal Physiology</i> , 2006, 291, F271-F281.	1.3	284
87	Cytoprotective Effects of Hypoxia against Cisplatin-Induced Tubular Cell Apoptosis: Involvement of Mitochondrial Inhibition and p53 Suppression. <i>Journal of the American Society of Nephrology: JASN</i> , 2006, 17, 1875-1885.	3.0	63
88	Renal Cyst Development in Mice with Conditional Inactivation of the von Hippel-Lindau Tumor Suppressor. <i>Cancer Research</i> , 2006, 66, 2576-2583.	0.4	322
89	pVHL Function Is Essential for Endothelial Extracellular Matrix Deposition. <i>Molecular and Cellular Biology</i> , 2006, 26, 2519-2530.	1.1	81
90	Protection of HIF-1-deficient primary renal tubular epithelial cells from hypoxia-induced cell death is glucose dependent. <i>American Journal of Physiology - Renal Physiology</i> , 2005, 289, F1217-F1226.	1.3	33

#	ARTICLE	IF	CITATIONS
91	Hypoxia inducible factor 1 \hat{A} regulates T cell receptor signal transduction. Proceedings of the National Academy of Sciences of the United States of America, 2005, 102, 17071-17076.	3.3	109
92	Inactivation of the Arylhydrocarbon Receptor Nuclear Translocator (Arnt) Suppresses von Hippel-Lindau Disease-Associated Vascular Tumors in Mice. Molecular and Cellular Biology, 2005, 25, 3163-3172.	1.1	132
93	Decreased Growth of Vhl \hat{A} Fibrosarcomas Is Associated with Elevated Levels of Cyclin Kinase Inhibitors p21 and p27. Molecular and Cellular Biology, 2005, 25, 4565-4578.	1.1	71
94	Inflammatory Hypoxia: Role of Hypoxia-Inducible Factor. Cell Cycle, 2005, 4, 255-257.	1.3	137
95	The VHL tumor suppressor in development and disease: Functional studies in mice by conditional gene targeting. Seminars in Cell and Developmental Biology, 2005, 16, 564-574.	2.3	66
96	Deletion of Vhlh in chondrocytes reduces cell proliferation and increases matrix deposition during growth plate development. Development (Cambridge), 2004, 131, 2497-2508.	1.2	119
97	Vhlh Gene Deletion Induces Hif-1-Mediated Cell Death in Thymocytes. Molecular and Cellular Biology, 2004, 24, 9038-9047.	1.1	100
98	Hypoxic induction of Ctgfis directly mediated by Hif-1. American Journal of Physiology - Renal Physiology, 2004, 287, F1223-F1232.	1.3	262
99	Epithelial hypoxia-inducible factor-1 is protective in murine experimental colitis. Journal of Clinical Investigation, 2004, 114, 1098-1106.	3.9	484
100	Epithelial hypoxia-inducible factor-1 is protective in murine experimental colitis. Journal of Clinical Investigation, 2004, 114, 1098-1106.	3.9	358
101	DNA oligonucleotide microarray technology identifies fisp-12 among other potential fibrogenic genes following murine unilateral ureteral obstruction (UUO): Modulation during epithelial-mesenchymal transition. Kidney International, 2003, 64, 2079-2091.	2.6	44
102	HIF-1 \hat{A} Is Essential for Myeloid Cell-Mediated Inflammation. Cell, 2003, 112, 645-657.	13.5	1,862
103	Hypoxia-Induced Gene Expression Occurs Solely through the Action of Hypoxia-Inducible Factor 1 \hat{A} (HIF-1 \hat{A}): Role of Cytoplasmic Trapping of HIF-2 \hat{A} . Molecular and Cellular Biology, 2003, 23, 4959-4971.	1.1	164
104	Ineffective erythropoiesis in Stat5a \hat{A} /5b \hat{A} mice due to decreased survival of early erythroblasts. Blood, 2001, 98, 3261-3273.	0.6	625
105	Vascular tumors in livers with targeted inactivation of the von Hippel-Lindau tumor suppressor. Proceedings of the National Academy of Sciences of the United States of America, 2001, 98, 1583-1588.	3.3	357
106	A Lymphocyte-specific Ltk Tyrosine Kinase Isoform Is Retained in the Endoplasmic Reticulum in Association with Calnexin. Journal of Biological Chemistry, 1997, 272, 1297-1301.	1.6	7
107	The murine NF2 homologue encodes a highly conserved merlin protein with alternative forms. Human Molecular Genetics, 1994, 3, 407-411.	1.4	48
108	A novel moesin-, ezrin-, radixin-like gene is a candidate for the neurofibromatosis 2 tumor suppressor. Cell, 1993, 72, 791-800.	13.5	1,286

#	ARTICLE	IF	CITATIONS
109	Complete Human NF1 cDNA Sequence: Two Alternatively Spliced mRNAs and Absence of Expression in a Neuroblastoma Line. DNA and Cell Biology, 1992, 11, 727-734.	0.9	33