
Leonardo Giovannoni

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/5773151/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	Intralesional administration of L19-IL2/L19-TNF in stage III or stage IVM1a melanoma patients: results of a phase II study. Cancer Immunology, Immunotherapy, 2015, 64, 999-1009.	4.2	138
2	Targeting Interleukin-2 to the Bone Marrow Stroma for Therapy of Acute Myeloid Leukemia Relapsing after Allogeneic Hematopoietic Stem Cell Transplantation. Cancer Immunology Research, 2015, 3, 547-556.	3.4	39
3	The tumor-targeting immunocytokine F16-IL2 in combination with doxorubicin: dose escalation in patients with advanced solid tumors and expansion into patients with metastatic breast cancer. Cell Adhesion and Migration, 2015, 9, 14-21.	2.7	51
4	Potentiating the activity of rituximab against mantle cell lymphoma in mice by targeting interleukin-2 to the neovasculature. Leukemia Research, 2015, 39, 739-748.	0.8	14
5	Armed antibodies for cancer treatment: a promising tool in a changing era. Cancer Immunology, Immunotherapy, 2015, 64, 113-121.	4.2	28
6	Intralesional Treatment of Stage III Metastatic Melanoma Patients with L19–IL2 Results in Sustained Clinical and Systemic Immunologic Responses. Cancer Immunology Research, 2014, 2, 668-678.	3.4	81
7	Radioimmunotherapy with Tenarad, a 1311-labelled antibody fragment targeting the extra-domain A1 of tenascin-C, in patients with refractory Hodgkin's lymphoma. European Journal of Nuclear Medicine and Molecular Imaging, 2014, 41, 867-877.	6.4	40
8	Abundant in vitro expression of the oncofetal ED-B-containing fibronectin translates into selective pharmacodelivery of 131I-L19SIP in a prostate cancer patient. Journal of Cancer Research and Clinical Oncology, 2014, 140, 35-43.	2.5	24
9	Isolated limb perfusion with the tumorâ€ŧargeting human monoclonal antibody–cytokine fusion protein L19â€TNF plus melphalan and mild hyperthermia in patients with locally advanced extremity melanoma. Journal of Surgical Oncology, 2013, 107, 173-179.	1.7	72
10	Phase I/II study of the tumour-targeting human monoclonal antibody–cytokine fusion protein L19-TNF in patients with advanced solid tumours. Journal of Cancer Research and Clinical Oncology, 2013, 139, 447-455.	2.5	92
11	Antibody-Based Delivery of Interleukin-2 to Neovasculature Has Potent Activity Against Acute Myeloid Leukemia. Science Translational Medicine, 2013, 5, 201ra118.	12.4	97
12	Phase 0 Microdosing PET Study Using the Human Mini Antibody F16SIP in Head and Neck Cancer Patients. Journal of Nuclear Medicine, 2013, 54, 397-401.	5.0	47
13	Radretumab Radioimmunotherapy in Patients with Brain Metastasis: A 124I-L19SIP Dosimetric PET Study. Cancer Immunology Research, 2013, 1, 134-143.	3.4	63
14	The antibody-based targeted delivery of TNF in combination with doxorubicin eradicates sarcomas in mice and confers protective immunity. British Journal of Cancer, 2013, 109, 1206-1213.	6.4	78
15	Radioimmunotherapy with Radretumab in Patients with Relapsed Hematologic Malignancies. Journal of Nuclear Medicine, 2012, 53, 922-927.	5.0	65
16	A Dose-Escalation and Signal-Generating Study of the Immunocytokine L19-IL2 in Combination with Dacarbazine for the Therapy of Patients with Metastatic Melanoma. Clinical Cancer Research, 2011, 17, 7732-7742.	7.0	134
17	Expression, engineering and characterization of the tumor-targeting heterodimeric immunocytokine F8-IL12. Protein Engineering, Design and Selection, 2010, 23, 653-661.	2.1	33
18	The tumour-targeting human L19-IL2 immunocytokine: Preclinical safety studies, phase I clinical trial in patients with solid tumours and expansion into patients with advanced renal cell carcinoma. European Journal of Cancer, 2010, 46, 2926-2935.	2.8	149

#	Article	IF	CITATIONS
19	Expression of the oncofetal ED-B–containing fibronectin isoform in hematologic tumors enables ED-B–targeted 131I-L19SIP radioimmunotherapy in Hodgkin lymphoma patients. Blood, 2009, 113, 2265-2274.	1.4	153
20	A general method for the selection of high-level scFv and IgG antibody expression by stably transfected mammalian cells. Protein Engineering, Design and Selection, 2008, 22, 169-174.	2.1	24
21	Antibody-Mediated Delivery of Interleukin-2 to the Stroma of Breast Cancer Strongly Enhances the Potency of Chemotherapy. Clinical Cancer Research, 2008, 14, 6515-6524.	7.0	102
22	Immunoscintigraphic detection of the ED-B domain of fibronectin, a marker of angiogenesis, in patients with cancer. Clinical Cancer Research, 2003, 9, 571-9.	7.0	229
23	HISTIDYL TAGS AND STRUCTURAL STABILIZATION OF LINEAR PEPTIDES. Spectroscopy Letters, 2002, 35, 111-118.	1.0	1
24	Metal ion complexation and folding of linear peptides. Biophysical Chemistry, 2002, 97, 79-86.	2.8	6
25	NMR studies on Ni(II) induced cyclization of a histidine-tagged peptide. Journal of Peptide Science, 2002, 8, 634-641.	1.4	2
26	Recombinant antibodies for the selective targeting of tumor neovasculature. Current Opinion in Drug Discovery & Development, 2002, 5, 204-13.	1.9	1
27	A strategy for the isolation of catalytic activities from repertoires of enzymes displayed on phage 1 1Edited by J. Karn. Journal of Molecular Biology, 1999, 286, 617-633.	4.2	107
28	Increased binding affinity and valence of recombinant antibody fragments lead to improved targeting of tumoral angiogenesis. Cancer Research, 1999, 59, 347-52.	0.9	154