

Arjen ten Have

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/579350/publications.pdf>

Version: 2024-02-01

36
papers

3,872
citations

318942

23
h-index

445137

33
g-index

41
all docs

41
docs citations

41
times ranked

4384
citing authors

#	ARTICLE	IF	CITATIONS
1	Functional Classification and Characterization of the Fungal Glycoside Hydrolase 28 Protein Family. <i>Journal of Fungi</i> (Basel, Switzerland), 2022, 8, 217.	1.5	6
2	Structure-function analysis of Sedolisins: evolution of tripeptidyl peptidase and endopeptidase subfamilies in fungi. <i>BMC Bioinformatics</i> , 2018, 19, 464.	1.2	2
3	Molecular dynamics and structure function analysis show that substrate binding and specificity are major forces in the functional diversification of Equisins. <i>BMC Bioinformatics</i> , 2018, 19, 338.	1.2	3
4	HMMER Cut-off Threshold Tool (HMMERCTTER): Supervised classification of superfamily protein sequences with a reliable cut-off threshold. <i>PLoS ONE</i> , 2018, 13, e0193757.	1.1	16
5	Chlorogenic acid, anthocyanin and flavan-3-ol biosynthesis in flesh and skin of Andean potato tubers (<i>Solanum tuberosum</i> subsp. <i>andigena</i>). <i>Food Chemistry</i> , 2017, 229, 837-846.	4.2	57
6	Computational Functional Analysis of Lipid Metabolic Enzymes. <i>Methods in Molecular Biology</i> , 2017, 1609, 195-216.	0.4	2
7	Evolutionary and Functional Relationships in the Truncated Hemoglobin Family. <i>PLoS Computational Biology</i> , 2016, 12, e1004701.	1.5	36
8	The diversity of algal phospholipase D homologs revealed by biocomputational analysis. <i>Journal of Phycology</i> , 2015, 51, 943-962.	1.0	13
9	Chlorogenic Acid Biosynthesis Appears Linked with Suberin Production in Potato Tuber (<i>Solanum</i>) Tj ETQq1 1 0.784314 rgBT /Ove 2.4 49	2.4	49
10	Phospholipase D \hat{r} knock-out mutants are tolerant to severe drought stress. <i>Plant Signaling and Behavior</i> , 2015, 10, e1089371.	1.2	28
11	Extensive Expansion of A1 Family Aspartic Proteinases in Fungi Revealed by Evolutionary Analyses of 107 Complete Eukaryotic Proteomes. <i>Genome Biology and Evolution</i> , 2014, 6, 1480-1494.	1.1	17
12	The tomato phosphatidylinositol-phospholipase C2 (SIPLC2) is required for defense gene induction by the fungal elicitor xylanase. <i>Journal of Plant Physiology</i> , 2014, 171, 959-965.	1.6	26
13	Evolution and functional diversification of the small heat shock protein/ β -crystallin family in higher plants. <i>Planta</i> , 2012, 235, 1299-1313.	1.6	77
14	Genomic Analysis of the Necrotrophic Fungal Pathogens <i>Sclerotinia sclerotiorum</i> and <i>Botrytis cinerea</i> . <i>PLoS Genetics</i> , 2011, 7, e1002230.	1.5	902
15	Phosphatidic acid production in chitosan-elicited tomato cells, via both phospholipase D and phospholipase C/diacylglycerol kinase, requires nitric oxide. <i>Journal of Plant Physiology</i> , 2011, 168, 534-539.	1.6	86
16	The aspartic proteinase family of three <i>Phytophthora</i> species. <i>BMC Genomics</i> , 2011, 12, 254.	1.2	19
17	Identification of tomato phosphatidylinositol-specific phospholipase-C (PI-PLC) family members and the role of PLC4 and PLC6 in HR and disease resistance. <i>Plant Journal</i> , 2010, 62, 224-239.	2.8	127
18	Nitric Oxide and Phosphatidic Acid Signaling in Plants. <i>Plant Cell Monographs</i> , 2010, , 223-242.	0.4	9

#	ARTICLE	IF	CITATIONS
19	The Botrytis cinerea aspartic proteinase family. Fungal Genetics and Biology, 2010, 47, 53-65.	0.9	101
20	Quantitative resistance to Botrytis cinerea from Solanum neorickii. Euphytica, 2008, 159, 83-92.	0.6	27
21	Nitric Oxide Is Critical for Inducing Phosphatidic Acid Accumulation in Xylanase-elicited Tomato Cells. Journal of Biological Chemistry, 2007, 282, 21160-21168.	1.6	118
22	Partial stem and leaf resistance against the fungal pathogen Botrytis cinerea in wild relatives of tomato. European Journal of Plant Pathology, 2007, 117, 153-166.	0.8	32
23	Three QTLs for Botrytis cinerea resistance in tomato. Theoretical and Applied Genetics, 2007, 114, 585-593.	1.8	50
24	Nitric Oxide Functions as Intermediate in Auxin, Abscisic Acid, and Lipid Signaling Pathways. Plant Cell Monographs, 2006, , 113-130.	0.4	11
25	An aspartic proteinase gene family in the filamentous fungus Botrytis cinerea contains members with novel features. Microbiology (United Kingdom), 2004, 150, 2475-2489.	0.7	72
26	A tomato metacaspase gene is upregulated during programmed cell death in Botrytis cinerea-infected leaves. Planta, 2003, 217, 517-522.	1.6	125
27	The Role of Ethylene and Wound Signaling in Resistance of Tomato to Botrytis cinerea. Plant Physiology, 2002, 129, 1341-1351.	2.3	301
28	The Contribution of Cell Wall Degrading Enzymes to Pathogenesis of Fungal Plant Pathogens. , 2002, , 341-358.		68
29	Botrytis cinerea Endopolygalacturonase Genes Are Differentially Expressed in Various Plant Tissues. Fungal Genetics and Biology, 2001, 33, 97-105.	0.9	129
30	Regulation of endopolygalacturonase gene expression in Botrytis cinerea by galacturonic acid, ambient pH and carbon catabolite repression. Current Genetics, 2000, 37, 152-157.	0.8	131
31	Transgenic Expression of Pear PCIP in Tomato Limits Fungal Colonization. Molecular Plant-Microbe Interactions, 2000, 13, 942-950.	1.4	228
32	Infection Strategies of Botrytis cinerea and Related Necrotrophic Pathogens. , 2000, , 33-64.		115
33	Fungal and plant gene expression during synchronized infection of tomato leaves by Botrytis cinerea. European Journal of Plant Pathology, 1998, 104, 207-220.	0.8	170
34	The Endopolygalacturonase Gene Bcpg1 Is Required for Full Virulence of Botrytis cinerea. Molecular Plant-Microbe Interactions, 1998, 11, 1009-1016.	1.4	513
35	Ethylene biosynthetic genes are differentially expressed during carnation (Dianthus caryophyllus L.) flower senescence. , 1997, 34, 89-97.		144
36	Molecular cloning of two different ACC synthase PCR fragments in carnation flowers and organ-specific expression of the corresponding genes. Plant Molecular Biology, 1994, 26, 453-458.	2.0	57