

Keijo Viiri

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/5825861/publications.pdf>

Version: 2024-02-01

29
papers

2,031
citations

516710

16
h-index

454955

30
g-index

36
all docs

36
docs citations

36
times ranked

4366
citing authors

#	ARTICLE	IF	CITATIONS
1	Glacial history and colonization of Europe by the blue tit <i>Parus caeruleus</i> . <i>Journal of Avian Biology</i> , 2004, 35, 352-359.	1.2	927
2	Short RNAs Are Transcribed from Repressed Polycomb Target Genes and Interact with Polycomb Repressive Complex-2. <i>Molecular Cell</i> , 2010, 38, 675-688.	9.7	338
3	The interaction of PRC2 with RNA or chromatin is mutually antagonistic. <i>Genome Research</i> , 2016, 26, 896-907.	5.5	191
4	DNA-Binding and -Bending Activities of SAP30L and SAP30 Are Mediated by a Zinc-Dependent Module and Monophosphoinositides. <i>Molecular and Cellular Biology</i> , 2009, 29, 342-356.	2.3	53
5	Safety and efficacy of AMG 714 in patients with type 2 refractory coeliac disease: a phase 2a, randomised, double-blind, placebo-controlled, parallel-group study. <i>The Lancet Gastroenterology and Hepatology</i> , 2019, 4, 960-970.	8.1	52
6	The Dnmt3L ADD Domain Controls Cytosine Methylation Establishment during Spermatogenesis. <i>Cell Reports</i> , 2015, 10, 944-956.	6.4	39
7	Extracellular transglutaminase 2 has a role in cell adhesion, whereas intracellular transglutaminase 2 is involved in regulation of endothelial cell proliferation and apoptosis. <i>Cell Proliferation</i> , 2011, 44, 49-58.	5.3	36
8	Myosin IXB gene region and gluten intolerance: linkage to coeliac disease and a putative dermatitis herpetiformis association. <i>Journal of Medical Genetics</i> , 2007, 45, 222-227.	3.2	35
9	SAP30L interacts with members of the Sin3A corepressor complex and targets Sin3A to the nucleolus. <i>Nucleic Acids Research</i> , 2006, 34, 3288-3298.	14.5	34
10	Genome-Wide Transcriptomic Analysis of Intestinal Mucosa in Celiac Disease Patients on a Gluten-Free Diet and Postgluten Challenge. <i>Cellular and Molecular Gastroenterology and Hepatology</i> , 2021, 11, 13-32.	4.5	33
11	Genome-wide repression of eRNA and target gene loci by the ETV6-RUNX1 fusion in acute leukemia. <i>Genome Research</i> , 2016, 26, 1468-1477.	5.5	31
12	Polycomb Repressive Complex 2 Enacts Wnt Signaling in Intestinal Homeostasis and Contributes to the Instigation of Stemness in Diseases Entailing Epithelial Hyperplasia or Neoplasia. <i>Stem Cells</i> , 2017, 35, 445-457.	3.2	30
13	Histological, immunohistochemical and mRNA gene expression responses in coeliac disease patients challenged with gluten using PAXgene fixed paraffin-embedded duodenal biopsies. <i>BMC Gastroenterology</i> , 2019, 19, 189.	2.0	27
14	Dermatitis Herpetiformis Refractory to Gluten-free Dietary Treatment. <i>Acta Dermato-Venereologica</i> , 2016, 96, 82-86.	1.3	24
15	Modulation of Wnt/BMP pathways during corneal differentiation of hPSC maintains ABCG2-positive LSC population that demonstrates increased regenerative potential. <i>Stem Cell Research and Therapy</i> , 2019, 10, 236.	5.5	21
16	Phosphoinositides as Regulators of Protein-Chromatin Interactions. <i>Science Signaling</i> , 2012, 5, pe19.	3.6	20
17	Extensive reprogramming of the nascent transcriptome during iPSC to hepatocyte differentiation. <i>Scientific Reports</i> , 2019, 9, 3562.	3.3	19
18	Promoter-Targeted Histone Acetylation of Chromatinized Parvoviral Genome Is Essential for the Progress of Infection. <i>Journal of Virology</i> , 2016, 90, 4059-4066.	3.4	13

#	ARTICLE	IF	CITATIONS
19	Interactions of Functional Apolipoprotein E Gene Promoter Polymorphisms With Smoking on Aortic Atherosclerosis. <i>Circulation: Cardiovascular Genetics</i> , 2008, 1, 107-116.	5.1	12
20	Lumbo-sacral neural crest derivatives fate mapped with the aid of Wnt-1 promoter integrate but are not essential to kidney development. <i>Differentiation</i> , 2009, 77, 199-208.	1.9	12
21	Transglutaminase 2-specific coeliac disease autoantibodies induce morphological changes and signs of inflammation in the small-bowel mucosa of mice. <i>Amino Acids</i> , 2017, 49, 529-540.	2.7	12
22	Targeted deletion of keratin 8 in intestinal epithelial cells disrupts tissue integrity and predisposes to tumorigenesis in the colon. <i>Cellular and Molecular Life Sciences</i> , 2022, 79, 10.	5.4	11
23	Phylogenetic analysis of the SAP30 family of transcriptional regulators reveals functional divergence in the domain that binds the nuclear matrix. <i>BMC Evolutionary Biology</i> , 2009, 9, 149.	3.2	10
24	R ² -dependent disulfide bond formation in SAP30L corepressor protein: Implications for structure and function. <i>Protein Science</i> , 2016, 25, 572-586.	7.6	9
25	Apolipoprotein A4 Defines the Villus-Crypt Border in Duodenal Specimens for Celiac Disease Morphometry. <i>Frontiers in Immunology</i> , 2021, 12, 713854.	4.8	8
26	JAZF1-SUZ12 dysregulates PRC2 function and gene expression during cell differentiation. <i>Cell Reports</i> , 2022, 39, 110889.	6.4	6
27	Polycomb Repressive Complex 2 Regulates Genes Necessary for Intestinal Microfold Cell (M Cell) Development. <i>Cellular and Molecular Gastroenterology and Hepatology</i> , 2021, 12, 873-889.	4.5	5
28	PRC2 Regulated Atoh8 Is a Regulator of Intestinal Microfold Cell (M Cell) Differentiation. <i>International Journal of Molecular Sciences</i> , 2021, 22, 9355.	4.1	5
29	Alternative mRNA splicing of SAP30L regulates its transcriptional repression activity. <i>FEBS Letters</i> , 2008, 582, 379-384.	2.8	4