

# Thomas S Griffith

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/5844300/publications.pdf>

Version: 2024-02-01

149  
papers

12,454  
citations

41627

51  
h-index

30277

107  
g-index

155  
all docs

155  
docs citations

155  
times ranked

15105  
citing authors

#	ARTICLE	IF	CITATIONS
1	Toll-like receptor 7 and 8 imidazoquinoline-based agonist/antagonist pairs. <i>Bioorganic and Medicinal Chemistry Letters</i> , 2022, 59, 128548.	1.0	4
2	Lentivirus Mediated Pancreatic Beta-Cell-Specific Insulin Gene Therapy for STZ-Induced Diabetes. <i>Molecular Therapy</i> , 2021, 29, 149-161.	3.7	10
3	Lentiviral gene therapy vectors encoding VIP suppressed diabetes-related inflammation and augmented pancreatic beta-cell proliferation. <i>Gene Therapy</i> , 2021, 28, 130-141.	2.3	9
4	Novel TLR 7/8 agonists for improving NK cell mediated antibody-dependent cellular cytotoxicity (ADCC). <i>Scientific Reports</i> , 2021, 11, 3346.	1.6	17
5	Prolonged Reactive Oxygen Species Production following Septic Insult. <i>ImmunoHorizons</i> , 2021, 5, 477-488.	0.8	14
6	Current Update on Severe Acute Respiratory Syndrome Coronavirus 2 Vaccine Development with a Special Emphasis on Gene Therapy Viral Vector Design and Construction for Vaccination. <i>Human Gene Therapy</i> , 2021, 32, 541-562.	1.4	9
7	Sepsis, Cytokine Storms, and Immunopathology: The Divide between Neonates and Adults. <i>ImmunoHorizons</i> , 2021, 5, 512-522.	0.8	14
8	Severity of Sepsis Determines the Degree of Impairment Observed in Circulatory and Tissue-Resident Memory CD8 T Cell Populations. <i>Journal of Immunology</i> , 2021, 207, 1871-1881.	0.4	10
9	Sepsis and multiple sclerosis: Causative links and outcomes. <i>Immunology Letters</i> , 2021, 238, 40-46.	1.1	5
10	NK Cell-Derived IL-10 Supports Host Survival during Sepsis. <i>Journal of Immunology</i> , 2021, 206, 1171-1180.	0.4	19
11	Autoimmunity Increases Susceptibility to and Mortality from Sepsis. <i>ImmunoHorizons</i> , 2021, 5, 844-854.	0.8	3
12	Sepsis leads to lasting changes in phenotype and function of memory CD8 T cells. <i>ELife</i> , 2021, 10, .	2.8	19
13	Inducing Experimental Polymicrobial Sepsis by Cecal Ligation and Puncture. <i>Current Protocols in Immunology</i> , 2020, 131, e110.	3.6	25
14	TLR7/8 Agonist-Loaded Nanoparticles Augment NK Cell-Mediated Antibody-Based Cancer Immunotherapy. <i>Molecular Pharmaceutics</i> , 2020, 17, 2109-2124.	2.3	28
15	New Insights into the Immune System Using Dirty Mice. <i>Journal of Immunology</i> , 2020, 205, 3-11.	0.4	59
16	Worry and FRET: ROS Production Leads to Fluorochrome Tandem Degradation and impairs Interpretation of Flow Cytometric Results. <i>Immunity</i> , 2020, 52, 419-421.	6.6	6
17	Exploiting antibody biology for the treatment of cancer. <i>Immunotherapy</i> , 2020, 12, 255-267.	1.0	7
18	CD4 T Cell Responses and the Sepsis-Induced Immunoparalysis State. <i>Frontiers in Immunology</i> , 2020, 11, 1364.	2.2	83

#	ARTICLE	IF	CITATIONS
19	Polymicrobial Sepsis Impairs Antigen-Specific Memory CD4 T Cell-Mediated Immunity. <i>Frontiers in Immunology</i> , 2020, 11, 1786.	2.2	18
20	CD8 <sup>+</sup> T cells mediate ultraviolet A-induced immunomodulation in a model of extracorporeal photochemotherapy. <i>European Journal of Immunology</i> , 2020, 50, 725-735.	1.6	6
21	Sepsis impedes EAE disease development and diminishes autoantigen-specific naive CD4 T cells. <i>ELife</i> , 2020, 9, .	2.8	16
22	Microbial Exposure Enhances Immunity to Pathogens Recognized by TLR2 but Increases Susceptibility to Cytokine Storm through TLR4 Sensitization. <i>Cell Reports</i> , 2019, 28, 1729-1743.e5.	2.9	74
23	A wild microbiome improves mouse modeling of the human immune response. <i>Lab Animal</i> , 2019, 48, 337-338.	0.2	5
24	Cytomegalovirus Evades TRAIL-Mediated Innate Lymphoid Cell 1 Defenses. <i>Journal of Virology</i> , 2019, 93, .	1.5	11
25	Sepsis-Induced State of Immunoparalysis Is Defined by Diminished CD8 T Cell-Mediated Antitumor Immunity. <i>Journal of Immunology</i> , 2019, 203, 725-735.	0.4	21
26	Cutting Edge: Polymicrobial Sepsis Has the Capacity to Reinvigorate Tumor-Infiltrating CD8 T Cells and Prolong Host Survival. <i>Journal of Immunology</i> , 2019, 202, 2843-2848.	0.4	20
27	Paradoxical effects of obesity on T cell function during tumor progression and PD-1 checkpoint blockade. <i>Nature Medicine</i> , 2019, 25, 141-151.	15.2	539
28	Poly(D,L-lactide-co-glycolide) Nanoparticles as Delivery Platforms for TLR7/8 Agonist-Based Cancer Vaccine. <i>Journal of Pharmacology and Experimental Therapeutics</i> , 2019, 370, 715-724.	1.3	38
29	Combination of Sunitinib and PD-L1 Blockade Enhances Anticancer Efficacy of TLR7/8 Agonist-Based Nanovaccine. <i>Molecular Pharmaceutics</i> , 2019, 16, 1200-1210.	2.3	30
30	HIV-based lentivirus-mediated vasoactive intestinal peptide gene delivery protects against DIO animal model of Type 2 diabetes. <i>Gene Therapy</i> , 2018, 25, 269-283.	2.3	12
31	Polymeric nanoparticles encapsulating novel TLR7/8 agonists as immunostimulatory adjuvants for enhanced cancer immunotherapy. <i>Biomaterials</i> , 2018, 164, 38-53.	5.7	133
32	Therapeutic Potential of Lentivirus-Mediated Glucagon-Like Peptide-1 Gene Therapy for Diabetes. <i>Human Gene Therapy</i> , 2018, 29, 802-815.	1.4	16
33	Biliary tract instillation of a SMAC mimetic induces TRAIL-dependent acute sclerosing cholangitis-like injury in mice. <i>Cell Death and Disease</i> , 2018, 8, e2535-e2535.	2.7	9
34	Acidic pH-responsive polymer nanoparticles as a TLR7/8 agonist delivery platform for cancer immunotherapy. <i>Nanoscale</i> , 2018, 10, 20851-20862.	2.8	59
35	Polymicrobial Sepsis Chronic Immunoparalysis Is Defined by Diminished Ag-Specific T Cell-Dependent B Cell Responses. <i>Frontiers in Immunology</i> , 2018, 9, 2532.	2.2	48
36	Polymicrobial sepsis influences NK-cell-mediated immunity by diminishing NK-cell-intrinsic receptor-mediated effector responses to viral ligands or infections. <i>PLoS Pathogens</i> , 2018, 14, e1007405.	2.1	46

#	ARTICLE	IF	CITATIONS
37	Eradication of Established Tumors by Chemically Self-Assembled Nanoring Labeled T Cells. <i>ACS Nano</i> , 2018, 12, 6563-6576.	7.3	24
38	Cutting Edge: Elevated Leptin during Diet-Induced Obesity Reduces the Efficacy of Tumor Immunotherapy. <i>Journal of Immunology</i> , 2018, 201, 1837-1841.	0.4	53
39	Sepsis-Induced T Cell Immunoparalysis: The Ins and Outs of Impaired T Cell Immunity. <i>Journal of Immunology</i> , 2018, 200, 1543-1553.	0.4	143
40	The synergy between ionizing radiation and immunotherapy in the treatment of prostate cancer. <i>Immunotherapy</i> , 2017, 9, 1005-1018.	1.0	2
41	Design and Synthesis of N1-Modified Imidazoquinoline Agonists for Selective Activation of Toll-like Receptors 7 and 8. <i>ACS Medicinal Chemistry Letters</i> , 2017, 8, 1148-1152.	1.3	32
42	A Syngeneic Mouse Model of Metastatic Renal Cell Carcinoma for Quantitative and Longitudinal Assessment of Preclinical Therapies. <i>Journal of Visualized Experiments</i> , 2017, , .	0.2	16
43	TRAIL deletion prevents liver inflammation but not adipose tissue inflammation during murine diet-induced obesity. <i>Hepatology Communications</i> , 2017, 1, 648-662.	2.0	33
44	Enteric immunity, the gut microbiome, and sepsis: Rethinking the germ theory of disease. <i>Experimental Biology and Medicine</i> , 2017, 242, 127-139.	1.1	51
45	The current status of immunobased therapies for metastatic renal-cell carcinoma. <i>ImmunoTargets and Therapy</i> , 2017, Volume 6, 83-93.	2.7	14
46	Polymicrobial sepsis impairs bystander recruitment of effector cells to infected skin despite optimal sensing and alarming function of skin resident memory CD8 T cells. <i>PLoS Pathogens</i> , 2017, 13, e1006569.	2.1	47
47	Focal Therapy for Prostate Cancer: A Molecular Biology Approach with TRAIL. <i>Current Clinical Urology</i> , 2017, , 347-354.	0.0	0
48	CD8 T Cell-Independent Antitumor Response and Its Potential for Treatment of Malignant Gliomas. <i>Cancers</i> , 2016, 8, 71.	1.7	8
49	Clinical and Experimental Sepsis Impairs CD8 T-Cell-Mediated Immunity. <i>Critical Reviews in Immunology</i> , 2016, 36, 57-74.	1.0	55
50	Gut Microbial Membership Modulates CD4 T Cell Reconstitution and Function after Sepsis. <i>Journal of Immunology</i> , 2016, 197, 1692-1698.	0.4	31
51	Polymicrobial Sepsis Diminishes Dendritic Cell Numbers and Function Directly Contributing to Impaired Primary CD8 T Cell Responses In Vivo. <i>Journal of Immunology</i> , 2016, 197, 4301-4311.	0.4	48
52	Triptolide enhances the tumoricidal activity of TRAIL against renal cell carcinoma. <i>FEBS Journal</i> , 2015, 282, 4747-4765.	2.2	15
53	Alterations in Antigen-Specific Naive CD4 T Cell Precursors after Sepsis Impairs Their Responsiveness to Pathogen Challenge. <i>Journal of Immunology</i> , 2015, 194, 1609-1620.	0.4	55
54	Polymicrobial Sepsis Increases Susceptibility to Chronic Viral Infection and Exacerbates CD8+ T Cell Exhaustion. <i>Journal of Immunology</i> , 2015, 195, 116-125.	0.4	48

#	ARTICLE	IF	CITATIONS
55	Exploiting natural anti-tumor immunity for metastatic renal cell carcinoma. <i>Human Vaccines and Immunotherapeutics</i> , 2015, 11, 1612-1620.	1.4	16
56	Therapeutic applications of TRAIL receptor agonists in cancer and beyond. , 2015, 155, 117-131.		67
57	The Frequency of Naive and Early-Activated Hapten-Specific B Cell Subsets Dictates the Efficacy of a Therapeutic Vaccine against Prescription Opioid Abuse. <i>Journal of Immunology</i> , 2015, 194, 5926-5936.	0.4	40
58	Immunosuppression after Sepsis: Systemic Inflammation and Sepsis Induce a Loss of Naïve T-Cells but No Enduring Cell-Autonomous Defects in T-Cell Function. <i>PLoS ONE</i> , 2014, 9, e115094.	1.1	52
59	Tumor necrosis factor-related apoptosis-inducing ligand-induced apoptotic pathways in cancer immunosurveillance: molecular mechanisms and prospects for therapy. <i>Research and Reports in Biochemistry</i> , 2014, , 1.	1.6	3
60	GLP-1-mediated gene therapy approaches for diabetes treatment. <i>Expert Reviews in Molecular Medicine</i> , 2014, 16, e7.	1.6	14
61	Intravascular staining for discrimination of vascular and tissue leukocytes. <i>Nature Protocols</i> , 2014, 9, 209-222.	5.5	612
62	Polymicrobial Sepsis Alters Antigen-Dependent and -Independent Memory CD8 T Cell Functions. <i>Journal of Immunology</i> , 2014, 192, 3618-3625.	0.4	58
63	Effective TRAIL-based immunotherapy requires both plasmacytoid and CD8 <sup>+</sup> dendritic cells. <i>Cancer Immunology, Immunotherapy</i> , 2014, 63, 685-697.	2.0	19
64	Cellular Inhibitor of Apoptosis Protein cIAP2 Protects against Pulmonary Tissue Necrosis during Influenza Virus Infection to Promote Host Survival. <i>Cell Host and Microbe</i> , 2014, 15, 23-35.	5.1	141
65	Impact of sepsis on CD4 T cell immunity. <i>Journal of Leukocyte Biology</i> , 2014, 96, 767-777.	1.5	128
66	CpG-mediated modulation of MDSC contributes to the efficacy of Ad5-TRAIL therapy against renal cell carcinoma. <i>Cancer Immunology, Immunotherapy</i> , 2014, 63, 1213-1227.	2.0	32
67	Minimal changes in the systemic immune response after nephrectomy of localized renal masses <sup>11</sup> This work was supported by the University of Iowa Carver College of Medicine/Department of Urology Investigator Start-up Funds, NIH Grant CA181088-01 (to L.A.N.), and NIH Grant CA109446 (to T.S.G.). <i>Urologic Oncology: Seminars and Original Investigations</i> . 2014, 32, 589-600.	0.8	19
68	PMN and anti-tumor immunity <sup>11</sup> The case of bladder cancer immunotherapy. <i>Seminars in Cancer Biology</i> , 2013, 23, 183-189.	4.3	38
69	Sustained and Incomplete Recovery of Naive CD8 <sup>+</sup> T Cell Precursors after Sepsis Contributes to Impaired CD8 <sup>+</sup> T Cell Responses to Infection. <i>Journal of Immunology</i> , 2013, 190, 1991-2000.	0.4	73
70	Clinical utility of insulin and insulin analogs. <i>Islets</i> , 2013, 5, 67-78.	0.9	40
71	T-Cell-Mediated Immunity and the Role of TRAIL in Sepsis-Induced Immunosuppression. <i>Critical Reviews in Immunology</i> , 2013, 33, 23-40.	1.0	43
72	Diet-Induced Obesity Alters Dendritic Cell Function in the Presence and Absence of Tumor Growth. <i>Journal of Immunology</i> , 2012, 189, 1311-1321.	0.4	94

#	ARTICLE	IF	CITATIONS
73	Therapeutic potential of VIP vs PACAP in diabetes. <i>Journal of Molecular Endocrinology</i> , 2012, 49, R157-R167.	1.1	41
74	Activation of systemic antitumor immunity via TRAIL-induced apoptosis. <i>Oncolimmunology</i> , 2012, 1, 1178-1180.	2.1	4
75	TNF-related apoptosis-inducing ligand (TRAIL) exerts therapeutic efficacy for the treatment of pneumococcal pneumonia in mice. <i>Journal of Experimental Medicine</i> , 2012, 209, 1937-1952.	4.2	79
76	Eradication of Metastatic Renal Cell Carcinoma after Adenovirus-Encoded TNF-Related Apoptosis-Inducing Ligand (TRAIL)/CpG Immunotherapy. <i>PLoS ONE</i> , 2012, 7, e31085.	1.1	46
77	Cell Death in the Maintenance and Abrogation of Tolerance: The Five Ws of Dying Cells. <i>Immunity</i> , 2011, 35, 456-466.	6.6	86
78	Description of a Novel Murine Model for Ileocystoplasty and Early Histologic Changes. <i>Scientific World Journal</i> , The, 2011, 11, 1325-1331.	0.8	2
79	Sensitization of human bladder tumor cells to TNF-related apoptosis-inducing ligand (TRAIL)-induced apoptosis with a small molecule IAP antagonist. <i>Apoptosis: an International Journal on Programmed Cell Death</i> , 2011, 16, 13-26.	2.2	28
80	Tracing of islet graft survival by way of <i>in vivo</i> fluorescence imaging. <i>Diabetes/Metabolism Research and Reviews</i> , 2011, 27, 575-583.	1.7	11
81	Immune Unresponsiveness to Secondary Heterologous Bacterial Infection after Sepsis Induction Is TRAIL Dependent. <i>Journal of Immunology</i> , 2011, 187, 2148-2154.	0.4	56
82	The Magnitude of the T Cell Response to a Clinically Significant Dose of Influenza Virus Is Regulated by TRAIL. <i>Journal of Immunology</i> , 2011, 187, 4581-4588.	0.4	36
83	Systemic Immunological Tolerance to Ocular Antigens Is Mediated by TRAIL-Expressing CD8+ T Cells. <i>Journal of Immunology</i> , 2011, 186, 791-798.	0.4	24
84	The Plasticity of Regulatory T Cell Function. <i>Journal of Immunology</i> , 2011, 187, 4987-4997.	0.4	58
85	Advances in Viral Vector-Based TRAIL Gene Therapy for Cancer. <i>Cancers</i> , 2011, 3, 603-620.	1.7	11
86	TRAIL-expressing CD8+ T cells mediate tolerance following soluble peptide-induced peripheral T cell deletion. <i>Journal of Leukocyte Biology</i> , 2010, 88, 1217-1225.	1.5	18
87	Sepsis-Induced Apoptosis Leads to Active Suppression of Delayed-Type Hypersensitivity by CD8+ Regulatory T Cells through a TRAIL-Dependent Mechanism. <i>Journal of Immunology</i> , 2010, 184, 6766-6772.	0.4	63
88	Micro-recanalization in a biodegradable graft for reconstruction of the vas deferens is enhanced by sildenafil citrate. <i>Asian Journal of Andrology</i> , 2010, 12, 814-818.	0.8	1
89	Conatumumab, a fully human mAb against death receptor 5 for the treatment of cancer. <i>Current Opinion in Investigational Drugs</i> , 2010, 11, 688-98.	2.3	22
90	Adenovirus-Mediated TRAIL Gene (Ad5hTRAIL) Delivery into Pancreatic Islets Prolongs Normoglycemia in Streptozotocin-Induced Diabetic Rats. <i>Human Gene Therapy</i> , 2009, 20, 1177-1189.	1.4	31

#	ARTICLE	IF	CITATIONS
91	Activation-Induced CD154 Expression Abrogates Tolerance Induced by Apoptotic Cells. <i>Journal of Immunology</i> , 2009, 183, 6114-6123.	0.4	24
92	TNF-related apoptosis-inducing ligand (TRAIL): A new path to anti-cancer therapies. <i>European Journal of Pharmacology</i> , 2009, 625, 63-72.	1.7	163
93	The role of neutrophils and TNF-related apoptosis-inducing ligand (TRAIL) in bacillus Calmette-Guérin (BCG) immunotherapy for urothelial carcinoma of the bladder. <i>Cancer and Metastasis Reviews</i> , 2009, 28, 345-353.	2.7	44
94	Early microrecanalization of vas deferens following biodegradable graft implantation in bilaterally vasectomized rats. <i>Asian Journal of Andrology</i> , 2009, 11, 373-378.	0.8	6
95	The Use of Immunofluorescence in Microdissection Testicular Sperm Extraction. <i>Journal of Andrology</i> , 2009, 30, 548-551.	2.0	10
96	TRAIL-Deficient Mice Exhibit Delayed Regression of Retinal Neovascularization. <i>American Journal of Pathology</i> , 2009, 175, 2697-2708.	1.9	17
97	High TRAIL Death Receptor 4 and Decoy Receptor 2 Expression Correlates With Significant Cell Death in Pancreatic Ductal Adenocarcinoma Patients. <i>Pancreas</i> , 2009, 38, 154-160.	0.5	30
98	TRAIL Gene Therapy: From Preclinical Development to Clinical Application. <i>Current Gene Therapy</i> , 2009, 9, 9-19.	0.9	84
99	Induction of Tumor Cell Apoptosis by TRAIL Gene Therapy. <i>Methods in Molecular Biology</i> , 2009, 542, 315-334.	0.4	4
100	Molecular mechanisms of death ligand-mediated immune modulation: A gene therapy model to prolong islet survival in type 1 diabetes. <i>Journal of Cellular Biochemistry</i> , 2008, 104, 710-720.	1.2	27
101	Influenza-induced expression of functional tumor necrosis factor-related apoptosis-inducing ligand on human peripheral blood mononuclear cells. <i>Human Immunology</i> , 2008, 69, 634-646.	1.2	24
102	Tumor necrosis factor-related apoptosis inducing ligand-R4 decoy receptor expression is correlated with high Gleason scores, prostate-specific antigen recurrence, and decreased survival in patients with prostate carcinoma. <i>Urologic Oncology: Seminars and Original Investigations</i> , 2008, 26, 158-165.	0.8	33
103	Role of neutrophils in BCG immunotherapy for bladder cancer. <i>Urologic Oncology: Seminars and Original Investigations</i> , 2008, 26, 341-345.	0.8	100
104	TNF-related apoptosis-inducing ligand (TRAIL) is expressed throughout myeloid development, resulting in a broad distribution among neutrophil granules. <i>Journal of Leukocyte Biology</i> , 2008, 83, 621-629.	1.5	26
105	CD8 T Cells Utilize TRAIL to Control Influenza Virus Infection. <i>Journal of Immunology</i> , 2008, 181, 4918-4925.	0.4	176
106	High Levels of Endogenous Tumor Necrosis Factor-Related Apoptosis-Inducing Ligand Expression Correlate With Increased Cell Death in Human Pancreas. <i>Pancreas</i> , 2008, 36, 385-393.	0.5	27
107	Apoptotic Cells Induce Tolerance by Generating Helpless CD8+ T Cells That Produce TRAIL. <i>Journal of Immunology</i> , 2007, 178, 2679-2687.	0.4	81
108	Identification of the Mycobacterial Subcomponents Involved in the Release of Tumor Necrosis Factor-Related Apoptosis-Inducing Ligand from Human Neutrophils. <i>Infection and Immunity</i> , 2007, 75, 1265-1271.	1.0	39



#	ARTICLE	IF	CITATIONS
109	Activation of Tumor-Specific CD8+ T Cells after Intratumoral Ad5-TRAIL/CpG Oligodeoxynucleotide Combination Therapy. <i>Cancer Research</i> , 2007, 67, 11980-11990.	0.4	45
110	Cytomegalovirus and the role of interferon in the expression of tumor necrosis factor-related apoptosis-inducing ligand in the placenta. <i>American Journal of Obstetrics and Gynecology</i> , 2007, 197, 608.e1-608.e6.	0.7	8
111	Histone deacetylase inhibitors enhance Ad5-TRAIL killing of TRAIL-resistant prostate tumor cells through increased caspase-2 activity. <i>Apoptosis: an International Journal on Programmed Cell Death</i> , 2007, 12, 561-571.	2.2	45
112	Neutrophils and TRAIL: insights into BCG immunotherapy for bladder cancer. <i>Immunologic Research</i> , 2007, 39, 79-93.	1.3	39
113	Survivin inhibits apoptosis induced by TRAIL, and the ratio between survivin and TRAIL receptors is predictive of recurrent disease in neuroblastoma. <i>Journal of Pediatric Surgery</i> , 2006, 41, 1431-1440.	0.8	29
114	<i>Neisseria gonorrhoeae</i> delays the onset of apoptosis in polymorphonuclear leukocytes. <i>Cellular Microbiology</i> , 2006, 8, 1780-1790.	1.1	49
115	A vision of cell death: Fas ligand and immune privilege 10 years later. <i>Immunological Reviews</i> , 2006, 213, 228-238.	2.8	101
116	Induction of protective immunity to RM-1 prostate cancer cells with ALVAC-IL-2/IL-12/TNF- $\beta$ combination therapy. <i>International Journal of Cancer</i> , 2006, 119, 2632-2641.	2.3	15
117	TRAIL Deficiency Delays, but Does Not Prevent, Erosion in the Quality of "Helpless" Memory CD8 T Cells. <i>Journal of Immunology</i> , 2006, 177, 999-1006.	0.4	56
118	Histone Deacetylase Inhibitors Modulate the Sensitivity of Tumor Necrosis Factor-Related Apoptosis-Inducing Ligand-Resistant Bladder Tumor Cells. <i>Cancer Research</i> , 2006, 66, 499-507.	0.4	80
119	CD4+ T-cell help controls CD8+ T-cell memory via TRAIL-mediated activation-induced cell death. <i>Nature</i> , 2005, 434, 88-93.	13.7	547
120	Depsipeptide (FR901228) Enhances the Cytotoxic Activity of TRAIL by Redistributing TRAIL Receptor to Membrane Lipid Rafts. <i>Molecular Therapy</i> , 2005, 11, 542-552.	3.7	81
121	Immunostimulatory oligodeoxynucleotides induce apoptosis of B cell chronic lymphocytic leukemia cells. <i>Journal of Leukocyte Biology</i> , 2005, 77, 378-387.	1.5	90
122	Histone deacetylase inhibitors modulate renal cell carcinoma sensitivity to TRAIL/Apo-2L-induced apoptosis by enhancing TRAIL-R2 expression. <i>Cancer Biology and Therapy</i> , 2005, 4, 1104-1112.	1.5	59
123	Neutrophil stimulation with <i>Mycobacterium bovis</i> bacillus Calmette-Guèrin (BCG) results in the release of functional soluble TRAIL/Apo-2L. <i>Blood</i> , 2005, 106, 3474-3482.	0.6	112
124	Human B Cells Express Functional TRAIL/Apo-2 Ligand after CpG-Containing Oligodeoxynucleotide Stimulation. <i>Journal of Immunology</i> , 2004, 173, 892-899.	0.4	95
125	Tumor Necrosis Factor-Related Apoptosis-Inducing Ligand. <i>Cancer Research</i> , 2004, 64, 3386-3390.	0.4	167
126	Inhibition of the NF- $\kappa$ B pathway enhances TRAIL-mediated apoptosis in neuroblastoma cells. <i>Cancer Gene Therapy</i> , 2004, 11, 681-690.	2.2	54



#	ARTICLE	IF	CITATIONS
127	Expression of TNF-related apoptosis-inducing ligand (TRAIL) in megakaryocytes and platelets. <i>Experimental Hematology</i> , 2004, 32, 1073-1081.	0.2	38
128	The topoisomerase I inhibitor topotecan increases the sensitivity of prostate tumor cells to TRAIL/Apo-2L-induced apoptosis. <i>Cancer Chemotherapy and Pharmacology</i> , 2003, 52, 175-184.	1.1	14
129	Apoptosis, tolerance, and regulatory T cells - old wine, new wineskins. <i>Immunological Reviews</i> , 2003, 193, 111-123.	2.8	55
130	Structure/Function Analysis of the Murine CD95L Promoter Reveals the Identification of a Novel Transcriptional Repressor and Functional CD28 Response Element. <i>Journal of Biological Chemistry</i> , 2003, 278, 35950-35958.	1.6	22
131	Plasmacytoid Dendritic Cell-Derived IFN- $\gamma$ Induces TNF-Related Apoptosis-Inducing Ligand/Apo-2L-Mediated Antitumor Activity by Human Monocytes Following CpG Oligodeoxynucleotide Stimulation. <i>Journal of Immunology</i> , 2003, 171, 212-218.	0.4	67
132	Uptake of Apoptotic Antigen-Coupled Cells by Lymphoid Dendritic Cells and Cross-Priming of CD8+ T Cells Produce Active Immune Unresponsiveness. <i>Journal of Immunology</i> , 2002, 168, 5589-5595.	0.4	174
133	TRAIL: A Mechanism of Tumor Surveillance in an Immune Privileged Site. <i>Journal of Immunology</i> , 2002, 169, 4739-4744.	0.4	95
134	Induction and regulation of tumor necrosis factor-related apoptosis-inducing ligand/Apo-2 ligand-mediated apoptosis in renal cell carcinoma. <i>Cancer Research</i> , 2002, 62, 3093-9.	0.4	60
135	Induction of glioblastoma apoptosis using neural stem cell-mediated delivery of tumor necrosis factor-related apoptosis-inducing ligand. <i>Cancer Research</i> , 2002, 62, 7170-4.	0.4	201
136	Regulation of Fas Ligand-Induced Apoptosis by TNF. <i>Journal of Immunology</i> , 2001, 167, 3049-3056.	0.4	62
137	Inhibition of Murine Prostate Tumor Growth and Activation of Immunoregulatory Cells With Recombinant Canarypox Viruses. <i>Journal of the National Cancer Institute</i> , 2001, 93, 998-1007.	3.0	28
138	Suppression of Tumor Growth Following Intralesional Therapy with TRAIL Recombinant Adenovirus. <i>Molecular Therapy</i> , 2001, 4, 257-266.	3.7	90
139	Adenoviral-Mediated Transfer of the TNF-Related Apoptosis-Inducing Ligand/Apo-2 Ligand Gene Induces Tumor Cell Apoptosis. <i>Journal of Immunology</i> , 2000, 165, 2886-2894.	0.4	184
140	Tumoricidal activity of tumor necrosis factor-related apoptosis-inducing ligand in vivo. <i>Nature Medicine</i> , 1999, 5, 157-163.	15.2	2,377
141	Monocyte-mediated Tumoricidal Activity via the Tumor Necrosis Factor-related Cytokine, TRAIL. <i>Journal of Experimental Medicine</i> , 1999, 189, 1343-1354.	4.2	442
142	Human Dendritic Cells Mediate Cellular Apoptosis via Tumor Necrosis Factor-Related Apoptosis-Inducing Ligand (Trail). <i>Journal of Experimental Medicine</i> , 1999, 190, 1155-1164.	4.2	369
143	TRAIL: a molecule with multiple receptors and control mechanisms. <i>Current Opinion in Immunology</i> , 1998, 10, 559-563.	2.4	436
144	Inducible Nonlymphoid Expression of Fas Ligand Is Responsible for Superantigen-Induced Peripheral Deletion of T Cells. <i>Immunity</i> , 1998, 9, 711-720.	6.6	145

#	ARTICLE	IF	CITATIONS
145	Antiinflammatory Effects of CD95 Ligand (FasL)-induced Apoptosis. <i>Journal of Experimental Medicine</i> , 1998, 188, 887-896.	4.2	208
146	A vision of cell death: insights into immune privilege. <i>Immunological Reviews</i> , 1997, 156, 167-184.	2.8	167
147	Cell death and the immune response: a lesson from the privileged. <i>Journal of Clinical Immunology</i> , 1997, 17, 1-10.	2.0	15
148	CD95-Induced Apoptosis of Lymphocytes in an Immune Privileged Site Induces Immunological Tolerance. <i>Immunity</i> , 1996, 5, 7-16.	6.6	366
149	The immune response and the eye. TCR $\alpha$ -chain related molecules regulate the systemic immunity to antigen presented in the eye. <i>International Immunology</i> , 1995, 7, 1617-1625.	1.8	29