Ignacio Santa-Regina

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/5894986/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	Identification of Canola Roots Endophytic Bacteria and Analysis of Their Potential as Biofertilizers for Canola Crops with Special Emphasis on Sporulating Bacteria. Agronomy, 2021, 11, 1796.	3.0	15
2	TRY plant trait database – enhanced coverage and open access. Global Change Biology, 2020, 26, 119-188.	9.5	1,038
3	Pseudomonas edaphica sp. nov., isolated from rhizospheric soil of Cistus ladanifer L. in Spain. International Journal of Systematic and Evolutionary Microbiology, 2019, 69, 3141-3147.	1.7	13
4	Culturable bacterial diversity from the chestnut (Castanea sativa Mill.) phyllosphere and antagonism against the fungi causing the chestnut blight and ink diseases. AIMS Microbiology, 2017, 3, 293-314.	2.2	11
5	BAAD: a Biomass And Allometry Database for woody plants. Ecology, 2015, 96, 1445-1445.	3.2	122
6	Aboveground Biomass and Nutrient Pools in Two Evergreen Oak Stands of the Middle Moroccan Atlas Area. Arid Land Research and Management, 2013, 27, 188-202.	1.6	2
7	Adaptation, tolerance, and evolution of plant species in a pyrite mine in response to contamination level and properties of mine tailings: sustainable rehabilitation. Journal of Soils and Sediments, 2013, 13, 730-741.	3.0	65
8	Zinc, cadmium and thallium distribution in soils and plants of an area impacted by sphalerite-bearing mine wastes. Geoderma, 2013, 207-208, 25-34.	5.1	45
9	Mobility and phytoavailability of antimony in an area impacted by a former stibnite mine exploitation. Science of the Total Environment, 2013, 449, 260-268.	8.0	22
10	Antimony, arsenic and lead distribution in soils and plants of an agricultural area impacted by former mining activities. Science of the Total Environment, 2012, 439, 35-43.	8.0	74
11	Mobility and phytoavailability of arsenic in an abandoned mining area. Geoderma, 2011, 166, 153-161.	5.1	22
12	Arsenic distribution in soils and plants of an arsenic impacted former mining area. Environmental Pollution, 2011, 159, 2637-2647.	7.5	41
13	Diversity and community structure of culturable arsenic-resistant bacteria across a soil arsenic gradient at an abandoned tungsten–tin mining area. Chemosphere, 2011, 85, 129-134.	8.2	39
14	Arsenic, antimony, and other trace element contamination in a mine tailings affected area and uptake by tolerant plant species. Environmental Geochemistry and Health, 2011, 33, 353-362.	3.4	65
15	Effects of different drying processes on the concentrations of metals and metalloids in plant materials. Journal of Radioanalytical and Nuclear Chemistry, 2011, 289, 29-34.	1.5	8
16	Correlation among soil enzyme activities under different forest system management practices. Ecological Engineering, 2011, 37, 1123-1131.	3.6	165
17	Acinetobacter strains IH9 and OCI1, two rhizospheric phosphate solubilizing isolates able to promote plant growth, constitute a new genomovar of Acinetobacter calcoaceticus. Systematic and Applied Microbiology, 2009, 32, 334-341.	2.8	20
18	Mercury in soils and plants in an abandoned cinnabar mining area (SW Spain). Journal of Hazardous Materials, 2009, 168, 1319-1324.	12.4	55

#	Article	IF	CITATIONS
19	Evaluation of various chemical extraction methods to estimate plant-available arsenic in mine soils. Chemosphere, 2008, 70, 1459-1467.	8.2	84
20	Paenibacillus castaneae sp. nov., isolated from the phyllosphere of Castanea sativa Miller. International Journal of Systematic and Evolutionary Microbiology, 2008, 58, 2560-2564.	1.7	29
21	Cadmium and zinc in polluted mining soils and uptake by plants (El Losar mine, Spain). International Journal of Environment and Pollution, 2008, 33, 146.	0.2	18
22	Long-term effectiveness of sowing high and low diversity seed mixtures to enhance plant community development on ex-arable fields. Applied Vegetation Science, 2007, 10, 97.	1.9	36
23	Antimony and Arsenic Uptake by Plants in an Abandoned Mining Area. Communications in Soil Science and Plant Analysis, 2007, 38, 1255-1275.	1.4	49
24	Reclassification of Pseudomonas aurantiaca as a synonym of Pseudomonas chlororaphis and proposal of three subspecies, P. chlororaphis subsp. chlororaphis subsp. nov., P. chlororaphis subsp. aureofaciens subsp. nov., comb. nov. and P. chlororaphis subsp. aurantiaca subsp. nov., comb. nov International Journal of Systematic and Evolutionary Microbiology, 2007, 57, 1286-1290.	1.7	99
25	Longâ€ŧerm effectiveness of sowing high and low diversity seed mixtures to enhance plant community development on exâ€∎rable fields. Applied Vegetation Science, 2007, 10, 97-110.	1.9	93
26	Arsenic Bioavailability in Polluted Mining Soils and Uptake by Tolerant Plants (El Cabaco mine, Spain). Bulletin of Environmental Contamination and Toxicology, 2007, 79, 29-35.	2.7	30
27	Natural Diversity of Nodular Microsymbionts of Alnus glutinosa in the Tormes River Basin. Plant and Soil, 2006, 280, 373-383.	3.7	12
28	Forest Management and Plant Species Diversity in Chestnut Stands of Three Mediterranean Areas. Biodiversity and Conservation, 2006, 15, 1129-1142.	2.6	31
29	Burkholderia ferrariae sp. nov., isolated from an iron ore in Brazil. International Journal of Systematic and Evolutionary Microbiology, 2006, 56, 2421-2425.	1.7	45
30	Pseudomonas lutea sp. nov., a novel phosphate-solubilizing bacterium isolated from the rhizosphere of grasses. International Journal of Systematic and Evolutionary Microbiology, 2004, 54, 847-850.	1.7	59
31	Plant species diversity, plant biomass and responses of the soil community on abandoned land across Europe: idiosyncracy or above-belowground time lags. Oikos, 2003, 103, 45-58.	2.7	204
32	Nutrient pools to the soil through organic matter and throughfall under a Scots pine plantation in the Sierra de la Demanda, Spain. European Journal of Soil Biology, 2001, 37, 125-133.	3.2	30
33	Separating the chance effect from other diversity effects in the functioning of plant communities. Oikos, 2001, 92, 123-134.	2.7	132
34	Nutrient cycling in a natural beech forest and adjacent planted pine in northern Spain. Forestry, 2001, 74, 11-28.	2.3	49
35	Litter fall, decomposition and nutrient release in three semi-arid forests of the Duero basin, Spain. Forestry, 2001, 74, 347-358.	2.3	34
36	Foliar nutrient dynamics and nutrient-use efficiency in Castanea sativa coppice stands of southern Europe. Forestry, 2001, 74, 1-10.	2.3	20

Ignacio Santa-Regina

#	Article	IF	CITATIONS
37	Organic matter and nitrogen dynamics in a mature forest of common beech in the Sierra de la Demanda, Spain. Annals of Forest Science, 2001, 58, 301-314.	2.0	20
38	Nutrient Return to the Soil Through Litterfall and Throughfall Under Beech and Pine Stands of Sierra de la Demanda, Spain. Arid Land Research and Management, 2000, 14, 239-252.	0.3	17
39	Biomass estimation and nutrient pools in four Quercus pyrenaica in Sierra de Gata Mountains, Salamanca, Spain. Forest Ecology and Management, 2000, 132, 127-141.	3.2	41
40	Biomass, nutrient content, litterfall and nutrient return to the soil in Mediterranean oak forests. Forest Ecology and Management, 1999, 119, 39-49.	3.2	97
41	Organic matter dynamics in beech and pine stands of mountainous Mediterranean climate area. Annales Des Sciences Forestières, 1999, 56, 667-677.	1.2	11
42	Nutrient cycling in deciduous forest ecosystems of the Sierra de Gata mountains: aboveground litter production and potential nutrient return. Annales Des Sciences Forestières, 1998, 55, 749-769.	1.2	16
43	Nutrient cycling in deciduous forest ecosystems of the Sierra de Gata mountains: nutrient supplies to the soil through both litter and throughfall. Annales Des Sciences Forestières, 1998, 55, 771-784.	1.2	24
44	Title is missing!. Plant Ecology, 1997, 133, 49-56.	1.6	6