

Diego Moretti

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/5920643/publications.pdf>

Version: 2024-02-01

76
papers

3,144
citations

201674

27
h-index

161849

54
g-index

77
all docs

77
docs citations

77
times ranked

3581
citing authors

#	ARTICLE	IF	CITATIONS
1	Methodological Considerations for Investigating Iron Status and Regulation in Exercise and Sport Science Studies. <i>International Journal of Sport Nutrition and Exercise Metabolism</i> , 2022, 32, 359-370.	2.1	5
2	Iron Bioavailability from Infant Cereals Containing Whole Grains and Pulses: A Stable Isotope Study in Malawian Children. <i>Journal of Nutrition</i> , 2022, 152, 826-834.	2.9	3
3	The effect of a natural polyphenol supplement on iron absorption in adults with hereditary hemochromatosis. <i>European Journal of Nutrition</i> , 2022, 61, 2967-2977.	3.9	6
4	Measurement of long-term iron absorption and loss during iron supplementation using a stable isotope of iron (⁵⁷ Fe). <i>British Journal of Haematology</i> , 2021, 192, 179-189.	2.5	15
5	Direct assessment of body iron balance in women with and without iron supplementation using a long-term isotope dilution method in Benin and Switzerland. <i>American Journal of Clinical Nutrition</i> , 2021, 113, 1657-1669.	4.7	3
6	Iron homeostasis during anemia of inflammation: a prospective study of patients with tuberculosis. <i>Blood</i> , 2021, 138, 1293-1303.	1.4	20
7	Isotopic measurement of iron requirements in sub-Saharan African children. <i>American Journal of Clinical Nutrition</i> , 2021, 114, 986-996.	4.7	3
8	Threshold ferritin and hepcidin concentrations indicating early iron deficiency in young women based on upregulation of iron absorption. <i>EClinicalMedicine</i> , 2021, 39, 101052.	7.1	38
9	The <i>TM6SS6</i> variant (SNP rs855791) affects iron metabolism and oral iron absorption – a stable iron isotope study in Taiwanese women. <i>Haematologica</i> , 2021, 106, 2897-2905.	3.5	8
10	A Natural Low Phytic Acid Finger Millet Accession Significantly Improves Iron Bioavailability in Indian Women. <i>Frontiers in Nutrition</i> , 2021, 8, 791392.	3.7	2
11	Higher Extrusion Temperature Induces Greater Formation of Less Digestible Type V and Retrograded Starch in Iron-Fortified Rice Grains But Does Not Affect Iron Bioavailability: Stable Isotope Studies in Young Women. <i>Journal of Nutrition</i> , 2021, , .	2.9	2
12	The bioavailability of iron picolinate is comparable to iron sulfate when fortified into a complementary fruit yogurt: a stable iron isotope study in young women. <i>European Journal of Nutrition</i> , 2020, 59, 1371-1378.	3.9	7
13	Iron absorption from supplements is greater with alternate day than with consecutive day dosing in iron-deficient anemic women. <i>Haematologica</i> , 2020, 105, 1232-1239.	3.5	113
14	Oral iron supplementation in iron-deficient women: How much and how often?. <i>Molecular Aspects of Medicine</i> , 2020, 75, 100865.	6.4	64
15	Iron bioavailability from bouillon fortified with a novel ferric phytate compound: a stable iron isotope study in healthy women (part II). <i>Scientific Reports</i> , 2020, 10, 5339.	3.3	13
16	Polyphenol-rich tea decreases iron absorption from fortified wheat bread in Senegalese mother-child pairs and bioavailability of ferrous fumarate is sharply lower in children. <i>European Journal of Clinical Nutrition</i> , 2020, 74, 1221-1228.	2.9	12
17	A heat-stable microparticle platform for oral micronutrient delivery. <i>Science Translational Medicine</i> , 2019, 11, .	12.4	20
18	Asymptomatic <i>Helicobacter Pylori</i> Infection in Preschool Children and Young Women Does Not Predict Iron Bioavailability from Iron-Fortified Foods. <i>Nutrients</i> , 2019, 11, 2093.	4.1	8

#	ARTICLE	IF	CITATIONS
19	Zinc Absorption From Agronomically Biofortified Wheat Is Similar to Post-Harvest Fortified Wheat and Is a Substantial Source of Bioavailable Zinc in Humans. <i>Journal of Nutrition</i> , 2019, 149, 840-846.	2.9	32
20	High Bioavailability from Ferric Pyrophosphate-Fortified Bouillon Cubes in Meals is Not Increased by Sodium Pyrophosphate: a Stable Iron Isotope Study in Young Nigerian Women. <i>Journal of Nutrition</i> , 2019, 149, 723-729.	2.9	4
21	The Importance of Iron Status for Young Children in Low- and Middle-Income Countries: A Narrative Review. <i>Pharmaceuticals</i> , 2019, 12, 59.	3.8	36
22	Micronutrient-fortified rice can be a significant source of dietary bioavailable iron in schoolchildren from rural Ghana. <i>Science Advances</i> , 2019, 5, eaau0790.	10.3	18
23	Novel approaches to oral iron treatment. <i>HemaSphere</i> , 2019, 3, 109-111.	2.7	1
24	The Impact of Morning versus Afternoon Exercise on Iron Absorption in Athletes. <i>Medicine and Science in Sports and Exercise</i> , 2019, 51, 2147-2155.	0.4	32
25	Iron-containing micronutrient powders modify the effect of oral antibiotics on the infant gut microbiome and increase post-antibiotic diarrhoea risk: a controlled study in Kenya. <i>Gut</i> , 2019, 68, 645-653.	12.1	40
26	The opposing effects of acute inflammation and iron deficiency anemia on serum hepcidin and iron absorption in young women. <i>Haematologica</i> , 2019, 104, 1143-1149.	3.5	41
27	Optimization Routines for Enforcing One-to-One Matches in Record Linkage Problems. <i>R Journal</i> , 2019, 11, 185.	1.8	0
28	Athlete Iron Consumption: Timing Is Everything, But When Is Best?. <i>Medicine and Science in Sports and Exercise</i> , 2019, 51, 295-296.	0.4	0
29	Sensitivity of fatty acid desaturation and elongation to plasma zinc concentration: a randomised controlled trial in Beninese children. <i>British Journal of Nutrition</i> , 2018, 119, 610-619.	2.3	33
30	Combining food-based dietary recommendations using <sc>Optifood</sc> with zinc-fortified water potentially improves nutrient adequacy among 4- to 6-year-old children in <sc>Kisumu West</sc> district, <sc>Kenya</sc>. <i>Maternal and Child Nutrition</i> , 2018, 14, e12515.	3.0	15
31	An intensified training schedule in recreational male runners is associated with increases in erythropoiesis and inflammation and a net reduction in plasma hepcidin. <i>American Journal of Clinical Nutrition</i> , 2018, 108, 1324-1333.	4.7	22
32	Bouillon Cubes. , 2018, , 159-165.		10
33	Effectiveness of zinc-fortified water on zinc intake, status and morbidity in Kenyan pre-school children: a randomised controlled trial. <i>Public Health Nutrition</i> , 2018, 21, 2855-2865.	2.2	8
34	Iron Bioavailability from Ferric Pyrophosphate in Extruded Rice Cofortified with Zinc Sulfate Is Greater than When Cofortified with Zinc Oxide in a Human Stable Isotope Study. <i>Journal of Nutrition</i> , 2017, 147, jn241778.	2.9	10
35	Comparing intake estimations based on food composition data with chemical analysis in Malian women. <i>Public Health Nutrition</i> , 2017, 20, 1351-1361.	2.2	5
36	Zinc Absorption from Milk Is Affected by Dilution but Not by Thermal Processing, and Milk Enhances Absorption of Zinc from High-Phytate Rice in Young Dutch Women. <i>Journal of Nutrition</i> , 2017, 147, 1086-1093.	2.9	9

#	ARTICLE	IF	CITATIONS
37	The effect of lipids, a lipid-rich ready-to-use therapeutic food, or a phytase on iron absorption from maize-based meals fortified with micronutrient powders. <i>American Journal of Clinical Nutrition</i> , 2017, 105, ajcn142976.	4.7	6
38	Duration of exclusive breastfeeding is a positive predictor of iron status in 6- to 10-month-old infants in rural Kenya. <i>Maternal and Child Nutrition</i> , 2017, 13, .	3.0	20
39	Prediction of human iron bioavailability using rapid c-ELISAs for human plasma hepcidin. <i>Clinical Chemistry and Laboratory Medicine</i> , 2017, 55, 1186-1192.	2.3	6
40	Iron absorption from oral iron supplements given on consecutive versus alternate days and as single morning doses versus twice-daily split dosing in iron-depleted women: two open-label, randomised controlled trials. <i>Lancet Haematology</i> , 2017, 4, e524-e533.	4.6	276
41	Cold Extrusion but Not Coating Affects Iron Bioavailability from Fortified Rice in Young Women and Is Associated with Modifications in Starch Microstructure and Mineral Retention during Cooking. <i>Journal of Nutrition</i> , 2017, 147, 2319-2325.	2.9	8
42	Consumption of galacto-oligosaccharides increases iron absorption from a micronutrient powder containing ferrous fumarate and sodium iron EDTA: a stable-isotope study in Kenyan infants. <i>American Journal of Clinical Nutrition</i> , 2017, 106, 1020-1031.	4.7	61
43	Prebiotic galacto-oligosaccharides mitigate the adverse effects of iron fortification on the gut microbiome: a randomised controlled study in Kenyan infants. <i>Gut</i> , 2017, 66, 1956-1967.	12.1	123
44	Plant-Based Diets and Iron Status. , 2017, , 715-727.		8
45	A novel, high precision multiple-meal stable isotope method to compare iron absorption from extruded FePP-fortified rice containing different zinc compounds, citric acid/trisodium citrate and EDTA in Ghanaian children. <i>FASEB Journal</i> , 2017, 31, 436.5.	0.5	0
46	In Rwandese Women with Low Iron Status, Iron Absorption from Low-Phytic Acid Beans and Biofortified Beans Is Comparable, but Low-Phytic Acid Beans Cause Adverse Gastrointestinal Symptoms. <i>Journal of Nutrition</i> , 2016, 146, 970-975.	2.9	35
47	Cofortification of ferric pyrophosphate and citric acid/trisodium citrate into extruded rice grains doubles iron bioavailability through in situ generation of soluble ferric pyrophosphate citrate complexes. <i>American Journal of Clinical Nutrition</i> , 2016, 103, 1252-1259.	4.7	28
48	The effects of fat loss after bariatric surgery on inflammation, serum hepcidin, and iron absorption: a prospective 6-mo iron stable isotope study. <i>American Journal of Clinical Nutrition</i> , 2016, 104, 1030-1038.	4.7	38
49	The donation interval of 56 days requires extension to 180 days for whole blood donors to recover from changes in iron metabolism. <i>Blood</i> , 2016, 128, 2185-2188.	1.4	44
50	Efficacy and safety of hepcidin-based screen-and-treat approaches using two different doses versus a standard universal approach of iron supplementation in young children in rural Gambia: a double-blind randomised controlled trial. <i>BMC Pediatrics</i> , 2016, 16, 149.	1.7	21
51	Sodium pyrophosphate enhances iron bioavailability from bouillon cubes fortified with ferric pyrophosphate. <i>British Journal of Nutrition</i> , 2016, 116, 496-503.	2.3	27
52	Rural Beninese Children Are at Risk of Zinc Deficiency According to Stunting Prevalence and Plasma Zinc Concentration but Not Dietary Zinc Intakes. <i>Journal of Nutrition</i> , 2016, 146, 114-123.	2.9	24
53	In-home fortification with 2.5%mg iron as NaFeEDTA does not reduce anaemia but increases weight gain: a randomised controlled trial in Kenyan infants. <i>Maternal and Child Nutrition</i> , 2015, 11, 151-162.	3.0	22
54	Oral iron supplements increase hepcidin and decrease iron absorption from daily or twice-daily doses in iron-depleted young women. <i>Blood</i> , 2015, 126, 1981-1989.	1.4	372

#	ARTICLE	IF	CITATIONS
55	Efficacy of highly bioavailable zinc from fortified water: a randomized controlled trial in rural Beninese children. <i>American Journal of Clinical Nutrition</i> , 2015, 102, 1238-1248.	4.7	12
56	Iron fortification adversely affects the gut microbiome, increases pathogen abundance and induces intestinal inflammation in Kenyan infants. <i>Gut</i> , 2015, 64, 731-742.	12.1	477
57	The Donation Interval of 56 Days Requires Extension to 180 Days for Whole Blood Donors to Recover from Disturbances in Iron Homeostasis. <i>Blood</i> , 2015, 126, 774-774.	1.4	0
58	Bioavailability of iron, zinc, folic acid, and vitamin A from fortified maize. <i>Annals of the New York Academy of Sciences</i> , 2014, 1312, 54-65.	3.8	42
59	Three-month B vitamin supplementation in pre-school children affects folate status and homocysteine, but not cognitive performance. <i>European Journal of Nutrition</i> , 2014, 53, 1445-1456.	3.9	8
60	Relevance of dietary iron intake and bioavailability in the management of HFE hemochromatosis: a systematic review. <i>American Journal of Clinical Nutrition</i> , 2013, 98, 468-479.	4.7	29
61	Iron Status and Systemic Inflammation, but Not Gut Inflammation, Strongly Predict Gender-Specific Concentrations of Serum Hepcidin in Infants in Rural Kenya. <i>PLoS ONE</i> , 2013, 8, e57513.	2.5	47
62	Dephytinisation with Intrinsic Wheat Phytase and Iron Fortification Significantly Increase Iron Absorption from Fonio (<i>Digitaria exilis</i>) Meals in West African Women. <i>PLoS ONE</i> , 2013, 8, e70613.	2.5	22
63	Folate Catabolites in Spot Urine as Non-Invasive Biomarkers of Folate Status during Habitual Intake and Folic Acid Supplementation. <i>PLoS ONE</i> , 2013, 8, e56194.	2.5	12
64	Maize Porridge Enriched with a Micronutrient Powder Containing Low-Dose Iron as NaFeEDTA but Not Amaranth Grain Flour Reduces Anemia and Iron Deficiency in Kenyan Preschool Children. <i>Journal of Nutrition</i> , 2012, 142, 1756-1763.	2.9	36
65	Phytic Acid-to-Iron Molar Ratio Rather than Polyphenol Concentration Determines Iron Bioavailability in Whole-Cowpea Meal among Young Women. <i>Journal of Nutrition</i> , 2012, 142, 1950-1955.	2.9	25
66	Whole Cowpea Meal Fortified with NaFeEDTA Reduces Iron Deficiency among Ghanaian School Children in a Malaria Endemic Area. <i>Journal of Nutrition</i> , 2012, 142, 1836-1842.	2.9	27
67	Dietary intake of eicosapentaenoic acid (EPA) and docosahexaenoic acid (DHA) in children – a workshop report. <i>British Journal of Nutrition</i> , 2010, 103, 923-928.	2.3	29
68	Iron status and food matrix strongly affect the relative bioavailability of ferric pyrophosphate in humans. <i>American Journal of Clinical Nutrition</i> , 2006, 83, 632-638.	4.7	112
69	Extruded rice fortified with micronized ground ferric pyrophosphate reduces iron deficiency in Indian schoolchildren: a double-blind randomized controlled trial. <i>American Journal of Clinical Nutrition</i> , 2006, 84, 822-829.	4.7	132
70	Iron Fortification Reduces Blood Lead Levels in Children in Bangalore, India. <i>Pediatrics</i> , 2006, 117, 2014-2021.	2.1	77
71	Iron status and food matrix strongly affect the relative bioavailability of ferric pyrophosphate in humans. <i>FASEB Journal</i> , 2006, 20, A625.	0.5	1
72	Iron fortification reduces blood lead levels in children: a randomized, double-blind, controlled trial in Bangalore, India. <i>FASEB Journal</i> , 2006, 20, A131.	0.5	2

#	ARTICLE	IF	CITATIONS
73	Development and Evaluation of Iron-fortified Extruded Rice Grains. <i>Journal of Food Science</i> , 2005, 70, S330.	3.1	65
74	Particle Size Reduction and Encapsulation Affect the Bioavailability of Ferric Pyrophosphate in Rats. <i>Journal of Nutrition</i> , 2004, 134, 3301-3304.	2.9	71
75	Introduction of Iodized Salt to Severely Iodine-Deficient Children Does Not Provoke Thyroid Autoimmunity: A One-Year Prospective Trial in Northern Morocco. <i>Thyroid</i> , 2003, 13, 199-203.	4.5	63
76	Development of a dried whole-blood spot thyroglobulin assay and its evaluation as an indicator of thyroid status in goitrous children receiving iodized salt. <i>American Journal of Clinical Nutrition</i> , 2003, 77, 1453-1458.	4.7	76