

# Ricardo Plancarte-Sanchez

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/5945450/publications.pdf>

Version: 2024-02-01

20  
papers

682  
citations

1040056

9  
h-index

794594

19  
g-index

22  
all docs

22  
docs citations

22  
times ranked

470  
citing authors

#	ARTICLE	IF	CITATIONS
1	Breakthrough pain characteristics and syndromes in patients with cancer pain. An international survey. <i>Palliative Medicine</i> , 2004, 18, 177-183.	3.1	282
2	Superior Hypogastric Plexus Block for Pelvic Cancer Pain. <i>Anesthesiology</i> , 1990, 73, 236-239.	2.5	234
3	Complex Regional Pain Syndrome Type 2 (Causalgia) After Automated Laser Discectomy. <i>Spine</i> , 1997, 22, 459-461.	2.0	24
4	Intractable upper body pain in a pediatric patient relieved with cervical epidural opioid administration. <i>Journal of Pain and Symptom Management</i> , 1991, 6, 98-99.	1.2	20
5	Superior hypogastric plexus block and ganglion impar (Walther). <i>Techniques in Regional Anesthesia and Pain Management</i> , 2005, 9, 86-90.	0.2	18
6	Femoroplasty: A New Option for Femur Metastasis. <i>Pain Practice</i> , 2013, 13, 409-415.	1.9	14
7	A prospective, open-label, multicentre study of pregabalin in the treatment of neuropathic pain in Latin America. <i>International Journal of Clinical Practice</i> , 2010, 64, 1301-1309.	1.7	11
8	Anterior Cervical Approach for Stellate Ganglion and T2 to T3 Sympathetic Blocks: A Novel Technique. <i>Pain Practice</i> , 2005, 5, 244-248.	1.9	9
9	Antiallodynic Activity of Ceftriaxone and Clavulanic Acid in Acute Administration is Associated with Serum TNF- $\alpha$ Modulation and Activation of Dopaminergic and Opioidergic Systems. <i>Drug Development Research</i> , 2017, 78, 105-115.	2.9	9
10	Repeated administration of mazindol reduces spontaneous pain-related behaviors without modifying bone density and microarchitecture in a mouse model of complete Freund's adjuvant-induced knee arthritis. <i>Journal of Pain Research</i> , 2017, Volume 10, 1777-1786.	2.0	8
11	Ceftriaxone and clavulanic acid induce antiallodynia and anti-inflammatory effects in rats using the carrageenan model. <i>Journal of Pain Research</i> , 2018, Volume 11, 977-985.	2.0	8
12	Interventional treatment of cancer pain. <i>Seminars in Pain Medicine</i> , 2003, 1, 34-42.	0.4	7
13	Sympathetic block: Thoracic and lumbar. <i>Techniques in Regional Anesthesia and Pain Management</i> , 2005, 9, 91-96.	0.2	7
14	Palliative regional anesthesia for hospitalized patients. <i>Current Opinion in Anaesthesiology</i> , 1994, 7, 444-447.	2.0	4
15	RE: Yamada K, Ishihara Y, Saito T: Relief of intractable perineal pain by coccygeal nerve block in anterior sacrococcygeal ligament after surgery for rectal cancer. <i>J Anesthesia</i> (1994)8:52-54. <i>Journal of Anesthesia</i> , 1997, 11, 170-171.	1.7	3
16	Complex regional pain syndrome: A review. <i>Cirurgiã Y Cirujanos (English Edition)</i> , 2017, 85, 366-374.	0.0	3
17	Neurolytic Techniques for Cancer Pain Management. , 2006, , 501-507.		1
18	Regarding "Percutaneous Augmented Peripheral Osteoplasty in Long Bones of Oncologic Patients for Pain Reduction and Prevention of Impeding Pathologic Fracture: The Rebar Concept" <i>CardioVascular and Interventional Radiology</i> , 2016, 39, 477-478.	2.0	1

#	ARTICLE	IF	CITATIONS
19	Letter to the editor: Other ganglion impar blocks are also safe. <i>Techniques in Regional Anesthesia and Pain Management</i> , 2008, 12, 209-210.	0.2	0
20	Comments on splanchnic nerve neurolysis via the transdiscal approach under fluoroscopic guidance: a retrospective study. <i>Korean Journal of Pain</i> , 2022, 35, 356-357.	2.2	0