

Ricardo Plancarte-Sanchez

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/5945450/publications.pdf>

Version: 2024-02-01

20
papers

682
citations

1040056

9
h-index

794594

19
g-index

22
all docs

22
docs citations

22
times ranked

470
citing authors

#	ARTICLE	IF	CITATIONS
1	Comments on splanchnic nerve neurolysis via the transdiscal approach under fluoroscopic guidance: a retrospective study. Korean Journal of Pain, 2022, 35, 356-357.	2.2	0
2	Ceftriaxone and clavulanic acid induce antiallodynia and anti-inflammatory effects in rats using the carrageenan model. Journal of Pain Research, 2018, Volume 11, 977-985.	2.0	8
3	Antiallodynic Activity of Ceftriaxone and Clavulanic Acid in Acute Administration is Associated with Serum TNF- α Modulation and Activation of Dopaminergic and Opioidergic Systems. Drug Development Research, 2017, 78, 105-115.	2.9	9
4	Complex regional pain syndrome: A review. Cirug�a Y Cirujanos (English Edition), 2017, 85, 366-374.	0.0	3
5	Repeated administration of mazindol reduces spontaneous pain-related behaviors without modifying bone density and microarchitecture in a mouse model of complete Freund's adjuvant-induced knee arthritis. Journal of Pain Research, 2017, Volume 10, 1777-1786.	2.0	8
6	Regarding Percutaneous Augmented Peripheral Osteoplasty in Long Bones of Oncologic Patients for Pain Reduction and Prevention of Impeding Pathologic Fracture: The Rebar Concept. Cardiovascular and Interventional Radiology, 2016, 39, 477-478.	2.0	1
7	Femoroplasty: A New Option for Femur Metastasis. Pain Practice, 2013, 13, 409-415.	1.9	14
8	A prospective, open-label, multicentre study of pregabalin in the treatment of neuropathic pain in Latin America. International Journal of Clinical Practice, 2010, 64, 1301-1309.	1.7	11
9	Letter to the editor: Other ganglion impar blocks are also safe. Techniques in Regional Anesthesia and Pain Management, 2008, 12, 209-210.	0.2	0
10	Neurolytic Techniques for Cancer Pain Management. , 2006, , 501-507.		1
11	Anterior Cervical Approach for Stellate Ganglion and T2 to T3 Sympathetic Blocks: A Novel Technique. Pain Practice, 2005, 5, 244-248.	1.9	9
12	Superior hypogastric plexus block and ganglion impar (Walther). Techniques in Regional Anesthesia and Pain Management, 2005, 9, 86-90.	0.2	18
13	Sympathetic block: Thoracic and lumbar. Techniques in Regional Anesthesia and Pain Management, 2005, 9, 91-96.	0.2	7
14	Breakthrough pain characteristics and syndromes in patients with cancer pain. An international survey. Palliative Medicine, 2004, 18, 177-183.	3.1	282
15	Interventional treatment of cancer pain. Seminars in Pain Medicine, 2003, 1, 34-42.	0.4	7
16	Complex Regional Pain Syndrome Type 2 (Causalgia) After Automated Laser Discectomy. Spine, 1997, 22, 459-461.	2.0	24
17	RE: Yamada K, Ishihara Y, Saito T: Relief of intractable perineal pain by coccygeal nerve block in anterior sacrococcygeal ligament after surgery for rectal cancer. J Anesthesia (1994)8:52-54. Journal of Anesthesia, 1997, 11, 170-171.	1.7	3
18	Palliative regional anesthesia for hospitalized patients. Current Opinion in Anaesthesiology, 1994, 7, 444-447.	2.0	4

#	ARTICLE	IF	CITATIONS
19	Intractable upper body pain in a pediatric patient relieved with cervical epidural opioid administration. <i>Journal of Pain and Symptom Management</i> , 1991, 6, 98-99.	1.2	20
20	Superior Hypogastric Plexus Block for Pelvic Cancer Pain. <i>Anesthesiology</i> , 1990, 73, 236-239.	2.5	234