Konstantina Kyritsi

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/5948405/publications.pdf

Version: 2024-02-01

		840776	940533
18	368	11	16
papers	citations	h-index	g-index
18	18	18	297
all docs	docs citations	times ranked	citing authors

#	Article	IF	CITATIONS
1	Kupffer Cells. American Journal of Pathology, 2020, 190, 2185-2193.	3.8	80
2	Knockout of secretin receptor reduces biliary damage and liver fibrosis in Mdr2â^'/â^' mice by diminishing senescence of cholangiocytes. Laboratory Investigation, 2018, 98, 1449-1464.	3.7	41
3	Mast Cells Regulate Ductular Reaction and Intestinal Inflammation in Cholestasis Through Farnesoid X Receptor Signaling. Hepatology, 2021, 74, 2684-2698.	7.3	35
4	The Secretin/Secretin Receptor Axis Modulates Ductular Reaction and Liver Fibrosis through Changes in Transforming Growth Factor-β1–Mediated Biliary Senescence. American Journal of Pathology, 2018, 188, 2264-2280.	3.8	31
5	Downregulation of hepatic stem cell factor by Vivo-Morpholino treatment inhibits mast cell migration and decreases biliary damage/senescence and liver fibrosis in Mdr2â°'/â^' mice. Biochimica Et Biophysica Acta - Molecular Basis of Disease, 2019, 1865, 165557.	3.8	25
6	Mast Cells Promote Nonalcoholic Fatty Liver Disease Phenotypes and Microvesicular Steatosis in Mice Fed a Western Diet. Hepatology, 2021, 74, 164-182.	7.3	25
7	Modulation of the Tryptophan Hydroxylase 1/Monoamine Oxidaseâ€A/5â€Hydroxytryptamine/5â€Hydroxytryptamine Receptor 2A/2B/2C Axis Regulates Biliary Proliferation and Liver Fibrosis During Cholestasis. Hepatology, 2020, 71, 990-1008.	7.3	23
8	Functional Role of the Secretin/Secretin Receptor Signaling During Cholestatic Liver Injury. Hepatology, 2020, 72, 2219-2227.	7.3	18
9	Biliary damage and liver fibrosis are ameliorated in a novel mouse model lacking l-histidine decarboxylase/histamine signaling. Laboratory Investigation, 2020, 100, 837-848.	3.7	18
10	The Functional Roles of Immune Cells in Primary Liver Cancer. American Journal of Pathology, 2022, 192, 826-836.	3.8	17
11	Inhibition of Secretin/Secretin Receptor Axis Ameliorates NAFLD Phenotypes. Hepatology, 2021, 74, 1845-1863.	7.3	16
12	Amelioration of Large Bile Duct Damage by Histamine-2 Receptor Vivo-Morpholino Treatment. American Journal of Pathology, 2020, 190, 1018-1029.	3.8	13
13	Knockout of the Tachykinin Receptor 1 in the Mdr2â^//â^ (Abcb4â^//â^) Mouse Model of Primary Sclerosing Cholangitis Reduces Biliary Damage and Liver Fibrosis. American Journal of Pathology, 2020, 190, 2251-2266.	3.8	9
14	Melatonin receptor 1A, but not 1B, knockout decreases biliary damage and liver fibrosis during cholestatic liver injury. Hepatology, 2022, 75, 797-813.	7.3	9
15	The Effects of Taurocholic Acid on Biliary Damage and Liver Fibrosis Are Mediated by Calcitonin-Gene-Related Peptide Signaling. Cells, 2022, 11, 1591.	4.1	6
16	FGF1 Signaling Modulates Biliary Injury and Liver Fibrosis in the Mdr2â ⁻ '/â ⁻ ' Mouse Model of Primary Sclerosing Cholangitis. Hepatology Communications, 2022, 6, 1574-1588.	4.3	2
17	The protective effects of estrogen on biliary and liver damage are independent of ERâ€Î² signaling in female Mdr2 ^{â€/â€} mice. FASEB Journal, 2022, 36, .	0.5	0
18	Mast Cells Contribute to Hepatic Neurokinin1 Receptor Signaling, Subsequent Biliary Damage and Peribiliary Fibrosis Via TGFâ€Î²1 Signaling in MDR2â€Ĵ―Mouse Model of Primary Scelrosing Cholangitis. FASEB Journal, 2022, 36, .	0.5	0