

Konstantina Kyritsi

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/5948405/publications.pdf>

Version: 2024-02-01

18
papers

368
citations

840776

11
h-index

940533

16
g-index

18
all docs

18
docs citations

18
times ranked

297
citing authors

#	ARTICLE	IF	CITATIONS
1	Melatonin receptor 1A, but not 1B, knockout decreases biliary damage and liver fibrosis during cholestatic liver injury. <i>Hepatology</i> , 2022, 75, 797-813.	7.3	9
2	FGF1 Signaling Modulates Biliary Injury and Liver Fibrosis in the Mdr2 ^{-/-} Mouse Model of Primary Sclerosing Cholangitis. <i>Hepatology Communications</i> , 2022, 6, 1574-1588.	4.3	2
3	The Functional Roles of Immune Cells in Primary Liver Cancer. <i>American Journal of Pathology</i> , 2022, 192, 826-836.	3.8	17
4	The protective effects of estrogen on biliary and liver damage are independent of ER α signaling in female Mdr2 ^{-/-} mice. <i>FASEB Journal</i> , 2022, 36, .	0.5	0
5	Mast Cells Contribute to Hepatic Neurokinin1 Receptor Signaling, Subsequent Biliary Damage and Peribiliary Fibrosis Via TGF β 1 Signaling in MDR2 ^{-/-} Mouse Model of Primary Sclerosing Cholangitis. <i>FASEB Journal</i> , 2022, 36, .	0.5	0
6	The Effects of Taurocholic Acid on Biliary Damage and Liver Fibrosis Are Mediated by Calcitonin-Gene-Related Peptide Signaling. <i>Cells</i> , 2022, 11, 1591.	4.1	6
7	Mast Cells Promote Nonalcoholic Fatty Liver Disease Phenotypes and Microvesicular Steatosis in Mice Fed a Western Diet. <i>Hepatology</i> , 2021, 74, 164-182.	7.3	25
8	Inhibition of Secretin/Secretin Receptor Axis Ameliorates NAFLD Phenotypes. <i>Hepatology</i> , 2021, 74, 1845-1863.	7.3	16
9	Mast Cells Regulate Ductular Reaction and Intestinal Inflammation in Cholestasis Through Farnesoid X Receptor Signaling. <i>Hepatology</i> , 2021, 74, 2684-2698.	7.3	35
10	Modulation of the Tryptophan Hydroxylase 1/Monoamine Oxidase ^A /5 ^H HT ^A Hydroxytryptamine/5 ^H HT ^A Receptor 2A/2B/2C Axis Regulates Biliary Proliferation and Liver Fibrosis During Cholestasis. <i>Hepatology</i> , 2020, 71, 990-1008.	7.3	23
11	Knockout of the Tachykinin Receptor 1 in the Mdr2 ^{-/-} (Abcb4 ^{-/-}) Mouse Model of Primary Sclerosing Cholangitis Reduces Biliary Damage and Liver Fibrosis. <i>American Journal of Pathology</i> , 2020, 190, 2251-2266.	3.8	9
12	Functional Role of the Secretin/Secretin Receptor Signaling During Cholestatic Liver Injury. <i>Hepatology</i> , 2020, 72, 2219-2227.	7.3	18
13	Kupffer Cells. <i>American Journal of Pathology</i> , 2020, 190, 2185-2193.	3.8	80
14	Amelioration of Large Bile Duct Damage by Histamine-2 Receptor Vivo-Morpholino Treatment. <i>American Journal of Pathology</i> , 2020, 190, 1018-1029.	3.8	13
15	Biliary damage and liver fibrosis are ameliorated in a novel mouse model lacking l-histidine decarboxylase/histamine signaling. <i>Laboratory Investigation</i> , 2020, 100, 837-848.	3.7	18
16	Downregulation of hepatic stem cell factor by Vivo-Morpholino treatment inhibits mast cell migration and decreases biliary damage/senescence and liver fibrosis in Mdr2 ^{-/-} mice. <i>Biochimica Et Biophysica Acta - Molecular Basis of Disease</i> , 2019, 1865, 165557.	3.8	25
17	The Secretin/Secretin Receptor Axis Modulates Ductular Reaction and Liver Fibrosis through Changes in Transforming Growth Factor- β 1-Mediated Biliary Senescence. <i>American Journal of Pathology</i> , 2018, 188, 2264-2280.	3.8	31
18	Knockout of secretin receptor reduces biliary damage and liver fibrosis in Mdr2 ^{-/-} mice by diminishing senescence of cholangiocytes. <i>Laboratory Investigation</i> , 2018, 98, 1449-1464.	3.7	41