

# Karen R. Reed

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/5968791/publications.pdf>

Version: 2024-02-01

28  
papers

2,454  
citations

394421

19  
h-index

526287

27  
g-index

30  
all docs

30  
docs citations

30  
times ranked

3982  
citing authors

#	ARTICLE	IF	CITATIONS
1	NAP1L1: A Novel Human Colorectal Cancer Biomarker Derived From Animal Models of Apc Inactivation. <i>Frontiers in Oncology</i> , 2020, 10, 1565.	2.8	17
2	APC2 is critical for ovarian WNT signalling control, fertility and tumour suppression. <i>BMC Cancer</i> , 2019, 19, 677.	2.6	21
3	Spatiotemporal regulation of liver development by the Wnt/ $\beta$ -catenin pathway. <i>Scientific Reports</i> , 2018, 8, 2735.	3.3	20
4	Subtle Deregulation of the Wnt-Signaling Pathway Through Loss of Apc2 Reduces the Fitness of Intestinal Stem Cells. <i>Stem Cells</i> , 2018, 36, 114-122.	3.2	13
5	Functional redundancy between Apc and Apc2 regulates tissue homeostasis and prevents tumorigenesis in murine mammary epithelium. <i>Oncogene</i> , 2017, 36, 1793-1803.	5.9	25
6	Organoids as a Model for Colorectal Cancer. <i>Current Colorectal Cancer Reports</i> , 2016, 12, 281-287.	0.5	28
7	Secreted HMGB1 from Wnt activated intestinal cells is required to maintain a crypt progenitor phenotype. <i>Oncotarget</i> , 2016, 7, 51665-51673.	1.8	8
8	E-cadherin can limit the transforming properties of activating $\beta$ -catenin mutations. <i>EMBO Journal</i> , 2015, 34, 2321-2333.	7.8	83
9	Hunk/Mak-v is a negative regulator of intestinal cell proliferation. <i>BMC Cancer</i> , 2015, 15, 110.	2.6	15
10	Proteomic profiling of a mouse model of acute intestinal Apc deletion leads to identification of potential novel biomarkers of human colorectal cancer (CRC). <i>Biochemical and Biophysical Research Communications</i> , 2013, 440, 364-370.	2.1	30
11	Entopic overexpression of <i>Ascl2</i> does not accelerate tumorigenesis in Apc <sup>Min</sup> mice. <i>Gut</i> , 2012, 61, 1435-1438.	12.1	18
12	OC-018...Validation of two APC-dependent potential biomarkers of colorectal carcinogenesis. <i>Gut</i> , 2012, 61, A8.2-A8.	12.1	0
13	Conditional Disruption of Axin1 Leads to Development of Liver Tumors in Mice. <i>Gastroenterology</i> , 2012, 143, 1650-1659.	1.3	45
14	Rectal epithelial cell mitosis and expression of macrophage migration inhibitory factor are increased 3 years after Roux-en-Y gastric bypass (RYGB) for morbid obesity: implications for long-term neoplastic risk following RYGB. <i>Gut</i> , 2011, 60, 893-901.	12.1	42
15	Cyclin D2...Cyclin-Dependent Kinase 4/6 Is Required for Efficient Proliferation and Tumorigenesis following Apc Loss. <i>Cancer Research</i> , 2010, 70, 8149-8158.	0.9	79
16	Liver Zonation Occurs Through a $\beta$ -Catenin-Dependent, c-Myc-Independent Mechanism. <i>Gastroenterology</i> , 2009, 136, 2316-2324.e3.	1.3	142
17	A limited role for p53 in modulating the immediate phenotype of Apc loss in the intestine. <i>BMC Cancer</i> , 2008, 8, 162.	2.6	26
18	Deficiency of Mbd2 Attenuates Wnt Signaling. <i>Molecular and Cellular Biology</i> , 2008, 28, 6094-6103.	2.3	43

#	ARTICLE	IF	CITATIONS
19	B-catenin deficiency, but not Myc deletion, suppresses the immediate phenotypes of APC loss in the liver. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2008, 105, 18919-18923.	7.1	66
20	Myc deletion rescues Apc deficiency in the small intestine. <i>Nature</i> , 2007, 446, 676-679.	27.8	530
21	PPAR $\delta$ status and mismatch repair mediated neoplasia in the mouse intestine. <i>BMC Cancer</i> , 2006, 6, 113.	2.6	8
22	Apc deficiency predisposes to renal carcinoma in the mouse. <i>Oncogene</i> , 2005, 24, 8205-8210.	5.9	64
23	Cyclin D1 Is Not an Immediate Target of $\beta$ -Catenin following Apc Loss in the Intestine. <i>Journal of Biological Chemistry</i> , 2005, 280, 28463-28467.	3.4	92
24	Loss of Apc in vivo immediately perturbs Wnt signaling, differentiation, and migration. <i>Genes and Development</i> , 2004, 18, 1385-1390.	5.9	700
25	PPAR $\delta$ status and Apc-mediated tumourigenesis in the mouse intestine. <i>Oncogene</i> , 2004, 23, 8992-8996.	5.9	105
26	Paternal imprints can be established on the maternal Igf2-H19 locus without altering replication timing of DNA. <i>Human Molecular Genetics</i> , 2003, 12, 3123-3132.	2.9	19
27	Sequence conservation and variability of imprinting in the Beckwith-Wiedemann syndrome gene cluster in human and mouse. <i>Human Molecular Genetics</i> , 2000, 9, 1829-1841.	2.9	118
28	Syntenic Organization of the Mouse Distal Chromosome 7 Imprinting Cluster and the Beckwith-Wiedemann Syndrome Region in Chromosome 11p15.5. <i>Human Molecular Genetics</i> , 1998, 7, 1149-1159.	2.9	97