## Max Birchwood

## List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/5973085/publications.pdf

Version: 2024-02-01

295 papers 19,253 citations

67 h-index 130 g-index

337 all docs

337 docs citations

times ranked

337

10049 citing authors

#	Article	IF	Citations
1	The Psychosis High-Risk State. JAMA Psychiatry, 2013, 70, 107.	11.0	1,222
2	The Social Functioning Scale the Development and Validation of a New Scale of Social Adjustment for use in Family Intervention Programmes with Schizophrenic Patients. British Journal of Psychiatry, 1990, 157, 853-859.	2.8	957
3	The Omnipotence of Voices. British Journal of Psychiatry, 1994, 164, 190-201.	2.8	733
4	Early intervention in psychosis. British Journal of Psychiatry, 1998, 172, 53-59.	2.8	631
5	Prediction of Psychosis in Adolescents and Young Adults at High Risk. Archives of General Psychiatry, 2010, 67, 241.	12.3	575
6	A selfâ€report Insight Scale for psychosis: reliability, validity and sensitivity to change. Acta Psychiatrica Scandinavica, 1994, 89, 62-67.	4.5	568
7	The power and omnipotence of voices: subordination and entrapment by voices and significant others. Psychological Medicine, 2000, 30, 337-344.	4.5	437
8	Cognitive Therapy and Recovery from Acute Psychosis: a Controlled Trial. British Journal of Psychiatry, 1996, 169, 593-601.	2.8	407
9	Designing youth mental health services for the 21st century: examples from Australia, Ireland and the UK. British Journal of Psychiatry, 2013, 202, s30-s35.	2.8	387
10	Depression, demoralization and control over psychotic illness: a comparison of depressed and non-depressed patients with a chronic psychosis. Psychological Medicine, 1993, 23, 387-395.	4.5	326
11	Predicting relapse in schizophrenia: the development and implementation of an early signs monitoring system using patients and families as observers, a preliminary investigation. Psychological Medicine, 1989, 19, 649-656.	4.5	317
12	†Theory of mind' skills during an acute episode of psychosis and following recovery. Psychological Medicine, 1998, 28, 1101-1112.	4.5	299
13	The revised Beliefs About Voices Questionnaire (BAVQ–R). British Journal of Psychiatry, 2000, 177, 229-232.	2.8	269
14	Early detection and intervention evaluation for people at risk of psychosis: multisite randomised controlled trial. BMJ, The, 2012, 344, e2233-e2233.	6.0	266
15	Pathways to emotional dysfunction in first-episode psychosis. British Journal of Psychiatry, 2003, 182, 373-375.	2.8	256
16	The omnipotence of voices: testing the validity of a cognitive model. Psychological Medicine, 1997, 27, 1345-1353.	4.5	244
17	Cognitive approach to depression and suicidal thinking in psychosis. British Journal of Psychiatry, 2000, 177, 516-521.	2.8	242
18	Cognitive therapy for command hallucinations: randomised controlled trial. British Journal of Psychiatry, 2004, 184, 312-320.	2.8	237

#	Article	IF	CITATIONS
19	Psychotic-like experiences and correlation with distress and depressive symptoms in a community sample of adolescents and young adults. Schizophrenia Research, 2010, 119, 258-265.	2.0	235
20	Depression and Schizophrenia: Cause, Consequence or Trans-diagnostic Issue?. Schizophrenia Bulletin, 2017, 43, sbw097.	4.3	221
21	Interpersonal and role-related schema influence the relationship with the dominant †voice†in schizophrenia: a comparison of three models. Psychological Medicine, 2004, 34, 1571-1580.	4.5	215
22	Cognitive approach to depression and suicidal thinking in psychosis. British Journal of Psychiatry, 2000, 177, 522-528.	2.8	191
23	Social anxiety and the shame of psychosis: A study in first episode psychosis. Behaviour Research and Therapy, 2007, 45, 1025-1037.	3.1	190
24	A new scale (SES) to measure engagement with community mental health services. Journal of Mental Health, 2002, 11, 191-198.	1.9	187
25	Early intervention in psychosis. The critical period hypothesis. The British Journal of Psychiatry Supplement, 1998, 172, 53-9.	0.1	184
26	Cognitive Therapy and Recovery from Acute Psychosis: a Controlled Trial. British Journal of Psychiatry, 1996, 169, 602-607.	2.8	175
27	Predicting engagement with services for psychosis: insight, symptoms and recovery style. British Journal of Psychiatry, 2003, 182, 123-128.	2.8	171
28	Specific and Non-specific Effects of Educational Intervention with Families Living with a Schizophrenic Relative. British Journal of Psychiatry, 1987, 150, 645-652.	2.8	169
29	Cognitive therapy and recovery from acute psychosis: A controlled trial. British Journal of Psychiatry, 2000, 177, 8-14.	2.8	165
30	Early intervention in schizophrenia. British Journal of Psychiatry, 1997, 170, 2-5.	2.8	164
31	The Omnipotence of Voices II: the Beliefs About Voices Questionnaire (BAVQ). British Journal of Psychiatry, 1995, 166, 773-776.	2.8	161
32	The evolution of depression and suicidality in first episode psychosis. Acta Psychiatrica Scandinavica, 2010, 122, 211-218.	4.5	151
33	An exploration of evolved mental mechanisms for dominant and subordinate behaviour in relation to auditory hallucinations in schizophrenia and critical thoughts in depression. Psychological Medicine, 2001, 31, 1117-1127.	4.5	150
34	Clinical outcomes of Joint Crisis Plans to reduce compulsory treatment for people with psychosis: a randomised controlled trial. Lancet, The, 2013, 381, 1634-1641.	13.7	148
35	The future of cognitive-behavioural therapy for psychosis: not a quasi-neuroleptic. British Journal of Psychiatry, 2006, 188, 107-108.	2.8	144
36	Psychological pathways to depression in schizophrenia. European Archives of Psychiatry and Clinical Neuroscience, 2005, 255, 202-212.	3.2	143

#	Article	IF	Citations
37	Schizophrenia: early warning signs. Advances in Psychiatric Treatment, 2000, 6, 93-101.	0.5	141
38	Adapting to the challenge of psychosis: personal resilience and the use of sealing-over (avoidant) coping strategies. British Journal of Psychiatry, 2004, 185, 410-415.	2.8	140
39	Early Intervention in Schizophrenia. Australian and New Zealand Journal of Psychiatry, 1993, 27, 374-378.	2.3	136
40	Reducing duration of untreated psychosis: care pathways to early intervention in psychosis services. British Journal of Psychiatry, 2013, 203, 58-64.	2.8	135
41	From Environment to Therapy in Psychosis: A Real-World Momentary Assessment Approach. Schizophrenia Bulletin, 2011, 37, 244-247.	4.3	134
42	Cognitive behaviour therapy to prevent harmful compliance with command hallucinations (COMMAND): a randomised controlled trial. Lancet Psychiatry, the, 2014, 1, 23-33.	7.4	120
43	Families coping with schizophrenia: coping styles, their origins and correlates. Psychological Medicine, 1990, 20, 857-865.	4.5	113
44	Social anxiety disorder in first-episode psychosis: incidence, phenomenology and relationship with paranoia. British Journal of Psychiatry, 2009, 195, 234-241.	2.8	113
45	Disability in people clinically at high risk of psychosis. British Journal of Psychiatry, 2010, 197, 278-284.	2.8	113
46	The European Prediction of Psychosis Study (EPOS): integrating early recognition and intervention in Europe. World Psychiatry, 2005, 4, 161-7.	10.4	108
47	The Influence of Ethnicity and Family Structure on Relapse in First-Episode Schizophrenia. British Journal of Psychiatry, 1992, 161, 783-790.	2.8	102
48	Relatives and Patients as Partners in the Management of Schizophrenia. British Journal of Psychiatry, 1990, 156, 654-660.	2.8	101
49	Loss, humiliation and entrapment as appraisals of schizophrenic illness: A prospective study of depressed and nonâ€depressed patients. British Journal of Clinical Psychology, 1998, 37, 259-268.	3.5	101
50	Specific and Non-specific Effects of Educational Intervention for Families Living with Schizophrenia. British Journal of Psychiatry, 1992, 160, 806-814.	2.8	100
51	Pathways to Violent Behavior During First-Episode Psychosis. JAMA Psychiatry, 2013, 70, 1287.	11.0	99
52	A systematic review and meta-regression analysis of aggression during the First Episode of Psychosis. Acta Psychiatrica Scandinavica, 2013, 128, 413-421.	4.5	96
53	Axis I diagnoses and transition to psychosis in clinical high-risk patients EPOS project: Prospective follow-up of 245 clinical high-risk outpatients in four countries. Schizophrenia Research, 2012, 138, 192-197.	2.0	94
54	Duration of untreated psychosis and pathways to care in first-episode psychosis. British Journal of Psychiatry, 2002, 181, s73-s77.	2.8	93

#	Article	IF	CITATIONS
55	Is early intervention for psychosis a waste of valuable resources?. British Journal of Psychiatry, 2003, 182, 196-198.	2.8	92
56	Early attachment experience and recovery from psychosis. British Journal of Clinical Psychology, 1998, 37, 269-284.	3.5	87
57	Efficacy and Safety of Pharmacological and Psychological Interventions for the Treatment of Psychosis and Schizophrenia in Children, Adolescents and Young Adults: A Systematic Review and Meta-Analysis. PLoS ONE, 2015, 10, e0117166.	2.5	86
58	Expressed emotion as an adaptation to loss. British Journal of Psychiatry, 2005, 187, s59-s64.	2.8	85
59	Whither the Attenuated Psychosis Syndrome?. Schizophrenia Bulletin, 2012, 38, 1130-1134.	4.3	85
60	Cannabis use and age at onset of symptoms in subjects at clinical high risk for psychosis. Acta Psychiatrica Scandinavica, 2012, 125, 45-53.	4.5	82
61	Impact of contact on adolescents' mental health literacy and stigma: the SchoolSpace cluster randomised controlled trial. BMJ Open, 2016, 6, e009435.	1.9	81
62	Cannabis Use Is Associated With Increased Psychotic Symptoms and Poorer Psychosocial Functioning in First-Episode Psychosis: A Report From the UK National EDEN Study. Schizophrenia Bulletin, 2016, 42, 619-625.	4.3	80
63	Views of Young People in Early Intervention Services for First-Episode Psychosis in England. Psychiatric Services, 2011, 62, 882-887.	2.0	79
64	Drug and alcohol problems amongst individuals with severe mental health problems in an inner city area of the UK. Social Psychiatry and Psychiatric Epidemiology, 2001, 36, 448-455.	3.1	76
65	EARLY INTERVENTION IN PSYCHOTIC RELAPSE. Clinical Psychology Review, 2001, 21, 1211-1226.	11.4	75
66	Early intervention in psychosis: keeping faith with evidence-based health care. Psychological Medicine, 2010, 40, 399-404.	4.5	74
67	Acting on command hallucinations: A cognitive approach. British Journal of Clinical Psychology, 1997, 36, 139-148.	3.5	73
68	Comparing time use in individuals at different stages of psychosis and a non-clinical comparison group. Schizophrenia Research, 2015, 161, 188-193.	2.0	72
69	Barriers to shared decision making in mental health care: qualitative study of the Joint Crisis Plan for psychosis. Health Expectations, 2016, 19, 448-458.	2.6	71
70	Investigating trajectories of social recovery in individuals with first-episode psychosis: A latent class growth analysis. British Journal of Psychiatry, 2015, 207, 536-543.	2.8	70
71	Improving psychological adjustment following a first episode of psychosis: A randomised controlled trial of cognitive therapy to reduce post psychotic trauma symptoms. Behaviour Research and Therapy, 2009, 47, 454-462.	3.1	69
72	Social recovery therapy in combination with early intervention services for enhancement of social recovery in patients with first-episode psychosis (SUPEREDEN3): a single-blind, randomised controlled trial. Lancet Psychiatry,the, 2018, 5, 41-50.	7.4	67

#	Article	IF	Citations
73	Management of first-episode psychosis. Advances in Psychiatric Treatment, 2001, 7, 133-140.	0.5	67
74	REDIRECT: cluster randomised controlled trial of GP training in first-episode psychosis. British Journal of General Practice, 2009, 59, e183-e190.	1.4	66
75	Effect of delaying treatment of first-episode psychosis on symptoms and social outcomes: a longitudinal analysis and modelling study. Lancet Psychiatry,the, 2020, 7, 602-610.	7.4	66
76	Early intervention in schizophrenia: Theoretical background and clinical strategies. British Journal of Clinical Psychology, 1992, 31, 257-278.	3.5	63
77	The subjective experience and phenomenology of depression following first episode psychosis: A qualitative study using photo-elicitation. Journal of Affective Disorders, 2013, 149, 166-174.	4.1	63
78	The Critical Period for Early Intervention. Psychiatric Rehabilitation Skills, 2000, 4, 182-198.	0.1	62
79	Evidence for a Cognitive Model of Auditory Hallucinations. Journal of Nervous and Mental Disease, 2003, 191, 542-545.	1.0	62
80	The Trauma of First Episode Psychosis: The Role of Cognitive Mediation. Australian and New Zealand Journal of Psychiatry, 2004, 38, 327-333.	2.3	62
81	The <scp>UK</scp> national evaluation of the development and impact of <scp>E</scp> arly <scp>I</scp> ntervention <scp>S</scp> ervices (the <scp>N</scp> ational <scp>EDEN</scp> studies): study rationale, design and baseline characteristics. Microbial Biotechnology, 2014, 8, 59-67.	1.7	62
82	Mental health services for young people: matching the service to the need. British Journal of Psychiatry, 2013, 202, s1-s2.	2.8	61
83	Cognitive Predictors of Social and Occupational Functioning in Early Psychosis: A Systematic Review and Meta-analysis of Cross-Sectional and Longitudinal Data. Schizophrenia Bulletin, 2021, 47, 1243-1253.	4.3	59
84	Early detection and intervention evaluation for people at high-risk of psychosis-2 (EDIE-2): trial rationale, design and baseline characteristics. Microbial Biotechnology, 2011, 5, 24-32.	1.7	58
85	Adolescents and young adults who are not in employment, education, or training. BMJ, The, 2013, 347, f5270-f5270.	6.0	57
86	The needs of high and low expressed emotion families: a normative approach. Social Psychiatry and Psychiatric Epidemiology, 1993, 28, 11-16.	3.1	55
87	Acting on command hallucinations and dangerous behavior: A critique of the major findings in the last decade. Clinical Psychology Review, 2004, 24, 513-528.	11.4	55
88	Early intervention in psychosis: Opportunities for secondary prevention. British Journal of Clinical Psychology, 1996, 35, 487-502.	3.5	53
89	Ethnicity and pathways to care during first episode psychosis: the role of cultural illness attributions. BMC Psychiatry, 2015, 15, 287.	2.6	52
90	Early Intervention in Psychotic Relapse: Cognitive Approaches to Detection and Management. Behaviour Change, 1995, 12, 2-19.	1.3	51

#	Article	IF	CITATIONS
91	Antipsychotic medication versus psychological intervention versus a combination of both in adolescents with first-episode psychosis (MAPS): a multicentre, three-arm, randomised controlled pilot and feasibility study. Lancet Psychiatry,the, 2020, 7, 788-800.	7.4	51
92	Insight, duration of untreated psychosis and attachment in first-episode psychosis: prospective study of psychiatric recovery over 12-month follow-up. British Journal of Psychiatry, 2014, 205, 60-67.	2.8	50
93	Social anxiety disorder and shame cognitions in psychosis. Psychological Medicine, 2013, 43, 133-142.	4.5	49
94	What service users with psychotic disorders want in a mental health crisis or relapse: thematic analysis of joint crisis plans. Social Psychiatry and Psychiatric Epidemiology, 2014, 49, 1609-1617.	3.1	49
95	The course of negative symptom in first episode psychosis and the relationship with social recovery. Schizophrenia Research, 2016, 174, 165-171.	2.0	49
96	Acting on voices: Omnipotence, sources of threat, and safetyâ€seeking behaviours. British Journal of Clinical Psychology, 2008, 47, 201-213.	3.5	48
97	Towards a new paradigm of care: the International Declaration on Youth Mental Health. Microbial Biotechnology, 2013, 7, 103-108.	1.7	48
98	A multi-element psychosocial intervention for early psychosis (GET UP PIANO TRIAL) conducted in a catchment area of 10 million inhabitants: study protocol for a pragmatic cluster randomized controlled trial. Trials, 2012, 13, 73.	1.6	47
99	Layers of listening: Qualitative analysis of the impact of early intervention services for first-episode psychosis on carers' experiences. British Journal of Psychiatry, 2015, 207, 135-142.	2.8	47
100	Transitioning careâ€leavers with mental health needs: †they set you up to fail!'. Child and Adolescent Mental Health, 2017, 22, 138-147.	3.5	47
101	Development and implementation of early intervention services for young people with psychosis: case study. British Journal of Psychiatry, 2009, 194, 446-450.	2.8	46
102	Pathways to emotional dysfunction in first-episode psychosis. British Journal of Psychiatry, 2003, 182, 373-5.	2.8	46
103	Ethnicity and detention: are Black and minority ethnic (BME) groups disproportionately detained under the Mental Health Act 2007?. Social Psychiatry and Psychiatric Epidemiology, 2016, 51, 703-711.	3.1	45
104	Early intervention in psychosis. International Clinical Psychopharmacology, 1998, 13, S31-S40.	1.7	41
105	Delays in mental health services and duration of untreated psychosis. Psychiatric Bulletin, 2007, 31, 408-410.	0.3	41
106	Prediction of psychosis in clinical high-risk patients by the Schizotypal Personality Questionnaire. Results of the EPOS project. European Psychiatry, 2013, 28, 469-475.	0.2	41
107	Internalized stigma, emotional dysfunction and unusual experiences in young people at risk of psychosis. Microbial Biotechnology, 2015, 9, 133-140.	1.7	41
108	Improving Therapeutic Relationships. Qualitative Health Research, 2015, 25, 1637-1647.	2.1	40

#	Article	IF	Citations
109	Service users' views of moving on from early intervention services for psychosis: a longitudinal qualitative study in primary care. British Journal of General Practice, 2012, 62, e183-e190.	1.4	39
110	Combined severe mental health and substance use problems: What are the training and support needs of staff working with this client group?. Journal of Mental Health, 2001, 10, 131-140.	1.9	38
111	Predicting compliance with command hallucinations: Anger, impulsivity and appraisals of voices' power and intent. Schizophrenia Research, 2013, 147, 163-168.	2.0	38
112	Impact of early intervention services on duration of untreated psychosis: Data from the National EDEN prospective cohort study. Schizophrenia Research, 2014, 159, 1-6.	2.0	38
113	Seeking help for firstâ€episode psychosis: a family narrative. Microbial Biotechnology, 2016, 10, 334-345.	1.7	37
114	Cognitive approach to depression and suicidal thinking in psychosis. 2. Testing the validity of a social ranking model. British Journal of Psychiatry, 2000, 177, 522-8.	2.8	37
115	Control of Auditory Hallucinations through Occlusion of Monaural Auditory Input. British Journal of Psychiatry, 1986, 149, 104-107.	2.8	34
116	Impact of cognitive therapy on internalised stigma in people with at-risk mental states. British Journal of Psychiatry, 2013, 203, 140-145.	2.8	34
117	Youth services: meeting the mental health needs of adolescents. Irish Journal of Psychological Medicine, 2015, 32, 13-19.	1.0	34
118	A prospective study of dysfunctional thinking and the regulation of negative intrusive memories in bipolar 1 disorder: implications for affect regulation theory. Psychological Medicine, 2007, 37, 689.	4.5	33
119	Development and validation of multivariable prediction models of remission, recovery, and quality of life outcomes in people with first episode psychosis: a machine learning approach. The Lancet Digital Health, 2019, 1, e261-e270.	12.3	33
120	Psychotic experience subtypes, poor mental health status and helpâ€seeking behaviour in a community sample of young adults. Microbial Biotechnology, 2012, 6, 300-308.	1.7	32
121	Attachment: Developmental pathways to affective dysregulation in young people at ultraâ€high risk of developing psychosis. British Journal of Clinical Psychology, 2013, 52, 424-437.	3.5	32
122	Psychosocial outcome in patients at clinical high risk of psychosis: a prospective follow-up. Social Psychiatry and Psychiatric Epidemiology, 2013, 48, 303-311.	3.1	31
123	Randomised Controlled Trial of Joint Crisis Plans to Reduce Compulsory Treatment for People with Psychosis: Economic Outcomes. PLoS ONE, 2013, 8, e74210.	2.5	31
124	Pathways to care in subjects at high risk for psychotic disorders — A European perspective. Schizophrenia Research, 2014, 152, 400-407.	2.0	31
125	Can the therapeutic relationship predict 18 month outcomes for individuals with psychosis?. Psychiatry Research, 2014, 220, 585-591.	3.3	31
126	Personal beliefs about illness questionnaireâ€revised (PBIQâ€R): Reliability and validation in a first episode sample. British Journal of Clinical Psychology, 2012, 51, 448-458.	3.5	30

#	Article	IF	Citations
127	Preventing the Entrenchment of High Expressed Emotion in First Episode Psychosis: Early Developmental Attachment Pathways. Australian and New Zealand Journal of Psychiatry, 2000, 34, S191-S197.	2.3	29
128	Coâ€designing a virtual world with young people to deliver social cognition therapy in early psychosis. Microbial Biotechnology, 2020, 14, 37-43.	1.7	29
129	The national policy reforms for mental health services and the story of early intervention services in the United Kingdom. Journal of Psychiatry and Neuroscience, 2005, 30, 362-5.	2.4	29
130	Design of the BiRmingham Early Detection In untREated psyChosis Trial (REDIRECT): cluster randomised controlled trial of general practitioner education in detection of first episode psychosis [ISRCTN87898421]. BMC Health Services Research, 2005, 5, 19.	2.2	28
131	A prospective study of PTSD following recovery from firstâ€episode psychosis: The threat from persecutors, voices, and patienthood. British Journal of Clinical Psychology, 2012, 51, 418-433.	3.5	28
132	Short-term functional outcome and premorbid adjustment in clinical high-risk patients. Results of the EPOS project. European Psychiatry, 2014, 29, 371-380.	0.2	28
133	Social anxiety in first-episode psychosis: The role of childhood trauma and adult attachment. Journal of Affective Disorders, 2014, 163, 102-109.	4.1	28
134	Stigma and access to care in firstâ€episode psychosis. Microbial Biotechnology, 2019, 13, 1208-1213.	1.7	28
135	The person in recovery from acute and severe psychosis: The role of dependency, self-criticism, and efficacy American Journal of Orthopsychiatry, 2004, 74, 480-488.	1.5	27
136	Autobiographical memory and post-psychotic depression in first episode psychosis. British Journal of Clinical Psychology, 2004, 43, 97-104.	3.5	27
137	A preliminary evaluation of integrated treatment for co-existing substance use and severe mental health problems: Impact on teams and service users. Journal of Mental Health, 2006, 15, 577-591.	1.9	27
138	CRIMSON [CRisis plan IMpact: Subjective and Objective coercion and eNgagement] Protocol: A randomised controlled trial of joint crisis plans to reduce compulsory treatment of people with psychosis. Trials, 2010, 11, 102.	1.6	27
139	Views of early psychosis clinicians on discontinuation of antipsychotic medication following symptom remission in first episode psychosis. Microbial Biotechnology, 2016, 10, 355-361.	1.7	27
140	Informing people with schizophrenia about their illness: The effect of residual symptoms. Journal of Mental Health, 1992, 1, 61-70.	1.9	26
141	Prevalence of treatment resistance and clozapine use in early intervention services. BJPsych Open, 2020, 6, e107.	0.7	26
142	The contribution of shame to postâ€psychotic trauma. British Journal of Clinical Psychology, 2013, 52, 162-182.	3.5	25
143	Joint crisis planning in mental health care: the challenge of implementation in randomized trials and in routine care. World Psychiatry, 2015, 14, 281-283.	10.4	25
144	Early Signs Monitoring to Prevent Relapse in Psychosis and Promote Well-Being, Engagement, and Recovery: Protocol for a Feasibility Cluster Randomized Controlled Trial Harnessing Mobile Phone Technology Blended With Peer Support. JMIR Research Protocols, 2020, 9, e15058.	1.0	24

#	Article	IF	CITATIONS
145	The Concept of Schizophrenia and Phase-Specific Treatment: Cognitive-Behavioral Treatment in Pre-Psychosis and in Nonresponders. Journal of the American Academy of Psychoanalysis and Dynamic Psychiatry, 2003, 31, 209-228.	0.3	23
146	Cognitive behavioural therapy group work with voice hearers. Part 1. British Journal of Nursing, 2007, 16, 248-252.	0.7	23
147	The Strauss and Carpenter Prognostic Scale in subjects clinically at high risk of psychosis. Acta Psychiatrica Scandinavica, 2013, 127, 53-61.	4.5	23
148	Negative cognition, affect, metacognition and dimensions of paranoia in people at ultra-high risk of psychosis: a multi-level modelling analysis. Psychological Medicine, 2015, 45, 2675-2684.	4.5	23
149	Don't turn your back on the symptoms of psychosis: the results of a proof-of-principle, quasi-experimental intervention to reduce duration of untreated psychosis. BMC Psychiatry, 2016, 16, 127.	2.6	23
150	The COMMAND trial of cognitive therapy to prevent harmful compliance with command hallucinations: predictors of outcome and mediators of change. Psychological Medicine, 2018, 48, 1966-1974.	4.5	23
151	The development and implementation of an educational intervention on first episode psychosis for primary care. Medical Education, 2005, 39, 1006-1014.	2.1	22
152	Pilot randomised trial of a brief intervention for comorbid substance misuse in psychiatric inâ€patient settings. Acta Psychiatrica Scandinavica, 2016, 133, 298-309.	4.5	22
153	Carers' and service users' experiences of early intervention in psychosis services: implications for care partnerships. Microbial Biotechnology, 2018, 12, 410-416.	1.7	22
154	A Feasibility and Acceptability Trial of Social Cognitive Therapy in Early Psychosis Delivered Through a Virtual World: The VEEP Study. Frontiers in Psychiatry, 2020, 11, 219.	2.6	22
155	Early Intervention and Sustaining the Management of Vulnerability. Australian and New Zealand Journal of Psychiatry, 2000, 34, S181-S184.	2.3	21
156	Cognitive behavioural therapy group work with voice hearers. Part 2. British Journal of Nursing, 2007, 16, 292-295.	0.7	21
157	A randomised controlled feasibility trial for an educational school-based mental health intervention: study protocol. BMC Psychiatry, 2012, 12, 23.	2.6	21
158	Metacognitive beliefs as psychological predictors of social functioning: An investigation with young people at risk of psychosis. Psychiatry Research, 2018, 262, 520-526.	3.3	21
159	Duration of untreated psychosis and clinical outcomes of first episode psychosis: An observational and an instrumental variables analysis. Microbial Biotechnology, 2019, 13, 841-847.	1.7	21
160	Command Hallucinations: Cognitive Theory, Therapy, and Research. Journal of Cognitive Psychotherapy, 2003, 17, 67-84.	0.4	20
161	Early Intervention in Psychosis. Disease Management and Health Outcomes, 2005, 13, 53-63.	0.4	20
162	Depression predicts persistence of paranoia in clinical high-risk patients to psychosis: results of the EPOS project. Social Psychiatry and Psychiatric Epidemiology, 2016, 51, 247-257.	3.1	20

#	Article	IF	Citations
163	Psychosocial Family Intervention in Schizophrenia: A Review of Empirical Studies. British Journal of Psychiatry, 1992, 160, 272-275.	2.8	19
164	Preservation and compensation: The functional neuroanatomy of insight and working memory in schizophrenia. Schizophrenia Research, 2014, 152, 201-209.	2.0	19
165	Barriers and facilitators to partnership working between Early Intervention Services and the voluntary and community sector. Health and Social Care in the Community, 2008, 16, 493-500.	1.6	18
166	A multi-centre, randomised controlled trial of cognitive therapy to prevent harmful compliance with command hallucinations. BMC Psychiatry, 2011, 11, 155.	2.6	18
167	Prevalence of Psychotic-like Experiences in Young Adults With Social Anxiety Disorder and Correlation With Affective Dysregulation. Journal of Nervous and Mental Disease, 2013, 201, 1053-1059.	1.0	18
168	Through the looking glass: self-reassuring meta-cognitive capacity and its relationship with the thematic content of voices. Frontiers in Human Neuroscience, 2013, 7, 213.	2.0	18
169	Core outcome sets for use in effectiveness trials involving people with bipolar and schizophrenia in a community-based setting (PARTNERS2): study protocol for the development of two core outcome sets. Trials, 2015, 16, 47.	1.6	18
170	Experiences and satisfaction of children, young people and their parents with alternative mental health models to inpatient settings: a systematic review. European Child and Adolescent Psychiatry, 2020, 29, 1621-1633.	4.7	18
171	Predictors of engagement in first-episode psychosis. Schizophrenia Research, 2016, 175, 204-208.	2.0	17
172	Expressed emotion as a predictor of the first psychotic episode â€" Results of the European prediction of psychosis study. Schizophrenia Research, 2018, 199, 346-352.	2.0	17
173	Refining a model of collaborative care for people with a diagnosis of bipolar, schizophrenia or other psychoses in England: a qualitative formative evaluation. BMC Psychiatry, 2019, 19, 7.	2.6	17
174	Urban remediation: a new recovery-oriented strategy to manage urban stress after first-episode psychosis. Social Psychiatry and Psychiatric Epidemiology, 2020, 55, 273-283.	3.1	17
175	The CIRCuiTS study (Implementation of cognitive remediation in early intervention services): protocol for a randomised controlled trial. Trials, 2018, 19, 183.	1.6	16
176	Vulnerability to psychosocial disability in psychosis. Epidemiology and Psychiatric Sciences, 2019, 28, 140-145.	3.9	16
177	Components of therapy as mechanisms of change in cognitive therapy for people at risk of psychosis: Analysis of the EDIE-2 trial. British Journal of Psychiatry, 2015, 207, 123-129.	2.8	15
178	Shortâ€term outcome of substanceâ€induced psychotic disorder in a large <scp>UK</scp> first episode psychosis cohort. Acta Psychiatrica Scandinavica, 2016, 134, 321-328.	4.5	15
179	Preventing the Entrenchment of High Expressed Emotion in First Episode Psychosis: Early Developmental Attachment Pathways. Australian and New Zealand Journal of Psychiatry, 2000, 34, A191-A197.	2.3	14
180	Power and Perceived Expressed Emotion of Voices: Their Impact on Depression and Suicidal Thinking in Those Who Hear Voices. Clinical Psychology and Psychotherapy, 2013, 20, 199-205.	2.7	14

#	Article	IF	CITATIONS
181	Individualisation in crisis planning for people with psychotic disorders. Epidemiology and Psychiatric Sciences, 2014, 23, 353-359.	3.9	14
182	Childhood adversity predicts persistence of suicidal thoughts differently in females and males at clinical highâ€risk patients of psychosis. Results of the EPOS project. Microbial Biotechnology, 2019, 13, 935-942.	1.7	14
183	Psychological and social treatments: course and outcome. Current Opinion in Psychiatry, 1999, 12, 61-66.	6.3	14
184	Specialist community teams backed by years of quality research. The Psychiatrist, 2013, 37, 38-38.	0.3	14
185	Structure and stability of symptoms in first episode psychosis: a longitudinal network approach. Translational Psychiatry, 2021, 11, 567.	4.8	14
186	Perceived negative attitude of others as an early sign of psychosis. European Psychiatry, 2009, 24, 233-238.	0.2	13
187	Early intervention in psychosis services: the next generation. Microbial Biotechnology, 2014, 8, 1-2.	1.7	13
188	Distress, omnipotence, and responsibility beliefs in command hallucinations. British Journal of Clinical Psychology, 2017, 56, 303-309.	3.5	13
189	After the storm, Solar comes out: A new service model for children and adolescent mental health. Microbial Biotechnology, 2021, 15, 731-738.	1.7	13
190	Early Intervention and Sustaining the Management of Vulnerability. Australian and New Zealand Journal of Psychiatry, 2000, 34, A181-A184.	2.3	12
191	Usefulness of EQ-5D for evaluation of health-related quality of life in young adults with first-episode psychosis. Quality of Life Research, 2013, 22, 1055-1063.	3.1	12
192	CBT for psychosis: not a â€~quasi-neuroleptic'. British Journal of Psychiatry, 2014, 204, 488-489.	2.8	12
193	Costâ€effectiveness of early intervention services for psychosis and fidelity to national policy implementation guidance. Microbial Biotechnology, 2018, 12, 747-756.	1.7	12
194	Controversies and Growing Points in Cognitive-Behavioural interventions for People with Schizophrenia. Behavioural and Cognitive Psychotherapy, 1992, 20, 305-342.	1.2	11
195	Perceived negative attitude of others predicts transition to psychosis in patients at risk of psychosis. European Psychiatry, 2012, 27, 264-266.	0.2	11
196	Adolescent construction of mental illness: implication for engagement and treatment. Microbial Biotechnology, 2018, 12, 626-636.	1.7	11
197	Study protocol for a randomised controlled trial of CBT vs antipsychotics vs both in 14–18-year-olds: Managing Adolescent first episode Psychosis: a feasibility study (MAPS). Trials, 2019, 20, 395.	1.6	11
198	A High-Fidelity Virtual Environment for the Study of Paranoia. Schizophrenia Research and Treatment, 2013, 2013, 1-7.	1.5	10

#	Article	IF	Citations
199	Cognitive Behavioural Relating Therapy (CBRT) for Voice Hearers: A Case Study. Behavioural and Cognitive Psychotherapy, 2013, 41, 626-631.	1.2	10
200	Systematic Review of Cognitive-Behavioural Therapy for Social Anxiety Disorder in Psychosis. Brain Sciences, 2017, 7, 45.	2.3	10
201	Derivation of a prediction model for a diagnosis of depression in young adults: a matched case–control study using electronic primary care records. Microbial Biotechnology, 2018, 12, 444-455.	1.7	10
202	Service innovations. Psychiatric Bulletin, 2003, 27, 183-186.	0.3	10
203	Psychotherapies for Schizophrenia: A Review. , 2001, , 147-241.		9
204	A pilot study to assess the feasibility and impact of a brief motivational intervention on problem drug and alcohol use in adult mental health inpatient units: study protocol for a randomized controlled trial. Trials, 2014, 15, 308.	1.6	9
205	Informed, advance refusals of treatment by people with severe mental illness in a randomised controlled trial of joint crisis plans: demand, content and correlates. BMC Psychiatry, 2017, 17, 376.	2.6	9
206	Predictors of social and role outcomes in first episode psychosis: <scp>A prospective</scp> 12â€month study of social cognition, neurocognition and symptoms. Microbial Biotechnology, 2021, 15, 993-1001.	1.7	9
207	Psychotherapies for Schizophrenia: A Review. , 2003, , 153-247.		8
208	Cognitive Therapy for Command Hallucinations: Not a Quasi-Neuroleptic. Journal of Contemporary Psychotherapy, 2006, 36, 1-7.	1.2	8
209	Muslim faith healers' views on substance misuse and psychosis. Mental Health, Religion and Culture, 2012, 15, 653-673.	0.9	8
210	Abuse and dysfunctional affiliations in childhood: an exploration of their impact on voice-hearers' appraisals of power and expressed emotion. Psychosis, 2012, 4, 19-31.	0.8	8
211	Psychopathology and affect dysregulation across the continuum of psychosis: a multiple comparison group study. Microbial Biotechnology, 2014, 8, 221-228.	1.7	8
212	The COMMAND trial of cognitive therapy for harmful compliance with command hallucinations (CTCH): a qualitative study of acceptability and tolerability in the UK. BMJ Open, 2018, 8, e021657.	1.9	8
213	Lived experiences of negative symptoms in firstâ€episode psychosis: A qualitative secondary analysis. Microbial Biotechnology, 2019, 13, 773-779.	1.7	8
214	Community stakeholders' perspectives on youth mental health in India: Problems, challenges and recommendations. Microbial Biotechnology, 2021, 15, 716-722.	1.7	8
215	Commentary on Garety & Freeman I: â€~Cognitive approaches to delusions - A critical review of theories and evidence'. British Journal of Clinical Psychology, 1999, 38, 315-318.	3.5	7
216	Evaluating substance misuse interventions in psychosis: The limitations of the RCT with 'patient' as the unit of analysis. Journal of Mental Health, 2001, 10, 585-587.	1.9	7

#	Article	IF	Citations
217	Preventing Relapse of Psychotic Illness. Disease Management and Health Outcomes, 2002, 10, 395-407.	0.4	7
218	Don't turn your back on the symptoms of psychosis: a proof-of-principle, quasi-experimental public health trial to reduce the duration of untreated psychosis in Birmingham, UK. BMC Psychiatry, 2013, 13, 67.	2.6	7
219	Expressed Emotion and First Episodes of Schizophrenia. British Journal of Psychiatry, 1987, 151, 859-860.	2.8	6
220	Exploring the attitudes of staff working within mental health settings toward clients who use cannabis. Drugs: Education, Prevention and Policy, 2009, 16, 311-327.	1.3	6
221	Collaborating to Provide Early-Intervention Services to Persons in England With First-Episode Psychosis. Psychiatric Services, 2009, 60, 1484-1488.	2.0	6
222	NICE v. SIGN on psychosis and schizophrenia: Same roots, similar guidelines, different interpretations. British Journal of Psychiatry, 2016, 208, 316-319.	2.8	6
223	Frequency and preventative interventions for non-suicidal self-injury and suicidal behaviour in primary school-age children: a scoping review protocol. BMJ Open, 2017, 7, e017291.	1.9	6
224	Warwick-India-Canada (WIC) global mental health group: rationale, design and protocol. BMJ Open, 2021, 11, e046362.	1.9	6
225	Appraisals: Voices' power and purpose. , 2010, , 81-102.		6
226	The trauma of first episode psychosis: the role of cognitive mediation. Australian and New Zealand Journal of Psychiatry, 2004, 38, 327-333.	2.3	6
227	Pathways to emotional dysfunction in first-episode psychosis. British Journal of Psychiatry, 2003, 182, 373-375.	2.8	6
228	Early intervention and sustaining the management of vulnerability. Australian and New Zealand Journal of Psychiatry, 2000, 34, \$181-\$184.	2.3	5
229	TC5D EDEN: EVALUATING THE DEVELOPMENT AND IMPACT OF EARLY INTERVENTION SERVICES IN THE WEST MIDLANDS. Schizophrenia Research, 2006, 86, S50.	2.0	5
230	Treating prolactinoma and psychosis: medication and cognitive behavioural therapy. BMJ Case Reports, 2011, 2011, bcr0720103185-bcr0720103185.	0.5	5
231	The efficacy of psychotherapy in reducing post-psychotic trauma. Epidemiology and Psychiatric Sciences, 2011, 20, 127-131.	3.9	5
232	Measuring adherence in social recovery therapy with people with first episode psychosis. Behavioural and Cognitive Psychotherapy, 2020, 48, 82-90.	1.2	5
233	Development and Validation of a Nonremission Risk Prediction Model in First-Episode Psychosis: An Analysis of 2 Longitudinal Studies. Schizophrenia Bulletin Open, 2021, 2, sgab041.	1.7	5
234	New directions in the psychosocial approach to the psychoses. Journal of Mental Health, 1998, 7, 111-114.	1.9	4

#	Article	IF	Citations
235	Preventing the entrenchment of high expressed emotion in first episode psychosis: early developmental attachment pathways. Australian and New Zealand Journal of Psychiatry, 2000, 34, S191-S197.	2.3	4
236	The impact of training within a comprehensive dual diagnosis strategy: the Combined Psychosis and Substance Use (COMPASS) experience. Mental Health and Substance Use: Dual Diagnosis, 2012, 5, 206-216.	0.5	4
237	Using the crisis., 1995,, 116-148.		4
238	FC4D IMPROVING PSYCHOLOGICAL ADJUSTMENT FOLLOWING A FIRST EPISODE OF PSYCHOSIS: A RANDOMISED CONTROLLED TRIAL OF COGNITIVE THERAPY TO REDUCE POST-PSYCHOTIC TRAUMA SYMPTOMS. Schizophrenia Research, 2006, 86, S39.	2.0	3
239	Emotional and personality dysfunctions in early psychosis. , 2009, , 283-302.		3
240	A Consultation-Liaison Service on Integrated Treatment: A Program Description. Journal of Dual Diagnosis, 2013, 9, 149-157.	1.2	3
241	Physical illnesses, developmental risk factors and psychiatric diagnoses among subjects at risk of psychosis. European Psychiatry, 2013, 28, 135-140.	0.2	3
242	Cannabis use and abstention in first-episode psychosis: the participants'view. Mental Health and Substance Use: Dual Diagnosis, 2013, 6, 47-58.	0.5	3
243	Recruiting South Asians into a UK Mental Health Randomised Controlled Trial: Experiences of Field Researchers. Journal of Racial and Ethnic Health Disparities, 2014, 1, 181-193.	3.2	3
244	The impact of emotional acknowledgement and self-esteem on resilience to family dysfunction. European Psychiatry, 2017, 41, S734-S735.	0.2	3
245	A Synthetic Literature Review on the Management of Emerging Treatment Resistance in First Episode Psychosis: Can We Move towards Precision Intervention and Individualised Care?. Medicina (Lithuania), 2020, 56, 638.	2.0	3
246	Improved social functioning following social recovery therapy in first episode psychosis: Do social cognition and neurocognition change following therapy, and do they predict treatment response?. Schizophrenia Research, 2021, 228, 249-255.	2.0	3
247	Evaluation of a primary care-based collaborative care model (PARTNERS2) for people with diagnoses of schizophrenia, bipolar, or other psychoses: study protocol for a cluster randomised controlled trial. BJGP Open, 2021, 5, BJGPO.2021.0033.	1.8	3
248	Predicting and controlling relapse in schizophrenia: early signs monitoring., 1992,, 293-308.		3
249	Cognitive Therapy for command hallucinations. , 2010, , 103-122.		3
250	Interactive highly realistic virtual reality as a tool for understanding the genesis and treatment of psychotic symptoms. Studies in Health Technology and Informatics, 2009, 144, 138-40.	0.3	3
251	A case of monosymptomatic musical hallucinations. Irish Journal of Psychological Medicine, 1993, 10, 89-90.	1.0	2
252	â€~Schizophrenia postdrome': a study of lowâ€level psychotic experience after remission of firstâ€episode schizophrenia. Microbial Biotechnology, 2009, 3, 296-299.	1.7	2

#	Article	lF	Citations
253	Cognitive Theory and Therapy for Command Hallucinations. Journal of Experimental Psychopathology, 2012, 3, 537-551.	0.8	2
254	Protocol for a systematic review and meta-analysis of cognitive-behavioural therapy for social anxiety disorder in psychosis. Systematic Reviews, 2014, 3, 62.	5.3	2
255	Early intervention and identification strategies for young people at risk of developing mental health issues: working in partnership with schools in <scp>B</scp> irmingham, <scp>UK</scp> . Microbial Biotechnology, 2017, 11, 471-479.	1.7	2
256	Mental Health Hospital Admissions: a Teachable Moment and Window of Opportunity To Promote Change in Drug and Alcohol Misuse. International Journal of Mental Health and Addiction, 2019, 17, 22-40.	7.4	2
257	Status of primary and secondary mental healthcare of people with severe mental illness: an epidemiological study from the UK PARTNERS2 programme. BJPsych Open, 2021, 7, e53.	0.7	2
258	The impact of manic symptoms in firstâ€episode psychosis: Findings from the UK National EDEN study. Acta Psychiatrica Scandinavica, 2021, 144, 358-367.	4.5	2
259	Congratulations to Mike Startup and colleagues (Startup et al. 2004) for their excellent study replicating and extending our initial trial applying CBT to promote recovery from acute psychosis (Drury et al. 1996a,b, 2001). Psychological Medicine, 2005, 35, 152-153.	4.5	1
260	Commentary on Tiffin et al.?s ?From commitment to reality ? early intervention in psychosis services in England?. Reasons to be cheerful. Microbial Biotechnology, 2007, 1, 109-109.	1.7	1
261	Staff attitude towards cocaine/crack-cocaine use amongst individuals with severe mental health problems in an inner city area of the UK. Mental Health and Substance Use: Dual Diagnosis, 2008, 1, 205-215.	0.5	1
262	EARLY DETECTION AND INTERVENTION EVALUATION FOR PEOPLE AT-RISK OF PSYCHOSIS. Schizophrenia Research, 2012, 136, S17.	2.0	1
263	Poster #172 A MULTI-CENTRE RANDOMISED CONTROLLED TRIAL OF COGNITIVE THERAPY TO PREVENT HARMFUL COMPLIANCE WITH COMMAND HALLUCINATIONS. Schizophrenia Research, 2012, 136, S153.	2.0	1
264	G493â€How can health services effectively improve the mental health of young people leaving public care? the lync study. Archives of Disease in Childhood, 2015, 100, A212.1-A212.	1.9	1
265	Knitting up the ravelled sleeve of care: sleep and psychosis. Lancet Psychiatry, the, 2015, 2, 950-951.	7.4	1
266	The association between treatment beliefs and engagement in care in first episode psychosis. Schizophrenia Research, 2019, 204, 409-410.	2.0	1
267	Service innovations. Psychiatric Bulletin, 2003, 27, 183-186.	0.3	1
268	Understanding the Role of Emotion in Psychosis: Social Anxiety Disorder in First-Episode Psychosis. , 2011, , 89-110.		1
269	EPOS — General outlines of the "European prediction of psychosis study― Schizophrenia Research, 2003, 60, 5.	2.0	0
270	FK3 EMOTIONAL DYSFUNCTION IN PSYCHOSIS: A LOST CAUSE?. Schizophrenia Research, 2006, 86, S3.	2.0	O

#	Article	IF	Citations
271	FC7D EDEN: ADAPTING EIS IN THE WEST MIDLANDS TO THE SOCIO-CULTURAL CONTEXT. Schizophrenia Research, 2006, 86, S42.	2.0	O
272	0178 QUALITY OF LIFE AND RISK OF PSYCHOSIS. RESULTS OF THE EPOS PROJECT IN TURKU CENTRE. Schizophrenia Research, 2006, 86, S87.	2.0	0
273	0234 ANOMALOUS INTERPERSONAL SCHEMATA AMONG PEOPLE WITH FIRST-EPISODE PSYCHOSIS AND CO-MORBID SOCIAL ANXIETY. Schizophrenia Research, 2006, 86, S105.	2.0	O
274	0501 DOES ETHNICITY STILL INFLUENCE TREATMENT DELAY AND CARE PATHWAYS ONCE EARLY INTERVENTION IS ESTABLISHED?. Schizophrenia Research, 2006, 86, S125-S126.	2.0	0
275	0366 SCOTTISH STUDY OF PSYCHOSIS: ENGAGEMENT, ATTACHMENT AND RECOVERY (SSP: EAR)- STUDY OUTLINE. Schizophrenia Research, 2006, 86, S142-S143.	2.0	0
276	0435 EXPRESSED EMOTION AND PSYCHOPATHOLOGY IN SUBJECTS AT RISK FOR PSYCHOSIS. Schizophrenia Research, 2006, 86, S152.	2.0	0
277	0458 EXPRESSED EMOTION IN CLIENTS AT HIGH-RISK OF DEVELOPING PSYCHOSIS –RESULTS FROM THE EPOS PROJECT. Schizophrenia Research, 2006, 86, S152.	2.0	O
278	0176 NATIONAL EDEN: SOCIO-CULTURAL ISSUES IN IMPLEMENTING EIS ACROSS FIVE DIVERSE SITES IN ENGLAND. Schizophrenia Research, 2006, 86, S164.	2.0	0
279	EPOS - sample characteristics, transition rates and psychopathological predictors. European Psychiatry, 2007, 22, S64.	0.2	O
280	Premorbid adjustment in persons at high risk for psychosis. European Psychiatry, 2007, 22, S64.	0.2	0
281	Subjective quality of life and its changes in patients at risk of psychosis. European Psychiatry, 2008, 23, S68.	0.2	O
282	Transition to psychosis: Neuropsychological test results of the epos study. European Psychiatry, 2008, 23, S68-S69.	0.2	0
283	P03-118 - Perceived Negative Attitude Associates with Psychotic Experiences. European Psychiatry, 2010, 25, .	0.2	0
284	Treatment in the at-risk mental state for psychosis - a comparison of four european countries. European Psychiatry, 2011, 26, 2040-2040.	0.2	0
285	Subjective quality of life and its changes in patients at-risk for psychosis results of the EPOS study. European Psychiatry, 2011, 26, 2039-2039.	0.2	0
286	Course of psychopathology in the at-risk mental state - outcomes beyond transition to psychosis. European Psychiatry, 2011, 26, 2037-2037.	0.2	0
287	Poster #191 THE STRAUSS AND CARPENTER PROGNOSTIC SCALE IN SUBJECTS CLINICALLY AT HIGH RISK OF PSYCHOSIS. Schizophrenia Research, 2012, 136, S254.	2.0	O
288	THE PSYCHOSIS HIGH RISK STATE: IS IT VALID?. Schizophrenia Research, 2014, 153, S23-S24.	2.0	0

#	Article	IF	CITATIONS
289	Causal Connection Between Depression and Paranoia. European Psychiatry, 2015, 30, 113.	0.2	O
290	Substance misuse brief interventions during psychiatric hospital admissions. Advances in Dual Diagnosis, 2016, 9, .	0.5	0
291	The impact of the transition from primary school to secondary school on young adolescents. European Psychiatry, 2017, 41, S179-S180.	0.2	0
292	Prognostic models in first-episode psychosis – Authors' reply. The Lancet Digital Health, 2020, 2, e61.	12.3	0
293	A novel application of the Lego® Serious Play® methodology in mental health research: Understanding service users' experiences of the O‶9 mental health model in the United Kingdom. Microbial Biotechnology, 2021, , .	1.7	0
294	O-90â€Randomised controlled trial of joint crisis plans to reduce compulsory treatment for people with psychosis: Clinical outcomes and implementation. BMJ Supportive and Palliative Care, 2015, 5, A28.3-A28.	1.6	0
295	Treatment Targets in the Pre-psychotic Phase. , 0, , 75-91.		0