
seyed Akbar khodaparast

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/5975147/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	Discovery of a cryptic species, Erysiphe salicina sp. nov., and reconstruction of the phylogeny of powdery mildews on Populus and Salix spp Mycological Progress, 2022, 21, 1.	1.4	2
2	Studies on the secondary metabolism of Rosellinia and Dematophora strains (Xylariaceae) from Iran. Mycological Progress, 2022, 21, .	1.4	5
3	Resolution of the Hypoxylon fuscum Complex (Hypoxylaceae, Xylariales) and Discovery and Biological Characterization of Two of Its Prominent Secondary Metabolites. Journal of Fungi (Basel,) Tj ETQq1 1 0.78431	4 rg₿ ₮. \$Ove	rlo ck10 Tf 50
4	Phylogeny and taxonomy of the Erysiphe adunca complex (Erysiphaceae, Helotiales) on poplars and willows. Mycological Progress, 2021, 20, 517-537.	1.4	6
5	Phylogenetic and Functional Diversity of Saprolegniales and Fungi Isolated from Temperate Lakes in Northeast Germany. Journal of Fungi (Basel, Switzerland), 2021, 7, 968.	3.5	5
6	First report of powdery mildew caused by Golovinomyces bolayi on okra (Abelmoschus esculentus). Australasian Plant Disease Notes, 2020, 15, 1.	0.7	0
7	Phylogenetic structure of the Iranian capnodiaceous sooty mould fungi inferred from the sequences of rDNA regions and TEF1-a. Mycological Progress, 2020, 19, 155-169.	1.4	2
8	Taxonomical and functional diversity of Saprolegniales in Anzali lagoon, Iran. Aquatic Ecology, 2020, 54, 323-336.	1.5	16
9	Discovery of a new species of the Hypoxylon rubiginosum complex from Iran and antagonistic activities of Hypoxylon spp. against the Ash Dieback pathogen, Hymenoscyphus fraxineus, in dual culture. MycoKeys, 2020, 66, 105-133.	1.9	17
10	Association Analysis of Charcoal Rot Disease Resistance in Soybean. Plant Pathology Journal, 2019, 35, 189-199.	1.7	8
11	The contrasting roles of aquatic fungi and oomycetes in the degradation and transformation of polymeric organic matter. Limnology and Oceanography, 2019, 64, 2662-2678.	3.1	18
12	Diversity and pathogenicity of Botryosphaeriaceae species on forest trees in the north of Iran. European Journal of Forest Research, 2019, 138, 685-704.	2.5	19
13	A survey on Peniophora (Russulales, Basidiomycota) species in Iran. Nova Hedwigia, 2018, 107, 257-270.	0.4	1
14	Two new species of Pseudopyricularia from Iran. Mycological Progress, 2017, 16, 729-736.	1.4	2
15	A multiplex <scp>PCR</scp> â€based technique for identification of <i>Biscogniauxia mediterranea</i> and <i>Obolarina persica</i> causing charcoal disease of oak trees in Zagros forests. Forest Pathology, 2017, 47, e12330.	1.1	10
16	Declineâ€associated <i>Phaeoacremonium</i> species occurring on forest trees in the north of Iran. Forest Pathology, 2017, 47, e12368.	1.1	14
17	Notes on the genus Leveillula (Erysiphaceae): a new unrecorded species and notes on Leveillula infecting Ficus, Cucurbita and Tropaeolum in Iran. Phytotaxa, 2016, 260, 267.	0.3	4
18	Taxonomy and phylogenetic position of Phyllactinia takamatsui, a newly described powdery mildew on cotoneaster, based on molecular and morphological data. Mycological Progress, 2016, 15, 1.	1.4	1

#	Article	IF	CITATIONS
19	Rice grain discoloration effect on physical properties and head rice yield in three rice cultivars. Quality Assurance and Safety of Crops and Foods, 2016, 8, 283-288.	3.4	2
20	New records of polypores from Iran, with a checklist of polypores for Gilan Province Czech Mycology, 2016, 68, 139-148.	0.5	5
21	Evaluation of the virulence of <i>Sclerotium rolfsii</i> isolates on <i>Arachis hypogaea</i> and screening for resistant genotypes in greenhouse conditions. Hellenic Plant Protection Journal, 2015, 8, 1-11.	0.4	5
22	New records of cercosporoid hyphomycetes from Iran. Mycotaxon, 2012, 120, 157-169.	0.3	13
23	Additional rDNA ITS sequences and its phylogenetic consequences for the genus Leveillula with emphasis on conidium morphology. Mycological Progress, 2012, 11, 741-752.	1.4	21
24	Species pattern and phylogenetic relationships ofTrichodermastrains in rice fields of Southern Caspian Sea, Iran. Cereal Research Communications, 2011, 39, 560-568.	1.6	16
25	Dominant variance has an important role in downy mildew resistance in cucumber. Horticulture Environment and Biotechnology, 2011, 52, 422-426.	2.1	3
26	First record of powdery mildew of castor- oil plant (Ricinus communis) caused by the anamorphic stage of Leveillula taurica in Iran. Australasian Plant Disease Notes, 2011, 6, 36-38.	0.7	3
27	Distribution and severity of damage by <i>Cryphonectria parasitica</i> in the chestnut stands in Guilan province, Iran. Forest Pathology, 2010, 40, 450-457.	1.1	3
28	<i>Malcolmia africana</i> , a new host for powdery mildew disease caused by <i>Erysiphe cruciferarum</i> in Iran. Australasian Plant Disease Notes, 2010, 5, 101.	0.7	3
29	Parental Line Selection for Cucumber Hybrid Seed Production by Principal Components Analysis. International Journal of Vegetable Science, 2010, 16, 316-325.	1.3	2
30	Species, host range, and geographical distribution of powdery mildew fungi in Iran. Mycotaxon, 2009, 108, 213-216.	0.3	17
31	Comprehensive molecular phylogenetic analysis and evolution of the genus Phyllactinia (Ascomycota:) Tj ETQq1 I	0.78431 2.5	4 rgBT /Over
32	Molecular and morphological characterization of Leveillula (Ascomycota: Erysiphales) on monocotyledonous plants. Mycological Research, 2007, 111, 673-679.	2.5	15
33	Multilocus phylogenetic analyses within Blumeria graminis, a powdery mildew fungus of cereals. Molecular Phylogenetics and Evolution, 2007, 44, 741-751.	2.7	75
34	Occurrence of Cryphonectria parasitica the causal agent of chestnut blight in Iran. Plant Pathology, 2006, 55, 815-815.	2.4	4
35	Three new species of the genus Leveillula from Iran. Mycoscience, 2002, 43, 459-461.	0.8	8
36	Phylogenetic structure of the genus Leveillula (Erysiphales: Erysiphaceae) inferred from the nucleotide sequences of the rDNA ITS region with special reference to the L. taurica species complex. Mycological Research, 2001, 105, 909-918.	2.5	51